

ON THE CIRCULAR COUÉTTE FLOW

BY

AKIRA TAKESHITA

1. Introduction.

In the present paper we shall study properties of special solutions of the 3-dimensional stationary and non-stationary Navier-Stokes equations from several points of view. The main solution considered in this paper is the classical Couétté flow between two rotating concentric cylinders.

There are mainly two reasons why we wish to treat this flow. First it is important to study precise properties of special solutions of the Navier-Stokes equations<sup>(1)</sup> in special cases along with the mathematical analysis for the equation in the general formulation, since we still not have complete theory for the problem of existence or non-existence of global in time regular solutions of the 3-dimensional N-S equations. Secondly, the Couétté flow itself has many properties which are quite interesting mathematically as well as physically. For example, as the celebrated experiment by G. I. Taylor in 1923 revealed and as was rigorously proved mathematically by W. Velte [6] in 1966, the Couétté flow is not necessarily the unique solution. And moreover, what is more interesting and seems even a peculiar phenomenon to mathematicians

---

(1) For simplicity, we call it an N-S equation.

is the experimental fact that in experiments the Couéte flow is actually observed in one circumstance and in another it is not, although mathematically it is a solution for both cases. In the latter case another flow different from the Couéte flow is observed. The explanations tried by physicists for this phenomenon are done from the stand-point of the stability theory (See, for example, C. C. Lin [3]). But at the present state of the mathematical treatment of the N-S equations where we do not know whether unique and regular global in time solution of the 3-dimensional N-S equation really exists or not, the stability theory is confronted with theoretical difficulties.

In this paper we shall treat problems related to the Couéte flows in general cases. First we shall study the problem whether or not for any given  $T (> 0)$  there exists a regular solution in the interval  $[0, T]$  of the corresponding non-stationary N-S equation for every initial data given near the Couéte flow. After establishing an affirmative answer to this question, we next prove the differentiability in the sense of Fréchet of the evolution operator which gives a mathematical foundation to the linear stability theory. Thirdly, we shall discuss the eigenvalue problem for the Fréchet derivative of the evolution operator at the Couéte flow and show that the Couéte flow is unstable for infinitesimal perturbations under certain circumstances. Finally we shall prove that the Couéte flow is an isolated solution under almost all circumstances.

The author wishes to express hearty thanks to Prof. H. Fujita who showed interests in this work and encouraged him with stimulating conversations.

## 2. Formulation of the problem and the results.

We consider the non-stationary and stationary N-S equations in a domain  $G$  between two concentric cylinders of radii  $R_1$  and  $R_2 (> R_1)$ . More precisely  $G$  is defined by  $G = \{x = (x_1, x_2, x_3) \in \mathbb{R}^3 ; R_1^2 < x_1^2 + x_2^2 < R_2^2\}$ . The two cylinders rotate with constant angular velocities ; the inner with  $\Omega_1$  and the outer with  $\Omega_2$  counter clockwise.

In  $G$  the N-S equation is expressed for the non-stationary motion

$$(NSE) \left\{ \begin{array}{l} \frac{\partial v}{\partial t} = \Delta v - \nabla_v v - \text{grad } q, \quad t > 0, x \in G \\ \text{div } v(x) = 0, \quad x \in G \\ v(0, x) = a(x) \\ \text{and the boundary condition of adherence at the boundary that} \\ \text{the fluid on the boundary move with the boundary.} \end{array} \right.$$

and for the stationary motion

$$(SE) \left\{ \begin{array}{l} \Delta v - \nabla_v v - \text{grad } q = 0 \\ \text{div } v = 0 \\ \text{the boundary condition of adherence at the boundary.} \end{array} \right.$$

The boundary condition will be given explicitly later. In these equations  $v$  is the velocity vector field of the fluid in question and  $q$  is a scalar function which is the pressure in the fluid and the unknowns are  $v$  and  $q$ .  $\nabla$  is the canonical affine connection in  $\mathbb{R}^3$  and  $\Delta$  is the

Laplacian. In the sequel to treat the equations (NSE) and (SE) effectively, we adopt two coordinate systems, the cartesian coordinate system  $(x_1, x_2, x_3)$  and the cylindrical coordinate system  $(r, \phi, z)$ . In these coordinate systems a vector field  $v$  is expressed ; in the

first,  $v = (v_1, v_2, v_3) = \sum_{i=1}^3 v_i \frac{\partial}{\partial x_i}$  and in the latter,  $v = (v_r, v_\phi, v_z)$   
 $= v_r \frac{\partial}{\partial r} + \frac{v_\phi}{r} \frac{\partial}{\partial \phi} + v_z \frac{\partial}{\partial z}$ . The well-known Couéte flow is thus expressed

by  $W = (A + \frac{B}{r^2}) \frac{\partial}{\partial \phi}$  and  $q_0 = \int^r \frac{1}{\rho} (A\rho + \frac{B}{\rho}) d\rho$  where

$A = (R_2^2 \Omega_2 - R_1^2 \Omega_1) / (R_2^2 - R_1^2)$  and  $B = R_1^2 R_2^2 (\Omega_1 - \Omega_2) / (R_2^2 - R_1^2)$ . This is a solution of (SE) for all  $R_1, R_2, \Omega_1, \Omega_2$ . We preserve the letter  $w$  to denote the Couéte flow exclusively in this paper. It is the aim of this paper to study the properties of the Couéte flow and those of the solutions of the equations (NSE) and (SE) near the Couéte flow.

To that end, we consider a portion  $G_h$  of  $G$  and treat the equations in  $G_h$ .  $G_h$  is defined by  $G_h = \{x \in G ; 0 < x_3 < h\}$  and the union of the lower and the upper bottom is denoted by  $\delta G_h$ . In  $G_h$  we consider the following initial value problem and boundary value problem.

$$(IVP) \left\{ \begin{array}{l} \frac{\partial v}{\partial t} = \Delta v - \nabla_v v - \text{grad } q \\ \text{div } v = 0 \\ v(x_1, x_2, 0) = v(x_1, x_2, h), \quad \frac{\partial v}{\partial x_3}(x_1, x_2, 0) = \frac{\partial v}{\partial x_3}(x_1, x_2, h) \\ v = (A + \frac{B}{R_i^2}) \frac{\partial}{\partial \phi} \quad \text{for } r^2 = x_1^2 + x_2^2 = R_i^2, \quad i = 1, 2. \end{array} \right.$$

and

$$(BVP) \left\{ \begin{array}{l} \Delta v - \Delta_v v - \text{grad } q = 0 \\ \text{div } v = 0 \\ v(x_1, x_2, 0) = v(x_1, x_2, h), \quad \frac{\partial v}{\partial x_3}(x_1, x_2, 0) = \frac{\partial v}{\partial x_3}(x_1, x_2, h). \\ v = (A + \frac{B}{R_i^2}) \frac{\partial}{\partial \phi} \quad \text{for } r = R_i, \quad i = 1, 2. \end{array} \right.$$

In order to treat the problem in a functional analysis setting, we introduce some function spaces and operators.  $L^2 = L^2(G_h)$  is a Hilbert space of all  $R^3$ -valued functions  $v = (v_1(x), v_2(x), v_3(x))$  defined in

$G_h$  for which the norm  $\|v\| = \left\{ \int_{G_h} \sum_{i=1}^3 v_i^2(x) dx \right\}^{1/2}$  is finite.

$C_{0,\sigma}^\infty \equiv C_{0,\sigma}^\infty(G_h)$  is a space of all  $R^3$ -valued functions

$\varphi = (\varphi_1(x), \varphi_2(x), \varphi_3(x))$  such that (i) every component  $\varphi_j \in C^\infty(\overline{G_h})$

(ii)  $\varphi_j \equiv 0$  near  $G_h \equiv \{x ; x_1^2 + x_2^2 = R_i^2, \quad i = 1, 2, \quad 0 \leq x_3 \leq h\}$

(iii)  $\text{div } \varphi = 0$ . (iv)  $\varphi(x_1, x_2, 0) = \varphi(x_1, x_2, h)$ ,

$\frac{\partial}{\partial x_3}(x_1, x_2, 0) = \frac{\partial}{\partial x_3}(x_1, x_2, h)$ .  $L_\sigma^2 \equiv L_\sigma^2(G_h)$  is the completion of

$C_{0,\sigma}^\infty$  with respect to the norm of  $L^2(G_h)$ . By  $P$  we denote the

orthogonal projection of  $L^2$  onto  $L_\sigma^2$ . For  $\varphi \in C_{0,\sigma}^\infty$  we define an

operator  $A$  by  $A\varphi = -P\Delta\varphi$ . It is easy to verify that  $A$  is a strictly

positive symmetric operator in the Hilbert space  $L_\sigma^2$ . The positivity is

verified by the Poincaré inequality. We take the Friedrichs extension of

$A$  which we denote also by the same letter  $A$ . Then  $A$  is a strictly

## 6

positive self-adjoint operator with compact inverse  $A^{-1}$ . For real  $\gamma$  we denote  $A^\gamma$  the fractional power of  $A$  and by  $\mathcal{D}(A^\gamma)$  the domain of definition of  $A^\gamma$  endowed with its graph norm  $\|\varphi\|_\gamma = \|A^\gamma \varphi\|$ . Transforming the unknowns from  $(v, q)$  to  $(\tilde{v}, p)$  by the identities  $v = u + w$ ,  $q = p + q_0$ , and making use of the above notations, the equations (IVP) and (BVP) are transformed (formally) into the following abstract evolution equation (EE) and an operator equation (E) in  $L_\sigma^2$ , respectively.

$$(EE) \quad \begin{cases} \frac{du}{dt} = -Au - P(\nabla_u u + \nabla_w u + \nabla_u w) \\ u(0) = a \end{cases}$$

and

$$(E) \quad Au + P(\Delta u + \nabla_w u + \nabla_u w) = 0.$$

In (EE)  $u = u(t)$  is regarded as an  $L_\sigma^2$ -valued function defined on  $\{t \geq 0\}$ . In order to investigate the integrability of the equation (EE), we introduce the following integral equation (IE),

$$(IE) \quad u(t) = e^{-tA} a - \int_0^t e^{-(t-s)A} P(\nabla_{u(s)} u(s) + \nabla_w u(s) + \nabla_{u(s)} w) ds$$

where by  $e^{-tA}$  we denote the semi-group of operators generated by  $-A$ .

If we can prove the existence of solution  $u(t)$  of (IE) with certain regularity property, it is easy to verify that it is a regular solution of (EE). Hence we shall be engaged exclusively in (IE).

Now we can state our theorems which we are going to prove in this paper.

Theorem 1 (an existence theorem)

(1) For every  $r > 0$  there exists  $T > 0$  such that there exists uniquely a solution of (IE) on the interval  $[0, T]$  for every  $a \in (A^{1/2})$  with  $\|A^{1/2}a\| < r$ .

(2) For every  $T > 0$  there exists  $r > 0$  such that the statement in (1) holds.

Theorem 2 (differentiability of the evolution operator)

The evolution operator  $S_t : \mathcal{D}(A^{1/2}) \rightarrow \mathcal{D}(A^{1/2})$  is Fréchet differentiable at  $0 \in \mathcal{D}(A^{1/2})$  for any  $t > 0$ .

Theorem 3 (eigenvalue problem for the Fréchet derivative of the evolution operator)

For any  $\omega = (\omega_1, \omega_2) \in S^1$  (the 1-sphere) such that  $(R_2^2 \omega_2^2 - R_1^2 \omega_1^2)(\omega_2 - \omega_1) > 0$  there exists  $\rho = \rho_\omega > 0$  such that the Fréchet derivative of the evolution operator  $S_t$  at 0 has real positive eigenvalue greater than 1 for every  $t > 0$ .

Theorem 4 0 is an isolated solution of (E) in  $\mathcal{D}(A^\gamma)$  with  $\gamma > \frac{3}{4}$  for almost all  $\Omega = (\Omega_1, \Omega_2) \in \mathbb{R}^2$ .

## 3. Existence theorems.

First we state two lemmas concerning the operators  $A$  and  $e^{-tA}$ .

For the statement of Lemma 1, we introduce the operator  $B$  defined as follows. The domain of definition of  $B$  is  $\mathcal{D}(B) = W_2^2(G_h) \cap H_2^1(G_h)$

where  $W_2^m(G_h)$  is a  $L^2$ -Sobolev space of order  $m$ , and  $H_2^2(G_h) =$

$\{u \in W_2^1(G_h) ; u|_{G_h} = 0, u(x_1, x_2, 0) = u(x_1, x_2, h), \frac{\partial u}{\partial x_3}(x_1, x_2, 0) =$

$\frac{\partial u}{\partial x_3}(x_1, x_2, h)\}$ . And for  $u \in \mathcal{D}(B)$ ,  $Bu = -\Delta u$ .

Lemma 1. For  $0 < \gamma < 1$ ,  $\mathcal{D}(A^\gamma) = \mathcal{D}(B^\gamma) \cap L_\sigma^2$ . And therefore  $\mathcal{D}(A^\gamma) \subset C(\bar{G}_h)$  for  $\gamma > \frac{3}{4}$  and  $\sup_{x \in G_h} |u(x)| \leq C_\gamma \|A^\gamma u\|$  for  $u \in \mathcal{D}(A^\gamma)$ .

We can prove the lemma by the interpolation theory of Lions and a certain fact concerning  $\mathcal{D}(A)$ . For details, see H. Fujita-H. Morimoto [2] where analogical result is proved.

Lemma 2. For  $0 < \gamma < e$ ,  $\|A^\gamma e^{-tA}\| \leq t^{-\gamma}$ .

Proof. The proof is easy if we use the spectral representation of  $A$  and so we omit the proof.

We note here that by virtue of Lemma 1, the nonlinear operator  $P \nabla_u v$  is well-defined for every  $u \in \mathcal{D}(A^\gamma)$  if  $\gamma > \frac{3}{4}$ , and  $v \in (A^{1/2})$

and we have  $\|P \nabla_u v\| \leq C_\gamma \|A^\gamma u\| \|A^{1/2} v\|$ . We note also that in the

cartesian coordinate system,  $\nabla_u v = \sum_{j=1}^3 \left( \sum_{i=1}^3 u_i \frac{\partial v_j}{\partial x_i} \right) \frac{\partial}{\partial x_j} \equiv (u \cdot \nabla) v$

and so the above estimate is an immediate consequence.

Now we are ready to study the integral equation (IE). For that purpose we introduce a function space  $\Psi_T^Y$ , for  $T (> 0)$  and  $(\frac{3}{4} <) \gamma (< 1)$ .  $\Psi_T^Y$  is a space of all  $\mathcal{D}(A^{1/2})$ -valued functions  $u(t)$  defined on the interval  $[0, T]$  such that (i)  $u(t) \in C([0, T]; \mathcal{D}(A^{1/2})) \cap C((0, T]; \mathcal{D}(A^Y))$ , (ii) and the norm

$$\|u\| = \sup_{0 \leq t \leq T} \|A^{1/2}u(t)\| + \sup_{0 < t \leq T} \sup_{0 < s \leq t} s^{\gamma - \frac{1}{2}} \|A^Y u(s)\|$$

is finite.

We are going to obtain solutions of the integral equation (IE) in the class  $\Psi_T^Y$  by the iteration method. We use the following iteration scheme.

$$(3.1) \quad \begin{cases} u_0(t) = 0, \\ u_{n+1}(t) = e^{-tA} a - \int_0^t e^{-(t-s)A} P[(u_n(s) \cdot \nabla)u_n(s) + (w \cdot \nabla)u_n(s) + (u_n(s) \cdot \nabla)w] ds. \end{cases}$$

First we must verify that the iteration is possible in  $\Psi_T^Y$ .

To that end we introduce functions  $K_u(t)$ ,  $M_u(t)$  for functions  $u(t)$  in  $\Psi_T^Y$ . They are defined as follows.

$$K_u(t) = \sup_{0 \leq s \leq t} s^{\gamma - \frac{1}{2}} \|A^Y u(s)\|, \quad M_u(t) = \max_{0 \leq s \leq t} \|A^{1/2}u(s)\|.$$

And in addition we define operators  $\mathcal{A}, \mathcal{B}, \mathcal{C}$  by

$$\begin{aligned} \mathcal{A}(u)(t) &= \int_0^t e^{-(t-s)A} P(u(s) \cdot \nabla)u(s) ds \\ \mathcal{B}(u)(t) &= \int_0^t e^{-(t-s)A} P(w \cdot \nabla)u(s) ds \\ \mathcal{C}(u)(t) &= \int_0^t e^{-(t-s)A} P(u(s) \cdot \nabla)w ds \end{aligned}$$

Then we obtain the following estimates which justify the iteration.

Lemma 3.

$$\begin{aligned} \sup_{0 < s \leq t} s^{\gamma - \frac{1}{2}} \|A^\gamma A(u)(s)\| &\leq t^{1-\gamma} C_{\gamma B_1} K_u(t) M_u(t) \\ \sup_{0 < s \leq t} s^{\gamma - \frac{1}{2}} \|A^\gamma B(u)(s)\| &\leq t^{1/2} \frac{C_\gamma}{1-\gamma} \|A^\gamma \omega\| M_u(t) \\ \sup_{0 < s \leq t} s^{\gamma - \frac{1}{2}} \|A^\gamma C(u)(s)\| &\leq t^{1-\gamma} C_{\gamma B_1} \|A^{1/2} \omega\| K_u(t) \\ \sup_{0 < s \leq t} \|A^{1/2} A(u)(s)\| &\leq t^{1-\gamma} C_{\gamma B_1} K_u(t) M_u(t) \\ \sup_{0 < s \leq t} \|A^{1/2} B(u)(s)\| &\leq t^{1/2} 2C_\gamma \|A^\gamma \omega\| M_u(t) \\ \sup_{0 < s \leq t} \|A^{1/2} C(u)(s)\| &\leq t^{1-\gamma} C_{\gamma B_1} \|A^{1/2} \omega\| K_u(t) \end{aligned}$$

Here we write  $B_1 = B(1-\gamma, \frac{3}{2} - \gamma)$  where  $B(\cdot, \cdot)$  is the beta function.

Proof. We prove the first estimate only. The others are proved similarly.

$$\begin{aligned} \int_0^s \|A^\gamma e^{-(s-\sigma)A} \cdot P(u(\sigma) \cdot \nabla) u(\sigma)\| d\sigma &\leq \int_0^s (s-\sigma)^{-\gamma} C_\gamma \|A^\gamma u(\sigma)\| \|A^{1/2} u(\sigma)\| d\sigma \\ &\leq C_\gamma \int_0^s (s-\sigma)^{-\gamma} \sigma^{\frac{1}{2} - \gamma} K_u(s) M_u(s) d\sigma = C_\gamma K_u(s) M_u(s) \int_0^1 (1-\rho)^{-\gamma} \rho^{\frac{1}{2} - \gamma} s^{\frac{3}{2} - 2\gamma} d\rho \\ &= s^{\frac{3}{2} - 2\gamma} C_\gamma B(1-\gamma, \frac{3}{2} - \gamma) K_u(s) M_u(s). \end{aligned}$$

Hence we have

$$\sup_{0 < s \leq t} s^{\gamma - \frac{1}{2}} \|A^\gamma A(u)(s)\| \leq C_{\gamma B_1} \sup_{0 < s \leq t} s^{1-\gamma} K_u(s) M_u(s) = C_{\gamma B_1} t^{1-\gamma} K_u(t) M_u(t).$$

In the last equality we use the fact that  $K_u(t)$  and  $M_u(t)$  are increasing functions and the assumption that  $\frac{3}{4} < \gamma < 1$ . Q. E. D.

By Lemma 3, we have

$$\sup_{0 < s \leq t} s^{\gamma - \frac{1}{2}} \| A [A(u(s)) + B(u(s)) + C(u(s))] \| \leq t^{1-\gamma} [C_1 K_u(t) M_u(t) + C_2 K_u(t) + C_3 t^{\gamma - \frac{1}{2}} M_u(t)]$$

and

$$\sup_{0 < s \leq t} \| A^{1/2} [A(u(s)) + B(u(s)) + C(u(s))] \| \leq t^{1-\gamma} [C_1 K_u(t) M_u(t) + C_2 K_u(t) + C_3 M_u(t)]$$

Therefore, defining  $N_u(t) = \max\{K_u(t), M_u(t)\}$  and  $\mathcal{D}(u) = A(u) + B(u) + C(u)$ , we have

$$(3.2) \quad N_{\{u\}}(t) \leq t^{1-\gamma} [C_1 N_u^2(t) + (c_2 + c_3 t^{\gamma - \frac{1}{2}}) N_u(t)]$$

where we used  $c_1, c_2, c_3$  to denote positive constants depending only on  $\gamma$  and the Couéte flow  $w$ .

We now return to the iteration scheme (3.1). By the estimate

(3.2) we have the following recurrence inequality

$$(3.3) \quad N_{u_{n+1}}(t) \leq \| A^{1/2} a \| + t^{1-\gamma} [c_1 N_{u_n}^2(t) + (c_2 + c_3 t^{\gamma - \frac{1}{2}}) N_{u_n}(t)].$$

By a simple consideration we have for every  $u_n$

$$(3.4) \quad N_{u_n}(t) \leq X(t)$$

if  $c_2 t^{1-\gamma} + c_3 t^{1/2} < 1$  and  $\Delta(t) \geq 0$

where we define

$$(3.5) \quad \Delta(t) = (c_2 t^{1-\gamma} + c_3 t^{1/2} - 1)^2 - 4c_1 t^{1-\gamma} \| A^{1/2} a \|^2$$

and

$$(3.6) \quad \chi(t) = [1 - (c_2 t^{1-\gamma} + c_3 t^{1/2}) - \Delta(t)^{1/2}] / 2c_1 t^{1-\gamma}.$$

We can now study the convergence of the iteration. Setting  $v_n(t) = u_{n+1}(t) - u_n(t)$ , we have

$$v_0(t) = e^{-tA} a \quad \text{and}$$

$$v_n(t) = - \int_0^t e^{-(t-s)A} P[(v_{n-1}(s) \cdot \nabla) u_n(s) + (u_{n-1}(s) \cdot \nabla) v_{n-1}(s) + (w \cdot \nabla) v_n(s) + (v_{n-1}(s) \cdot \nabla) w] ds.$$

In order to estimate  $v_n(t)$  we define

$$D_n(t) = \sup_{0 < s \leq t} s^{\gamma - \frac{1}{2}} \|A^\gamma v_n(s)\|, \quad E_n(t) = \sup_{0 < s \leq t} \|A^{1/2} v_n(s)\|,$$

$$F_n(t) = \max\{D_n(t), E_n(t)\}, \quad M_n(t) = K_{u_n}(t), \quad M_n(t) = M_{u_n}(t), \quad N_n(t) = N_{u_n}(t).$$

Then we have

$$F_n(t) \leq t^{1-\gamma} (d_1 \chi(t) + d_2 \|A^\gamma w\| t^{\gamma - \frac{1}{2}} + d_3 \|A^{1/2} w\|) F_{n-1}(t) \equiv \rho(t, w) F_n(t)$$

and

$$F_0(t) \leq \|A^{1/2} a\|$$

where  $d_1, d_2, d_3$  are positive constants depending only on  $\gamma$ . If we

note that for every fixed  $w$ ,  $\chi(t)$  tends to 0 as  $t$  tends to 0,

we immediately see that there exists a positive  $T$  depending only on

$$\|A^\gamma w\| \quad \text{such that} \quad \sum_{n=0}^{\infty} F_n(t) \quad \text{converges uniformly in} \quad t \in [0, T].$$

Hence we see, noting that  $A$  is a closed operator and has a continuous

inverse, that  $u(t) = \lim_{n \rightarrow \infty} u_n(t)$  exists in  $L^2_\sigma$  and  $\mathcal{D}(A^{1/2})$  for

$t \in [0, T]$  and in  $\mathcal{D}(A^\gamma)$  for every  $t \in (0, T]$  and that the former convergence is uniform on  $[0, T]$  and the latter locally uniform in  $(0, T]$ . The fact that the limit function belongs to  $\Psi_T^\gamma$  is evident. The estimates above show that for a given  $r > 0$  we can choose  $T > 0$  such that the limit function exists on the interval  $[0, T]$  for every initial data  $a \in \mathcal{D}(A^{1/2})$  such that  $\|A^{1/2}a\| \leq r$ .

For the proof of assertion (2) of Theorem 1, we note that (3.5) and (3.6) show that  $\chi(t) \rightarrow 0$  when  $\|A^{1/2}a\| \rightarrow 0$ . This and the fact that in the inequality  $c_2 T^{1-\gamma} + c_3 T < 1$   $c_2$  and  $c_3$  depend only on  $\gamma$  indicate that for arbitrary given  $T' > 0$  we can choose  $r > 0$  such that we can construct a solution  $u(t) \in \Psi_{T'}^\gamma$  for every  $u(0) = a \in \mathcal{D}(A^{1/2})$  such that  $\|A^{1/2}a\| \leq r$ .

Thus we have proved the following

Theorem 1.

(1) For every given  $r > 0$  we can choose  $T > 0$  such that for every  $a \in \mathcal{D}(A^{1/2})$  with  $\|A^{1/2}a\| \leq r$ , there exists a solution  $u(t)$  of the integral equation (IE) in the interval  $[0, T]$  which belongs to the class  $\Psi_T^\gamma$ .

(2) For every given  $T > 0$  we can choose  $r > 0$  such that there exists a solution of (IE) on  $[0, T]$  for every  $a \in \mathcal{D}(A^{1/2})$  with  $\|A^{1/2}a\| \leq r$ .

Remark. We have not mentioned the uniqueness of the solution.

We have omitted the proof since it is not difficult to prove the uniqueness of the solution in the class  $\Psi_T^\gamma$ .

4. Differentiability of the evolution operator  $S_t$ .

First we remember the definition of the Fréchet derivative.

Definition. Let  $X, Y$  be Banach spaces and  $\Phi$  be a continuous mapping defined in a neighbourhood  $U$  of an element  $a \in X$ . A bounded linear operator  $A$  is said to be the Fréchet derivative of  $\Phi$  at  $a$  if

$$\Phi(a + h) - \Phi(a) = Ah + o(\|h\|) \quad \text{as } a + h \text{ tends to } a \text{ in } U.$$

And when this is the case,  $\Phi$  is said to be Fréchet differentiable at  $a$ .

We now return to the integral equation (IE) and define the evolution operator  $S_t : \mathcal{D}(A^{1/2}) \rightarrow \mathcal{D}(A^{1/2})$  by  $S_t a = u(t)$  where  $u(t)$  is the solution (IE) with initial data  $a$ . By Theorem 1, we know that for any given  $T > 0$  we can find a neighbourhood  $U$  of  $0 \in \mathcal{D}(A^{1/2})$  where  $S_T$  is defined everywhere. Hence we can talk about the Fréchet differentiability of  $S_T$  at  $0$ . We fix  $U$  and  $T$  above. For  $h \in U$ ,  $S_t$  satisfies the following integral equation by definition

$$S_t h = e^{-tA} h - \int_0^t e^{-(t-s)A} P[(S_s h \cdot \nabla) S_s h + (w \cdot \nabla) S_s h + (S_s \cdot \nabla) w] ds.$$

An inspection indicates that the Fréchet derivative of  $S_t$  at  $0 \in \mathcal{D}(A^{1/2})$  which we denote by  $DS_t$  must satisfy the following integral equation if it exists

$$(4.1) \quad DS_t h = e^{-tA} h - \int_0^t e^{-(t-s)A} P[(w \cdot \nabla) DS_s h + (DS_s h \cdot \nabla) w] ds.$$

In order to integrate the equation (4.1) we introduce the function space  $\Psi_t^Y$  again and define an operator by

$$\Gamma(f)(t) = e^{-tA}h - \int_0^t e^{-(t-s)A} P[(w \cdot \nabla)f(s) + (f(s) \cdot \nabla)w] ds.$$

The operator  $\Gamma$  is well-defined since we have the following estimate

$$(4.2) \quad \max\{ \sup_{0 < s \leq t} s^{\gamma - \frac{1}{2}} \|A^\gamma \Gamma(f)(s)\|, \sup_{0 < s \leq t} \|A^{1/2} \Gamma(f)(s)\| \} \leq \|A^{1/2} h\| + c(t^{1-\gamma} K_f(t) + t^{1/2} M_f(t)),$$

with positive constant  $c$  depending only on  $\gamma$  and  $w$  (the Couéte flow).

And we have for  $f_1, f_2 \in \Psi_T^\gamma$ ,

$$(4.3) \quad \|\Gamma(f_1) - \Gamma(f_2)\| \leq c_1 \tau^{1-\gamma} \|A^\gamma w\| (1 + c_2 \tau^{\gamma - \frac{1}{2}}) \|f_1 - f_2\|$$

with positive constants  $c_1, c_2$  depending only on  $\gamma$ . We choose

$\tau > 0$  such that

$$(4.4) \quad c_1 \tau^{1-\gamma} \|A^\gamma w\| (1 + c_2 \tau^{\gamma - \frac{1}{2}}) < 1.$$

Then there exists uniquely in  $\Psi_T^\gamma$  a solution which we denote by  $f(t; h)$ .

Next we shall prove, making use of  $f(t; h)$ , that there exists

$0 < \tau' < \tau$  such that  $S_{\tau'}$  is Fréchet differentiable at 0. We set

$g(t; h) = S_t h$ , then we have

$$(4.5) \quad f(t; h) - g(t; h) = \int_0^t e^{-(t-s)A} \cdot P[(g(s; h) \cdot \nabla)g(s; h)] ds + \int_0^t e^{-(t-s)A} \cdot P\{(w \cdot \nabla)(f(s; h) - g(s; h)) + [(f(s; h) - g(s; h)) \cdot \nabla]w\} ds.$$

Then what we have to prove is that there exists  $\tau' > 0$  such that

$$\|A^{1/2}(f(\tau'; h) - g(\tau'; h))\| / \|A^{1/2}h\| \rightarrow 0 \text{ as } \|A^{1/2}h\| \rightarrow 0.$$

Estimating (4.2), we have

$$\| \| f(t ; h) - g(t ; h) \| \| \leq c_3 \tau'^{1-\gamma} \| A^\gamma w \| (1 + c_4 \tau'^{\gamma-\frac{1}{2}}) \| \| f(t ; h) - g(t ; h) \| \| + c_5 \tau'^{1-\gamma} \| A^\gamma w \| L(\tau')^2$$

with positive constants  $c_3, c_4, c_5$  depending only on  $\gamma$  where the norm is for functions  $f, g \in \Psi_{\tau'}^\gamma$ , and

$$L(t) = \max \left\{ \sup_{0 < s \leq t} s^{\gamma-\frac{1}{2}} \| A^\gamma g(s ; h) \|, \sup_{0 < s \leq t} \| A^{1/2} g(s ; h) \| \right\}.$$

If we choose  $\tau' > 0$  such that

$$(4.6) \quad c_3 \tau'^{1-\gamma} \| A^\gamma w \| (1 + c_4 \tau'^{\gamma-\frac{1}{2}}) < \rho < 1,$$

we have

$$\| \| f(t ; h) - g(t ; h) \| \| < (1 - \rho)^{-1} c_5 \tau'^{1-\gamma} \| A^\gamma w \| L(\tau')^2.$$

This implies that

$$f(\tau' ; h) - g(\tau' ; h) = O(\| A^{1/2} h \|^2)$$

which was to be proved. Thus we have proved that  $S_{\tau'}$  is Fréchet differentiable at 0. By the estimates (4.2) and (4.4), we see that  $\tau'$  is determined only  $\gamma$  and  $w$  and so, by the chain rule for Fréchet derivatives, we have the following

Theorem 2.

For every  $T > 0$  the evolution operator  $S_t$  is Fréchet differentiable at  $0 \in \mathcal{D}(A^{1/2})$ .

## 5. The eigenvalue problem.

In this section we study the eigenvalue problem of the linear operator  $E = DS_t$  which is the Fréchet derivative at zero of the evolution operator  $S_t$ . What we wish to know is whether  $E$  has a <sup>real positive</sup> eigenvalue ~~greater than 1~~ greater than 1 or not. If this is the case, the corresponding Couéte flow must be unstable, or we define that it is unstable.

This problem can be reduced to investigate whether the eigenvalue problem

$$(5.1) \quad -\Delta u - P\nabla_u w - P\nabla_w y = \lambda u$$

admits real positive eigenvalues or not. It is easy to verify that

(5.1) is equivalent to

$$(5.2) \quad \Delta u - \nabla_u w - \nabla_w u - \text{grad } p = \lambda u, \quad \text{div } u = 0$$

with a suitable scalar function  $p$ . We adopt the cylindrical coordinate system  $(r, \phi, z)$ , where  $u = (u_r, u_\phi, u_z) = u_r \frac{\partial}{\partial r} + \frac{u_\phi}{r} \frac{\partial}{\partial \phi} + u_z \frac{\partial}{\partial z}$  and the Couéte flow  $w$  is expressed as  $w = (w_r, w_\phi, w_z) = (A + \frac{B}{r^2}) \frac{\partial}{\partial \phi}$ .

We recall that

$$A = (R_2^2 \Omega_2 - R_1^2 \Omega_1) / (R_2^2 - R_1^2), \quad B = R_1^2 R_2^2 (\Omega_1 - \Omega_2) / (R_2^2 - R_1^2).$$

In the cylindrical coordinate system, (5.2) is expressed as

$$(5.3) \left\{ \begin{array}{l} (\Delta - \frac{1}{r^2})u_r - \frac{2}{r^2} \frac{\partial}{\partial \phi} \phi - \mathcal{N}(w)u_r + \frac{w}{r} \frac{u}{\phi} - \mathcal{N}(u)w_r + \frac{u}{r} \frac{w}{\phi} - \frac{\partial}{\partial r} = \lambda u_r \\ (\Delta - \frac{1}{r^2})u_\phi - \frac{2}{r^2} \frac{\partial u_r}{\partial \phi} - \mathcal{N}(w)u_\phi - \frac{w}{r} \frac{u}{\phi} - \mathcal{N}(u)w_\phi - \frac{u}{r} \frac{w}{\phi} - \frac{1}{r} \frac{\partial}{\partial \phi} = \lambda u_\phi \\ u_z - \mathcal{N}(w)u_z - \mathcal{N}(u)w_z - \frac{\partial}{\partial z} = \lambda u_z \\ \frac{1}{r} \frac{\partial}{\partial r} (ru_r) + \frac{1}{r} \frac{\partial u}{\partial \phi} + \frac{\partial u_z}{\partial z} = 0 \end{array} \right.$$

where

$$\Delta = \frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial^2}{\partial \phi^2} + \frac{\partial^2}{\partial z^2} \quad \text{and} \quad \mathcal{N}(v) = v_r \frac{\partial}{\partial r} + \frac{v_\phi}{r} \frac{\partial}{\partial \phi} + v_z \frac{\partial}{\partial z}.$$

We seek  $u$  and  $p$  which are independent of the variable  $\phi$ . Then (5.3)

reduces to

$$(5.4) \left\{ \begin{array}{l} (\Delta - \frac{1}{r^2})u_r + 2(A + \frac{B}{r^2})u_\phi - \frac{\partial p}{\partial r} = \lambda u_r \\ (\Delta - \frac{1}{r^2})u_\phi - 2Au_r = \lambda u_\phi \\ \Delta u_z - \frac{\partial p}{\partial z} = \lambda u_z \\ \frac{1}{r} \frac{\partial}{\partial r} (ru_r) + \frac{\partial u_z}{\partial z} = 0 \end{array} \right.$$

Introducing a stream function  $f$  by the relations  $ru_r = \frac{\partial}{\partial z} (rf)$  and

$ru_z = -\frac{\partial}{\partial r} (rf)$ , we obtain from (5.4)

$$(5.5) \left\{ \begin{array}{l} L(L - \lambda)f + 2(A + \frac{B}{r^2})u_\phi = 0 \\ (L - \lambda)u_\phi - 2Af = 0 \end{array} \right.$$

where we write  $L = \frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} + \frac{\partial^2}{\partial z^2} - \frac{1}{r^2}$ . On  $f$  we impose an

additional boundary condition that  $f, \frac{\partial f}{\partial r} = 0$  for  $r = R_1, R_2$ .

We set  $f(r, z) = f(r) \cos z$ ,  $u_\phi(r, z) = u(r) \sin z$  with  $\sigma = \frac{2\pi}{h}$ . The

boundary condition is reduced to the condition  $f(R_i) = f'(R_i) = u(R_i)$ ,

for  $i = 1, 2$ . Hence we have the following system of linear ordinary

differential equations

$$(5.6) \quad \begin{cases} (\mathcal{D} - \sigma^2)(\mathcal{D}^2 - \sigma^2 - \lambda)f(r) + 2\left(A + \frac{B}{r^2}\right)\sigma u(r) = 0 \\ (\mathcal{D} - \sigma^2 - \lambda)u(r) + 2Af(r) = 0 \\ f(R_i) = f'(R_i) = u(R_i) = 0, \quad i = 1, 2 \end{cases}$$

where we write  $\mathcal{D} = \frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{1}{r^2}$ . In order to investigate the

system (5.6), we consider the following two boundary value problems

for ordinary differential operators  $\mathcal{D} - \mu$  for  $\mu \geq 0$ .

$$(BVP-1) \quad \begin{cases} (\mathcal{D} - \mu)g(r) = (r), \quad r \in (R_1, R_2) \\ g(R_i) = 0, \quad i = 1, 2. \end{cases}$$

$$(VBP-2) \quad \begin{cases} (\mathcal{D} - \mu_1)(\mathcal{D} - \mu_2)g(r) = (r), \quad r \in (R_1, R_2) \\ g(R_i) = g'(R_i) = 0, \quad i = 1, 2 \end{cases}$$

with  $\mu, \mu_i \geq 0$ .

The next lemma is useful.

Lemma 4. Let  $G(r, r'; \mu)$  and  $H(r, r'; \mu_1, \mu_2)$  be the Green

functions for (BVP-1) and (BVP-2) respectively. Then  $G$  and  $H$  are negative valued and positive valued respectively almost everywhere.

This lemma may be proved by an explicit construction of the kernels making use of the Bessel functions but it can be proved by repetition of elementary discussions. The details are left to the reader.

We now return to the system (5.6). From (5.6) we have

$$(5.7) \quad f(r) = - \int_{R_1}^{R_2} H(r, r' ; \sigma^2, \sigma^2 + \lambda) 2\sigma \left( A + \frac{B}{r'^2} \right) u(r') dr'$$

and

$$(5.8) \quad u(r) = - \int_{R_1}^{R_2} G(r, r' ; \sigma^2 + \lambda) 2A f(r') dr'.$$

Hence we have

$$(5.9) \quad f(r) = \int_{R_1}^{R_2} K(r, r' ; \lambda) f(r') dr',$$

$$\text{where } K(r, s ; \lambda) = 4\sigma \int_{R_1}^{R_2} H(r, r' ; \sigma^2, \sigma^2 + \lambda) A \left( A + \frac{B}{r'^2} \right) G(r', s ; \sigma^2, \lambda) ds.$$

From (5.7), (5.8) we have

$$(5.10) \quad f(r) = \int_{R_1}^{R_2} K(r, s ; \sigma, \lambda) f(s) ds.$$

Now the problem is reduced to that whether the integral operator  $K$  defined by the kernel  $K(r, s ; \sigma, \lambda)$  has 1 as its eigenvalue or not. For that purpose the following lemma of Jentzsch (See Schmeidler [4]) is useful.

Lemma 5. Let  $K(r, s)$  be a continuous kernel on the interval  $[R_1, R_2]$  which is positive almost everywhere. Then the integral operator

$Kf(r) = \int_{R_1}^{R_2} K(r,s)f(s)ds$  has a positive eigenvalue.

What we have to do next is to investigate the signature of the

$$\text{function } k(r) = A\left(A + \frac{B}{r^2}\right) = \frac{R_2^2 \Omega_2^2 - R_1^2 \Omega_1^2}{R_2^2 - R_1^2} \left[ (R_2^2 \Omega_2^2 - R_1^2 \Omega_1^2) + \frac{R_1^2 R_2^2 (\Omega_1^2 - \Omega_2^2)}{r^2} \right]$$

For a fixed  $\omega = (\omega_1, \omega_2) \in S^1$  (the 1-sphere) we set  $(\Omega_1, \Omega_2) = \rho(\omega_1, \omega_2)$ ,  $\rho \geq 0$ , and

$$k(r; \rho, \omega) = \rho^2 \frac{R_2^2 \omega_2^2 - R_1^2 \omega_1^2}{R_2^2 - R_1^2} \left[ (R_2^2 \omega_2^2 - R_1^2 \omega_1^2) + \frac{R_1^2 R_2^2 (\omega_1^2 - \omega_2^2)}{r^2} \right] \equiv \rho^2 \tilde{k}(r; \omega)$$

Then the kernel  $K$  is expressed by

$$(5.12) \quad K(r,s; \lambda) = \rho^2 4\sigma \int_{R_1}^{R_2} H(r,r'; \sigma^2, \sigma^2 + \lambda) \tilde{k}(r'; \omega) K(r',s; \sigma^2, \lambda) dr' \\ \equiv \rho^2 L(r,s; \lambda, \omega).$$

Hence, making use of Lemma 4 and Lemma 5, we have

Theorem 3.

For every fixed  $\omega = (\omega_1, \omega_2) \in S^1$ , such that  $(R_2^2 \omega_2^2 - R_1^2 \omega_1^2)(\omega_2 - \omega_1) > 0$  there exists  $\rho(\omega) > 0$  such that for every  $\rho > \rho(\omega)$  the corresponding Couéte flow is unstable under infinitesimal perturbations.

It suffices only to notice that the integral equation (5.10) is reduced to

$$f(r) = \rho^2 \int_{R_1}^{R_2} L(r,s; \lambda, \omega) f(s) ds.$$

## 6. Isolatedness of the Couétté flow.

What we are going to do in this section is to investigate whether or not 0 is an isolated solution of the equation

$$(E) \quad Au - P(u \cdot \nabla)u + P(w \cdot \nabla)u + P(u \cdot \nabla)w = 0$$

where  $w$  is the Couétté flow. We take  $(1 >) \gamma (> \frac{3}{4})$  and work in  $\mathcal{D}(A^\gamma)$ . Assume that 0  $\in \mathcal{D}(A^\gamma)$  is not an isolated solution of (E).

Then there exists solutions  $u_n (\neq 0)$  of (E) for  $n = 1, 2, \dots$  such

that  $\lim_{n \rightarrow \infty} \|A^\gamma u_n\| = 0$ . Set  $\phi_n = u_n / \|A^\gamma u_n\|$ . Then we have

$$(6.1) \quad A\phi_n + \|A^\gamma u_n\| P(\phi_n \cdot \nabla)\phi_n + P(w \cdot \nabla)\phi_n + P(\phi_n \cdot \nabla)w = 0.$$

As for the second term, we see that

$$\|P(\phi_n \cdot \nabla)\phi_n\| \leq C_\gamma \|A^\gamma \phi_n\| \|A^{1/2} \phi_n\| \leq C_\gamma \|A^{\frac{1}{2}-\gamma}\| \|A^\gamma \phi_n\|^2 = C_\gamma \|A^{\frac{1}{2}-\gamma}\|$$

and so  $\lim_{n \rightarrow \infty} \|A^\gamma u_n\| P(\phi_n \cdot \nabla)\phi_n = 0$ . Hence

$$\lim_{n \rightarrow \infty} \{A\phi_n + P(w \cdot \nabla)\phi_n + P(\phi_n \cdot \nabla)w\} = 0 \text{ strongly in } L^2_\sigma.$$

And so by the boundedness of  $A^{\gamma-1}$  we have

$$(6.2) \quad \lim_{n \rightarrow \infty} \{A^\gamma \phi_n + A^{\gamma-1} [P(w \cdot \nabla)\phi_n + P(\phi_n \cdot \nabla)w]\} = 0.$$

By the compactness of the inclusions  $\mathcal{D}(A^\gamma) \rightarrow \mathcal{D}(A^{1/2}) \rightarrow L^2_\sigma$  and the equality  $\|\nabla \varphi\| = \|A^{1/2} \varphi\|$  for  $\varphi \in \mathcal{D}(A^{1/2})$  we see that

$\lim_n [P(w \cdot \nabla)\phi_n + P(\phi_n \cdot \nabla)w]$  exists strongly in  $L^2_\sigma$ , and this implies

that  $\phi_\infty = \lim_{n \rightarrow \infty} \phi_n$  exists strongly in  $\mathcal{D}(A^\gamma)$ . Hence we have

$$(6.3) \quad \begin{cases} A^\gamma \phi_\infty + A^{\gamma-1} [P(w \cdot \nabla) \phi_\infty + P(\phi_\infty \cdot \nabla) w] = 0 \\ \|A^\gamma \phi_\infty\| = 1. \end{cases}$$

This implies that the operator  $B$  defined by  $B\varphi = -A^{-1} [P(w \cdot \nabla)\varphi + P(\varphi \cdot \nabla)w]$  in  $L_\sigma^2$  with domain of definition  $\mathcal{D}(A^{1/2})$  has 1 as its eigenvalue.

We now ~~remember~~ <sup>recall</sup> the explicit form of  $w$ .

$$w = \frac{w}{r} \frac{\partial}{\partial \phi} = \rho \frac{1}{R_2^2 - R_1^2} [R_2^2 \omega_2 - R_1^2 \omega_1 + \frac{R_1^2 R_2^2 (\omega_1 - \omega_2)}{r^2}] \frac{\partial}{\partial \phi}$$

where  $\rho \geq 0$ ,  $\omega = (\omega_1, \omega_2) \in S^1$ . We set  $w = \rho W_\omega$ . Then  $B\varphi = -\rho A^{-1} [P(W_\omega \cdot \nabla)\varphi + P(\varphi \cdot \nabla)W_\omega]$ . By the compactness of the operator  $A^{-1}$  and the equality  $\|\nabla \varphi\| = \|A^{1/2} \varphi\|$  for  $\varphi \in \mathcal{D}(A^{1/2})$  we can prove that  $B$  can be extended to the whole space  $L_\sigma^2$  and the resulting operator is a compact operator. Hence we have

Theorem 4. The Couétté flow is an isolated solution in  $W_2^\gamma(G_h)$  ( $\gamma > \frac{3}{4}$ ) for almost all  $\Omega = (\Omega_1, \Omega_2) \in \mathbb{R}^2$ .

REFERENCES

- [1] H. Fujita-Kato, T., On the Navier-Stokes initial value problem I,  
Arch. Rational Mech. Anal. 16 (1964), 269-315.
- [2] H. Fujita-Morimoto, H., On fractional powers of the Stokes operator,  
Proc. Japan Acad. Vol. 46, No. 10 (1970), 1141-1143.
- [3] E. C. Lin, The theory of hydrodynamical stability, Oxford (1953).
- [4] W. Schmeidler, Integralgleichungen mit Anwendungen in Physik und  
Technik, Bd. I. Berlin 1950.
- [5] A. Takeshita, On the reproductive property of the 2-dimensional  
Navier-Stokes equations,  
J. Fac. Sci. Univ. Tokyo, I, Vol. XVI, part 3, (1970), 279-311.
- [6] W. Velte, Stabilität und Verzweigungen der Navier-Stokeschen  
Gleichungen beim Taylorproblem, Arch. Rational Anal. 22 (1966),  
1-14.