

On the stability of incompressible viscous fluid motions  
past objects

by <sup>^</sup>Kyuya MASUDA (Dept. of Math., Univ. of  
Tokyo, Komaba )

Let  $E$  be the exterior domain in 3-space. Let us consider  
the steady flow in  $E$  governed by

$$(1) \quad \begin{cases} -\nu \Delta w + (w \cdot \nabla) w + \nabla p = 0, \\ \operatorname{div} w = 0 \end{cases}$$

$$(2) \quad w(x) \rightarrow w^\infty \quad (|x| \rightarrow \infty)$$

$$(3) \quad w(x) = b(x) \quad (x \in \partial E)$$

where the viscosity coefficient  $\nu$  is a positive constant,  
 $w^\infty$  is some fixed constant vector,  $b$  is some prescribed  
function on  $E$ .

R. Finn showed that if  $w^\infty - b$  is "small" enough, then  
there exists a smooth solution  $w$  with

$$\sup_{x \in E} |x| |w(x) - w^\infty| < \infty$$
$$\nabla w \in L^3(E)$$

1

Given the disturbance  $u_0 \in L^2(\mathcal{E})$  to  $w$ . Then the perturbed flow  $v$  is governed by

$$(4) \quad \begin{cases} \frac{\partial v}{\partial t} - \nu \Delta v + (v \cdot \nabla) v - \nabla p_2 = 0 \\ \operatorname{div} v = 0 \end{cases}$$

$$(5) \quad \begin{cases} \lim_{|x| \rightarrow \infty} v(x, t) = w^\infty, & v(x, t) = b(x) \quad (x \in \partial \mathcal{E}, t > 0) \\ \lim_{t \downarrow 0} v(x, t) = w(x) + u_0(x) \end{cases}$$

Now our result is:

Assume that

- (i)  $\sup_{x \in \mathcal{E}} |x| |w(x) - w^\infty| < \frac{1}{2}$
- (ii)  $\nabla w \in L^3(\mathcal{E})$
- (iii)  $\operatorname{div} w = 0$

Then every weak solution  $v$  of (4), (5) becomes analytic (in  $t$  and  $x$ ) after some definite time  $T_0$ , and then converges to steady flow  $w$  uniformly in  $x$  on  $\mathcal{E}$  like

$$|v(x, t) - w(x)| \leq M t^{-1/8}, \quad (t \rightarrow \infty)$$

( $M$ ; constant)