

On The Symplectic Lazard Ring

by Kazumoto KOJIMA

0. Introduction

In 'Elementary proofs of some results of cobordism theory using Steenrod operations' (Advances in Math., 7 (1971), 29-56. ), D.Quillen determined complex cobordism ring  $MU_*$  using the formal group theory. This method is not applicable directly for the symplectic case.

However there are some works along ~~in~~<sup>in</sup> this line. Espetially, Buhstaber-Novikov studied two-valued formal groups and gave some applications to symplectic cobordism ring  $MSp_*$ .

We will define symplectic formal system using formal power series like as (two-valued) formal group, and construct a geometrical example of symplectic formal system. To construct this geometrical example, We need some stable maps between the complex (or symplectic) projective and quasiprojective space.

Moreover, we can construct a ring associated with symplectic formal system. We denote the symplectic Lazard ring  $LMSp$  as the associated ring for the universal symplectic formal system.

Then, we can construct a homomorphism  $\theta: LMSp \rightarrow MSp_*/Torsion$ . By some calculations and the result of R.Ōkita ('On the  $MSp$  Hattori-Stong problem', Osaka J. math. 13 (1976), 547-566.), we can conclude that if we apply the rational indecomposable functor  $Q( )$ , then  $Q(\theta)$  is an isomorphism.

### 1. Stable maps

There is a symplectification map  $q : \mathbb{C}P^\infty \longrightarrow \mathbb{H}P^\infty$ .

Since  $q$  is a fibre bundle whose fibre is  $S^2$ , there is a Becker-Gottlieb transfer  $t : \mathbb{H}P_+^\infty \xrightarrow{(s)} \mathbb{C}P_+^\infty$ .

Let  $F$  be  $\mathbb{C}$  or  $\mathbb{H}$  and  $S_F^n$  unit sphere in  $F^n$ .

Let  $G_n(\mathbb{C}) = U(n)$  and  $G_n(\mathbb{H}) = Sp(n)$ .

The quasiprojective space  $Q_n(F)$  is defined to be the space of generalized reflections, that is, the image of

$$\phi : S_F^n \times S_F^1 \longrightarrow G_n(F)$$

where  $\phi(u, q)$  is the automorphism which leaves  $v$  fixed if  $\langle u, v \rangle = 0$  and sends  $u$  to  $uq$ .

We may define  $Q_n(F)$  as the space obtained  $S_F^n \times S_F^1$  by imposing the equivalence relation  $(u, q) \sim (ug, g^{-1}qg)$  ( $g \in S_F^1$ ), and collapsing  $S_F^n \times 1$  to a point.

By the second definition, we can easily show that  $Q_n(\mathbb{C}) \approx \Sigma(\mathbb{C}P_+^{n-1})$ .

We put  $\widehat{\mathbb{C}P}^n = Q_n(\mathbb{C})$  and  $\widehat{\mathbb{H}P}^n = Q_n(\mathbb{H})$ . Clearly

we have a symplectification map  $\tilde{q} : \widehat{\mathbb{C}P}^\infty \longrightarrow \widehat{\mathbb{H}P}^\infty$ .

Now we construct a map from  $\widehat{\mathbb{H}P}^n$  to  $\widehat{\mathbb{C}P}^{2n}$ .

Let  $z \in \mathbb{H}^n$  and  $z = x + jy$  where  $x, y \in \mathbb{C}^n$ .

We denote complexification map  $c : \mathbb{H}^n \longrightarrow \mathbb{C}^{2n}$  by setting  $c(z) = x \oplus y \in \mathbb{C}^{2n}$ .

Let  $q = a + jb \in \mathbb{H}$  where  $a, b \in \mathbb{C}$ . Since  $S_C^1$  is a maximal torus of  $S_H^1$ , there is a  $g \in S_H^1$  such that  $g^{-1}qg \in S_C^1$ . If  $g^{-1}qg = e^{i\mathcal{L}t}$ , where  $-1 < t < 0$ , then  $(gj)^{-1}qgj = e^{-i\mathcal{L}t}$ .

Thus there is a  $g \in S_H^1$  such that  $g^{-1}qg = e^{i\mathcal{L}t}$  where  $0 \leq t \leq 1$ .

So a representative element of  $\widetilde{HP}^n$  can be taken as  $(x + jy, e^{i\pi t})$  where  $x, y \in \mathbb{C}^n$  and  $0 \leq t \leq 1$ .

We define  $\tilde{t}_n : \widetilde{HP}^n \rightarrow \widehat{CP}^{2n}$  by the equation  $\tilde{t}_n[(x + jy, e^{i\pi t})] = [(x \oplus y, e^{2i\pi t})]$ .

Then the following proposition holds.

Proposition. The diagram

$$\begin{array}{ccc} \widetilde{HP}^n & \xrightarrow{\tilde{t}_n} & \widehat{CP}^{2n} \\ \downarrow j & & \downarrow j \\ SP(n) & \xrightarrow{c} & U(2n) \end{array} \quad \text{commutes up to homotopy.}$$

By the theorem of Becker-Segal,

$Q(HP^\infty) \simeq BSp \times F$  ~~as an infinite loop space~~ where  $Q(\ )$  is a stabilize functor  $\varinjlim_n \Omega^n S^n(\ )$ .

So we have a map  $r : \Sigma \widetilde{HP}^\infty \rightarrow Q(HP^\infty)$  such that the diagram

$$\begin{array}{ccc} \Sigma \widetilde{HP}^\infty & \xrightarrow{j} & \Sigma Sp \\ r \downarrow & & \downarrow \wr \\ Q(HP^\infty) & \xrightarrow{j} & BSp \end{array} \quad \text{commutes up to homotopy.}$$

We may regard  $r$  as a stable map  $r : \Sigma \widetilde{HP}^\infty \xrightarrow{(s)} HP^\infty$ .

We put  $\overline{HP}^\infty = \Sigma^{-1} \widetilde{HP}^\infty$ ,  $\overline{q} = \Sigma^{-1} \overline{q}$  and  $\overline{t} = \Sigma^{-1} \overline{t}$ .

Then we have following stable maps :

$$\begin{array}{l} CP_+^\infty \xrightarrow{q} HP_+^\infty \xrightarrow{t} CP_+^\infty, \\ CP_+^\infty \xrightarrow{\overline{q}} \overline{HP}_+^\infty \xrightarrow{\overline{t}} CP_+^\infty \quad \text{and} \\ \Sigma^2 \overline{HP}^\infty \xrightarrow{r} HP^\infty. \end{array}$$

We can easily calculate the homomorphisms induced by these maps on the ordinaly

homology theory.

Let  $y^{\text{MSP}}$  be the euler class of MSP,  $y$  the class of ordinaly homology.

Then in H MSP-theory, we have  $y^{\text{MSP}} = h(y) = \sum_{i \geq 0} h_i y^{i+1}$ .

Let  $x$  be the complex euler class of the ordinaly homology H.

Now we can define the symplectic formal system.

Let  $R$  be a commutative ring with unit and  $R[[X, \bar{X}, Y, \bar{Y}]]$  formal power series ring with four variables  $X, \bar{X}, Y$  and  $\bar{Y}$ .

Definition 4.1. A symplectic formal system is a set of formal power series  $E(X), F_k(X, \bar{X}, Y, \bar{Y})$  and  $G_k(X, \bar{X}, Y, \bar{Y})$  (for  $k \geq 1$ ) such that satisfy

$$(i) \quad E(X) = \sum_{i \geq 1} a_i X^i,$$

$$F_k(X, \bar{X}, Y, \bar{Y}) = \sum_{i, j \geq 0} b_{i, j}^{(k)} X^i \cdot Y^j + \sum_{i, j \geq 1} c_{i, j}^{(k)} \bar{X} \cdot X^{i-1} \bar{Y} \cdot Y^{j-1},$$

$$G_k(X, \bar{X}, Y, \bar{Y}) = \sum_{i, j \geq 0} d_{i, j}^{(k)} (\bar{X} \cdot X^{i-1} Y^j + \bar{Y} \cdot Y^{i-1} X^j)$$

and under  $\bar{X}^2 = E(X), \bar{Y}^2 = E(Y)$ , satisfy also

$$(ii) \quad (\text{unitary relation}) \quad b_{1,0}^{(1)} = d_{1,0}^{(1)} = 1, \quad b_{n,0}^{(1)} = d_{n,0}^{(1)} = 0 \quad \text{for } n \neq 1,$$

(iii) (associative relation)

$$D(F_1(X, \bar{X}, Y, \bar{Y}), G_1(X, \bar{X}, Y, \bar{Y}), Z, \bar{Z})$$

$$= D(X, \bar{X}, F_1(Y, \bar{Y}, Z, \bar{Z}), G_1(Y, \bar{Y}, Z, \bar{Z})) \quad \text{for } D = F_1 \text{ or } G_1,$$

$$(iv) \quad (\text{commutative relation}) \quad b_{i,j}^{(1)} = b_{j,i}^{(1)}, \quad c_{i,j}^{(1)} = c_{j,i}^{(1)},$$

$$(v) \quad (\text{differntial relation}) \quad c_{1,1}^{(1)} = -2, \quad c_{1,n}^{(1)} = c_{n,1}^{(1)} = 0 \quad \text{for } n \neq 1,$$

$$(vi) \quad (\text{power relation}) \quad F_k(X, \bar{X}, Y, \bar{Y}) = (F_1(X, \bar{X}, Y, \bar{Y}))^k,$$

$$G_k(X, \bar{X}, Y, \bar{Y}) = G_1(X, \bar{X}, Y, \bar{Y}) F_{k-1}(X, \bar{X}, Y, \bar{Y}) \quad \text{and}$$

$$(vii) \quad (\text{squar relation}) \quad (G_1(X, \bar{X}, Y, \bar{Y}))^2 = E(F_1(X, \bar{X}, Y, \bar{Y})).$$

Definition 4.2.

Let  $\Gamma = \{E, F_k, G_k\}$  be a symplectic formal system over  $R$ .

An associated symplectic ring for  $\Gamma$ ,  $R_\Gamma$ , is the subring of  $R$  which is generated by the elements  $8a_i, 4b_{i,j}^{(2k-1)}, 2b_{i,j}^{(2k)}, c_{i,j}^{(k)}, 4d_{i,j}^{(k)}$  and  $1$ .

Now we can define symplectic Lazard ring  $\text{LMSp}$  as follows.

Let  $S$  be  $Z[a_i, b_{i,j}^{(k)}, c_{i,j}^{(k)}, d_{i,j}^{(k)}]$  where  $a_i, b_{i,j}^{(k)}, c_{i,j}^{(k)}$  and  $d_{i,j}^{(k)}$  are variables and  $I$  the ideal of relations that appear in (i) ~ (vii) of (4.1).

Then we get a universal symplectic formal system over  $S/I$ .

We denote  $\Gamma_{\text{univ}}$  as this system over  $S/I$  and do  $\text{LMSp}$  as  $(S/I)_{\Gamma_{\text{univ}}}$ .

Next we want to construct a symplectic formal system over  $H_*(\text{MSp})$ .

For simplicity, we denote  $f(x)$  and  $\bar{f}(x)$  as  $h(-x^2)$  and  $\frac{1}{2} \frac{d}{dx} h(-x^2)$

$H_*(\text{MSp})[[x]]$  where  $h(-x^2)$  is as previous.

We denote a symplectic formal system  $\Gamma_H$  by setting,

$$E^H(f(x)) = (\bar{f}(x))^2,$$

$$F_k^H(f(x), \bar{f}(x), f(y), \bar{f}(y)) = (f(x+y))^k \quad \text{and}$$

$$G_k^H(f(x), \bar{f}(x), f(y), \bar{f}(y)) = \bar{f}(x+y) \cdot (f(x+y))^{k-1} \quad \text{for } k \geq 1.$$

Then the relations (i) ~ (vii) except (v) are almost trivial.

Proposition 4.4. In  $\Gamma_H$ , differential relation holds.

We have a ring homomorphism  $\theta': \text{LMSp} \rightarrow H_*(\text{MSp})_{\Gamma_H}$  by the universality.

Theorem.  $\text{Im}(\theta') \subseteq \text{Im}(\text{Hurewicz homomorphism} : \text{MSp}_* \rightarrow H_*(\text{MSp}))$ .