

ON IMBEDDINGS OF SOME SEPARABLE EXTENSIONS

IN GALOIS EXTENSIONS

by

Takasi NAGAHARA

Department of Mathematics, Okayama University

Throughout B will mean a ring with identity element 1 , and all ring extensions of B will be assumed to have the (common) identity element 1 . As a sequel of imbedding theorems in field theory, we have the following problem: *If a ring extension A/B is f. g. projective and separable then can A/B be imbedded in a Galois extension N/B ?*

First, we consider a commutative ring A which is a ring extension of B . By [1], M. Auslander and O. Goldman proved that A/B is f. g. free and separable then A/B can be imbedded in a Galois extension. Moreover, in [8], O. E. Villamayor proved that this result is also true for any projective separable extension with projective rank. However, more generally, there holds that *any projective separable extension can be imbedded in a Galois extension* ([6]).

Now, let $B[X]$ be a polynomial ring over a commutative ring B . Then, if a monic $f \in B[X]$ is separable then the discriminant $\delta(f)$ is invertible in B , and conversely; when this is the case, the factor ring $B[X]/(f)$ (which is a ring extension of B) can be imbedded in a Galois extension N/B such that $N = B[x_1, \dots, x_n]$, $B[x_1] = B[X]/(f)$, and $f = (X - x_1) \cdots (X - x_n)$, which will be called *a splitting ring of f* ([5]). In case B is irreducible, there exists an (irreducible) polynomial closure

$C(B)$ so that any separable polynomial in $B[X]$ has a splitting ring in $C(B)$, and $C(B)$ is generated by the subrings of $C(B)$ which are splitting rings of separable polynomials in $B[X]$ ([2], [4]).

Next, we consider a non commutative ring B and a skew polynomial ring $B[X;\rho]$ of ρ -automorphisms type. Then, for a polynomial f in $B[X;\rho]$ of degree 2, f is Galois if and only if f is $\tilde{\rho}$ -separable ([3], [7], [8]). Moreover, if $f \in B[X;\rho]$ is $\tilde{\rho}$ -separable then $B[X;\rho]/(f)$ can be imbedded in a Galois extension which is a splitting ring of f . Further, in case B is simple, we can discuss (simple) splitting rings of separable polynomials in $B[X;\rho]$ and (simple) polynomial closures $C(B;\rho)$ in various ways. Those results will be seen later in a paper (to appear).

References

- [1] M. Auslander and O. Goldman: The Brauer group of a commutative ring, Trans. Amer. Math. Soc. 97 (1960), 367-409.
- [2] F. DeMeyer: Separable polynomials over a commutative ring, Rocky Mt. J. Math. 2 (1972), 299-310.
- [3] S. Ikehata: On separable polynomials and Frobenius polynomials in skew polynomial rings, Math. J. Okayama Univ. 22 (1980), 115-129.
- [4] G. J. Janusz: Separable algebras over a commutative rings, Trans. Amer. Math. Soc. 122 (1966), 461-479.
- [5] T. Nagahara: On separable polynomials over a commutative ring II, Math. J. Okayama Univ. 15 (1972), 149-162.
- [6] T. Nagahara: Imbeddings of some separable extensions in Galois extensions II, Math. J. Okayama Univ. 18 (1976), 189-194.
- [7] T. Nagahara: On separable polynomials of degree 2 in skew polynomial rings, Math. J. Okayama Univ. 19 (1976), 64-95.
- [8] T. Nagahara: On separable polynomials of degree 2 in skew polynomial rings II, Math. J. Okayama Univ. 21 (1979), 167-177.
- [9] O. E. Villamayor: Separable algebras and Galois extensions, Osaka J. Math. 4 (1967), 161-171.