

Normal affine subalgebras of a polynomial ring

阪大理 宮西正直 (Masayoshi Miyanishi)

Introduction. Let  $R := \mathbb{C}[x_1, \dots, x_n]$  be a polynomial ring in  $n$ -variables over the complex number field  $\mathbb{C}$ . A cofinite subalgebra of  $R$  is a  $\mathbb{C}$ -subalgebra  $A$  of  $R$  such that  $R$  is an  $A$ -module of finite type. We consider exclusively a normal cofinite subalgebra  $A$  of  $R$ . The following results are known by far:

1.  $A$  is finitely generated over  $\mathbb{C}$ , and all invertible elements of  $A$  are constants, i.e.,  $A^* = \mathbb{C}^*$ .

2. Let  $X$  be the normal affine variety defined by  $A$ . Then  $H_i(X; \mathbb{Z})$  is finite for all  $i \geq 1$ ,  $X$  is simply connected and  $\text{Pic}$  is trivial (Gurjar [3] and Kumar [4]). Therefore  $A$  is factorial if  $X$  is nonsingular.

3. If  $n = 2$  and  $A$  is regular,  $A$  is then a polynomial ring in two variables over  $\mathbb{C}$  (Miyanishi [6]).

4. Suppose  $n = 2$ . Then  $X$  has at worst quotient singularities (Brieskorn [1]). Moreover, if  $X$  is affine-ruled, i.e.,  $X$ , by definition, contains a non-empty Zariski open set of the form  $U_0 \times \mathbb{C}$ ,  $X$  has at worst cyclic quotient singularities (Miyanishi [6]).

We complement these results with the following:

THEOREM. Let  $A$  be a normal cofinite subalgebra of  $\mathbb{C}[x_1, x_2]$  and let  $X := \text{Spec } A$ . Then either  $X \simeq \mathbb{C}^2$  or  $X \simeq \mathbb{C}^2/G$ , where  $G$  is a small finite subgroup of  $\text{GL}(2, \mathbb{C})$ . If  $A$  is factorial,  $X$  is isomorphic to a hypersurface in  $\mathbb{C}^3$  defined by  $x_1^2 + x_2^3 + x_3^5 = 0$ .

The theorem holds true even if we replace the ground field  $\mathbb{C}$

by an algebraically closed field of characteristic  $p \neq 2, 3, 5$ . Moreover, the result is viewed as a global version of the following result of Brieskorn [1]:

Among two-dimensional normal singular analytic local rings, a factorial one is isomorphic to  $\mathbb{C}\{x, y, z\}/(x^2 + y^3 + z^5)$ .

1. Proof of Theorem: Nonsingular case.

Let  $X$  be a nonsingular algebraic surface defined over  $\mathbb{C}$ . Then there exists an open immersion of  $X$  into a nonsingular projective surface  $V$  such that  $D := V - X$  consists of nonsingular curves which cross each other normally. Let  $K_V$  be the canonical divisor and denote by the same letter  $D$  the reduced effective divisor such that  $\text{Supp } D = V - X$ . Then we say that  $X$  has (logarithmic) Kodaira dimension  $\kappa(X) = -\infty$  if  $|n(D + K_V)| = \emptyset$  for every  $n > 0$ . Then the property  $\kappa(X) = -\infty$  is independent of the choice of an immersion  $X \hookrightarrow V$ .

In proving the theorem, the following characterization of  $\mathbb{C}^2$  plays a crucial role:

Let  $X = \text{Spec } A$  be a two-dimensional affine surface defined over  $\mathbb{C}$ . Then  $X \cong \mathbb{C}^2$  if and only if the following three conditions are satisfied:

(i)  $A$  is factorial, (ii)  $A^* = \mathbb{C}^*$ , (iii)  $X$  is affine-ruled.  
When  $X$  is nonsingular, the condition (iii) is equivalent to (iii)'  $\kappa(X) = -\infty$ .

(See Miyanishi [5; 6].)

Let  $X$  now be the same as in the theorem. Suppose  $X$  is nonsingular. Then  $A := \Gamma(X, \mathcal{O}_X)$  is factorial by virtue of a result of Gurjar-Kumar, and  $A^* = \mathbb{C}^*$  because  $A$  is a  $\mathbb{C}$ -subalgebra of

$R := \mathbb{C}[x_1, x_2]$ . Moreover, since there is a finite morphism  $\theta: \mathbb{C}^2 \rightarrow X$ , we have  $\kappa(X) = -\infty$ . Then  $X \cong \mathbb{C}^2$  by virtue of the above-mentioned characterization of  $\mathbb{C}^2$ .

## 2. Proof of Theorem: Singular case.

We shall assume below that  $X$  is singular. Set

$X' := X - \text{Sing } X$ ,  $S = \mathbb{C}^2$ ,  $\theta: S \rightarrow X$  the given finite morphism,  $S' := \theta^{-1}(X')$ ,  $\theta' := \theta|_{S'}: S' \rightarrow X'$ ,  $q': Y' \rightarrow X'$  the topological universal covering space of  $X'$ .

Then  $\kappa(S') = -\infty$ , and  $\theta'$  factors as

$$\theta' : S' \xrightarrow{\pi'} Y' \xrightarrow{q'} X' .$$

Hence  $Y'$  is a nonsingular algebraic surface, and  $q'$  is a finite étale Galois covering with group  $G$ . Let

$$A := \Gamma(X', \underline{O}_{X'}) = \Gamma(X, \underline{O}_X),$$

$B :=$  the integral closure of  $A$  in the function field  $\mathbb{C}(Y')$ ,

$$R := \mathbb{C}[x_1, x_2] = \Gamma(S, \underline{O}_S),$$

$$Y := \text{Spec } B,$$

$\pi: S \rightarrow Y$  and  $q: Y \rightarrow X$ : the finite morphisms induced by the canonical inclusions  $B \subset R$  and  $A \subset B$ , respectively.

Then we know that:

- (i)  $A = R \cap \mathbb{C}(X)$  and  $B = R \cap \mathbb{C}(Y')$ ;
- (ii)  $Y$  is a normal affine surface defined over  $\mathbb{C}$  such that  $Y'$  is an open set of  $Y$  with  $\dim(Y - Y') \leq 0$ ;
- (iii)  $\theta = q \cdot \pi$ ,  $\pi' = \pi|_{S'}$ , and  $q' = q|_{Y'}$ ;
- (iv)  $G$  acts regularly on  $Y$ , and  $X \cong Y/G$ .

On the other hand,  $Y'$  is simply connected by the definition, Pic

is a torsion group because  $S'$  is a finite covering of  $Y'$ , and  $\dim(Y-Y') \leq 0$ . Therefore  $\text{Pic } Y' = (0)$ , and the divisor class group  $\text{Cl}(Y)$  is trivial, i.e.,  $B$  is factorial. Since  $B \subset R$ , we have  $B^* = \mathbb{C}^*$ . Hence if  $Y'$  is affine-ruled, so is  $Y$ , and  $Y \simeq \mathbb{C}^2$  by virtue of the characterization theorem of  $\mathbb{C}^2$ . The group  $G$  then becomes a finite subgroup of  $\text{Aut } \mathbb{C}^2 = \text{Aut } \mathbb{C}[x_1, x_2]$ , which is, up to conjugation, a finite subgroup of  $\text{GL}(2, \mathbb{C})$ . Let  $N$  be the normal subgroup of  $G$  consisting of all pseudo-reflections. Then  $\mathbb{C}^2/N$  is isomorphic to  $\mathbb{C}^2$ , and  $X \simeq (Y/N)/(G/N) \simeq \mathbb{C}^2/(G/N)$ . Hence we may assume that  $G$  is small, i.e.,  $G$  contains no pseudo-reflections.

Note that  $\kappa(Y') = -\infty$  because  $S'$  is a finite covering of  $Y'$  and  $\kappa(S') = -\infty$ . We shall show that  $Y'$  is affine-ruled. By reductio absurdum, we assume that  $Y'$  is not affine-ruled. Then we have the following:

THEOREM (Tsunoda-Miyayishi [8]). There exist a Zariski open set  $U$  of  $Y'$  and a proper birational morphism  $\phi : U \rightarrow Z$  from  $U$  onto a nonsingular algebraic surface  $Z$  defined over  $\mathbb{C}$  such that:

- (i) Either  $U = Y'$  or  $Y'-U$  has pure dimension 1;
- (ii)  $Z$  is a Platonic  $\mathbb{C}^*$ -fiber space.

A nonsingular algebraic surface  $Z$  is called a Platonic  $\mathbb{C}^*$ -fiber space if there exists a surjective morphism  $f : Z \rightarrow \mathbb{P}_{\mathbb{C}}^1$  such that general fibers of  $f$  are isomorphic to  $\mathbb{C}^*$  and that  $f$  has exactly three singular fibers  $\Delta_i = \mu_i \Gamma_i$  ( $i = 0, 1, 2$ ;  $\mu_0 \leq \mu_1 \leq \mu_2$ ) with  $\Gamma_i \simeq \mathbb{C}^*$ , where  $\{\mu_0, \mu_1, \mu_2\} = \{2, 2, n\}$  ( $n \geq 2$ ),  $\{2, 3, 3\}$ ,  $\{2, 3, 4\}$  or  $\{2, 3, 5\}$ .

Since  $U \subset Y' \subset Y$  and  $Y$  is affine,  $U$  does not contain any complete curve. Therefore  $\phi : U \rightarrow Z$  is an isomorphism. We claim that  $U = Y'$ . Otherwise, since  $Y' - U$  has pure dimension 1 and  $\text{Pic } Y' = (0)$ , there exists an element  $b$  of  $B = \Gamma(Y', \mathcal{O}_{Y'})$  such that  $\text{Supp } (b)_{0, Y'} = \text{Supp}(Y' - U)$ . Hence  $b$  is invertible on  $U$  and  $b \notin \mathbb{C}^*$ . Meanwhile, there exists a completion  $W$  of  $U$  such that  $W$  is a normal projective surface,  $W$  has at worst quotient singularities and  $W - U$  consists of two connected components, the one being a single irreducible curve and the other being a single quotient singular point (see Example 1 in the Section 3). Since  $(b)_W$  has support on  $W - U$ ,  $b$  must be a constant, i.e.,  $b \in \mathbb{C}^*$ . This is a contradiction. Hence  $U = Y'$ . In order to complete the proof of the first assertion of the theorem we make use of the following:

THEOREM (Miyanishi [7]; Fujita [2]). Let  $\tilde{U} \rightarrow U$  be the topological universal covering space of  $U$ . Then  $\tilde{U}$  is an affine-ruled nonsingular algebraic surface. Moreover we have


$$\pi_1(U) \cong \begin{cases} D_{2n} & \text{if } \{\mu_0, \mu_1, \mu_2\} = \{2, 2, n\} \\ A_4 & \text{if } \{\mu_0, \mu_1, \mu_2\} = \{2, 3, 3\} \\ S_4 & \text{if } \{\mu_0, \mu_1, \mu_2\} = \{2, 3, 4\} \\ A_5 & \text{if } \{\mu_0, \mu_1, \mu_2\} = \{2, 3, 5\} \end{cases},$$

where  $D_{2n}$  is a dihedral group of order  $2n$ , (see Example 2 of the Section 3).

However  $Y'$  is simply connected by the definition. This is apparently a contradiction. Thus  $Y'$  is affine-ruled, and we are done.

### 3. Examples.

(1) Let  $T$  be a hypersurface  $x_1^2 + x_2^3 + x_3^5 = 0$  in  $\mathbb{C}^3$  and let  $T'$  be the minimal resolution of the unique singular point  $P := (0,0,0)$  of  $T$ . Then  $T'$  is embedded into a nonsingular projective surface  $V$  in such a way that, in the configuration below, the top solid lines represent the exceptional curves arising from the minimal resolution of singularity at  $P$ , and the bottom solid lines represent the curves attached to  $T'$  to compactify the surface  $T'$ :




where "solid line" = a nonsingular rational curve, "broken line with weight  $-1$ " = an exceptional curve of the first kind and  $l$  is a fiber of  $f$ .


Moreover  $V$  has a structure of a  $\mathbb{P}^1$ -fibration  $f : V \rightarrow \mathbb{C}$  over  $\mathbb{C} \cong \mathbb{P}_{\mathbb{C}}^1$  such that  $f$  has three singular fibers  $F_i$  ( $i = 0, 1, 2$ ) and two cross-sections  $S_0, S_1$  as indicated in the configuration. Let  $U := T - \{P\}$ . Then  $U \cong V$  (all solid lines), and  $U$  is a Platonic  $\Gamma^*$ -fiber space with the triple  $\{2, 3, 5\}$ . The top solid lines contract down back to the quotient singular point  $P$ , and the bottom solid lines (sprouting from  $S_1$ ) contract down to three cyclic quotient singular points. With these contractions performed, we obtain a normal projective surface  $W$  such that  $T$  is an open set

of  $W$  and  $W-T$  consists of a single irreducible curve which is the proper transform of  $S_1$ .

(2) Consider a Platonic  $\mathbb{C}^*$ -fiber space  $U$  with the triple  $\{2,2,n\}$ . In general, it can be embedded into a nonsingular projective surface  $V$ , whose boundary  $V-U$  has the configuration as shown in the following picture:



Moreover,  $V$  has a structure of a  $\mathbb{P}^1$ -fibration  $f : V \rightarrow C$  over  $C \cong \mathbb{P}^1_{\mathbb{C}}$  for which  $S_0$  and  $S_1$  are cross-sections and  $F_0, F_1$  and  $F_2$  are the only singular fibers. Let  $D$  be the reduced effective divisor on  $V$  supported by all solid lines. Then  $D + K_V \sim \ell - (A_2 + B_2 + E_3)$ , where  $\ell$  is a fiber of  $f$ . Hence  $2(E_3 + D + K_V) \sim A_0 + A_1 + B_0 + B_1$ . This implies that there exists a finite double covering  $\alpha : \hat{V} \rightarrow V$  ramified only over  $A_0 + A_1 + B_0 + B_1$ . The surface  $\hat{V}$  has a  $\mathbb{P}^1$ -fibration  $\hat{f} : \hat{V} \rightarrow \hat{C} \cong \mathbb{P}^1_{\mathbb{C}}$  which is induced by the  $\mathbb{P}^1$ -fibration  $f : V \rightarrow C$  and has the configuration given in the next page. Indeed,  $\hat{V}$  is the normalization of  $V \times_C \hat{C}$ , where  $\hat{C} \rightarrow C$  is the double covering ramified over the points  $f(F_0)$  and  $f(F_1)$ .



Now contracting  $\hat{A}_0, \hat{A}_1, \hat{B}_0$  and  $\hat{B}_1$ , we are reduced to the case where  $\hat{f}$  has only two singular fibers of the same form as  $F_2$ . It is now a good exercise to show that  $\hat{V}$  - (all solid lines) is affine-ruled.

4. Proof of Theorem: The second assertion.

We start with the following situation:

$G \subset GL(2, \mathbb{C})$  : a small finite subgroup,

$X := \mathbb{C}^2/G$  : a singular normal affine surface,

$P :=$  the unique singular point of  $X$  which is the image of the point of origin  $(0,0)$  of  $\mathbb{C}^2$ ,

$A := \Gamma(X, \mathcal{O}_X)$ .

Then  $A$  is factorial if and only if  $\text{Pic}(X - \{P\}) = (0)$ . A line bundle  $L$  on  $X - \{P\}$  is constructed from a multiplicative character  $\chi$  of  $G$  in the following way: Let  $L$  be a line bundle on  $X - \{P\}$  and let  $\theta : \mathbb{C}^2 \rightarrow X$  be the (finite) quotient morphism. Then  $\theta^*L$  is a trivial line bundle on  $\mathbb{C}^2 - \{0\}$ . The action of  $G$  on  $\theta^*L$  is given by



$$(x, t) \in (\mathbb{C}^2 - \{0\}) \times \mathbb{C} \longmapsto ({}^g x, \chi(g, x)t) \in (\mathbb{C}^2 - \{0\}) \times \mathbb{C},$$

where  $g \in G$  and  $\chi(g, x) \in \mathbb{C}^*$ . Moreover, we have

$$\chi(gg'; x) = \chi(g; {}^{g'} x) \chi(g'; x) \quad \text{for } g, g' \in G.$$

If  $g \in G$  is fixed, then we have

$$\chi(g; x) \in \Gamma(\mathbb{C}^2 - \{0\}, \underline{0}^*) = \mathbb{C}[x_1, x_2]^* = \mathbb{C}^*.$$

Hence  $\chi(g, x)$  is independent of  $x$ . Write  $\chi(g) = \chi(g; x)$ . Then  $\chi : G \rightarrow \mathbb{C}^*$  is a multiplicative character. Conversely, a multiplicative character  $\chi$  of  $G$  defines a line bundle  $L_\chi := (\mathbb{C}^2 - \{0\}) \times \mathbb{C} / G$  with respect to the action of  $G$  as described above. Thus we have a 1 - 1 correspondence between

$$L \in \text{Pic}(X - \{P\}) \longleftrightarrow \chi \in \hat{G}.$$

Then we have:

$$\begin{aligned} \text{Pic}(X - \{P\}) = (0) &\iff \hat{G} = \{1\} \\ &\iff G \text{ is a binary icosahedral} \\ &\quad \text{group in } \text{SL}(2, \mathbb{C}) \\ &\iff X \text{ is isomorphic to a} \\ &\quad \text{hypersurface } x_1^2 + x_2^3 + x_3^5 = 0 \\ &\quad \text{in } \mathbb{C}^3. \end{aligned}$$

This completes the proof of the theorem.

#### References

- [1] E. Brieskorn: Rationale Singularitäten komplexer Flächen.  
Inventiones math. 4 (1968), 336-358.
- [2] T. Fujita: On the topology of non-complete algebraic surfaces  
J. Fac. Sci. Univ. of Tokyo, Sec. IA 29 (1982), 503-566.

- [3] R. V. Gurjar: Topology of affine varieties dominated by an affine space. *Inventiones math.* 59 (1980), 221-225.
- [4] S. Kumar: Varieties dominated by  $\mathbb{C}^n$ . Preprint.
- [5] M. Miyanishi: Lectures on curves on rational and unirational surfaces. Tata Institute of Fundamental Research, Bombay.
- [6] M. Miyanishi: Non-complete algebraic surfaces. *Lecture Notes in Mathematics* 857, Springer, 1981.
- [7] M. Miyanishi: On the affine-ruledness of algebraic varieties. To appear in the *Proceedings of the France-Japan Conference in Algebraic Geometry (Tokyo : 1982)*.
- [8] S. Tsunoda and M. Miyanishi: On the structure of non-complete algebraic surfaces with logarithmic Kodaira dimension  $-\infty$  and with non-connected boundaries at infinity. Forthcoming.