

SPECTRAL THEORY ON UNIFORMLY DISTRIBUTED SEQUENCES

KENJI NAGASAKA

SHINSYU UNIVERSITY

1. INTRODUCTION.

We shall consider sequences of real numbers or of complex numbers. The difference between simply normality and normality is to consider the frequency of any kind of block  $B_k$ . For the sequence of real numbers modulo 1 the corresponding relation is like between uniform distributed sequences and completely uniformly distributed sequences.

Theory of the uniform distribution of sequences is a fruitful branch in mathematics where number theory, probability theory and harmonic analysis meet with. Taking a normal number  $x$  to base  $q$ , then the sequence  $(q^n x)$  is uniformly distributed modulo 1. Hence we start from a modification of uniformly distributed sequences modulo 1. Then we discuss the spectrum and the spectral measure of a given sequence. These notions are from Fourier analysis and shall explain certain properties of the sequence considered.

## 2. NORMAL SETS.

Let  $\Lambda = (\lambda_n)$  be an infinite sequence of real numbers.

A real number  $x$  is called  $\Lambda$ -normal if the sequence

$x\Lambda = (x\lambda_n)$  is uniformly distributed modulo 1. Taking

$\Lambda = (q^n)$ , then a  $\Lambda$ -normal number coincides with a usual normal

number to base  $q$ . The set of  $\Lambda$ -normal numbers is denoted by

$B(\Lambda)$ . A set  $E$  of real numbers is called normal if there

exists a sequence  $\Lambda$  such that  $E = B(\Lambda)$ .

Mendès France and his colleagues collaborated on this notion (7), (8), (9) and (10) just after his arrival to Bordeaux University. Examples of normal sets are the followings:

- i) the complement of any real algebraic field of finite degree;
- ii)  $m\mathbb{Z}^*$  with positive integer  $m$ ;
- iii)  $\mathbb{Q}^*$ .

This series of papers ended with Gérard Rauzy's paper which gave a complete characterization of the normal set:

THEOREM (RAUZY). A set  $B$  is normal if and only if

- i)  $0 \notin B$ ;
- ii) for every positive integer  $q$ ,  $qB \subset B$ ;
- iii) there exist continuous real functions  $f_1, f_2, \dots$  such

that  $\lim_{n \rightarrow \infty} f_n(x) = 0$  exactly for those  $x$ 's which are elements of  $B$ .

The remaining unsolved (?) problems on normal sets are as follows:

- 1) Characterize the number  $\mu$  such that  $B + \mu = B$  implies the normality of  $B$ . ( Every positive integer satisfies this property. )
- 2) A finite union of normal sets is normal or not.

In early 70's Mendès France considered the "depth" of a rational number using continued fraction expansions (13) and (14) and mentioned a relation to the formal language theory (16) and corresponding results on real algebraic numbers of second degree ( remind Lagrange's theorem on periodic continued fractions ) were proved by Henri Cohen and also by Peysant Leroux. Another related his paper to continued fractions is (24).

### 3. SPECTRUMS AND SPECTRAL MEASURES.

In analyzing a sequence of statistical data, the autocorrelation functions are the main powerful tool. This idea is back to Nobert Wiener in 1927 for time series analysis. The progress of Fourier analysis allows us to apply this idea on

arithmetical sequences.

From now on we considered arithmetical sequences

$\alpha = (\alpha(n))$  of complex numbers. The correlation function of

with lag  $h$  is defined by

$$\gamma_{\alpha}(h) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \alpha(n) \overline{\alpha(n+h)},$$

where  $\bar{z}$  is the complex conjugate of  $z$ . M. Bass defined a

pseudo-random sequence  $\alpha = (\alpha(n))$  satisfying following three

conditions:

- i) For each  $h$ ,  $\gamma_{\alpha}(h)$  is determined;
- ii)  $\gamma_{\alpha}(0) \neq 0$ ;
- iii)  $\lim_{h \rightarrow \infty} \gamma_{\alpha}(h) = 0$ .

Mendès France proved the next theorem in his thesis (5):

THEOREM ( MENDES FRANCE ). If a real number  $x$  is normal to the base  $r$ , then the sequence

$$n \mapsto r_{|n|} (x) = \exp\left(\frac{2i\pi}{r} [x \cdot r^{|n|}]\right)$$

is pseudo-random and its correlation function is

$$\gamma(h) = \begin{cases} 1 & h = 0 \\ 0 & \text{otherwise} \end{cases}.$$

A normal number is a model of a random sequence, hence this

theorem seems quite natural.

By replacing iii) by

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{h=1}^N |\gamma(h)|^2 = 0,$$

we call  $\alpha = (\alpha(n))$  is pseudo-random in the sense of

Bertrandias. From now on we used "pseudo-random" in the sense of

Bertrandias.

Let us introduce another notion: spectrum of a sequence

$\alpha = (\alpha(n))$ :

$$\text{sp}(\alpha) = \{ \lambda \in [0,1); \lim_{N \rightarrow \infty} \sup \frac{1}{N} \left| \sum_{k=1}^N \alpha(k) \exp(-2i\pi\lambda k) \right| < \infty \}$$

Van der Corput difference theorem said that pseudo-random

sequence has empty spectrum. The converse is not in general

true. But in certain class of sequences, these two notions are

equivalent. Indeed, for any  $q$ -multiplicative sequence empty

spectrum implies pseudo-random (19). An arithmetical function

$\alpha$  is called  $q$ -multiplicative if

$$\alpha(aq^k + b) = \alpha(aq^k) \alpha(b)$$

and  $\alpha(0) = 1$  where  $q$  is a given integer larger than 1. The

typical example of  $q$ -multiplicative sequences is the sum of

digits  $s_q(n)$ : a positive integer  $n$  is written  $q$ -adically

$$n = \sum_{k=0} a_k \cdot q^k$$

then

$$s_q(n) = \sum_{k=0}^n a_k.$$

From the Bochner-Herglotz representation theorem, there exists a bounded positive measure  $\Lambda_\alpha$  such that

$$\gamma_\alpha(h) = \int_{\mathbb{R}/\mathbb{Z}} \exp(2i\pi hx) d\Lambda_\alpha(x).$$

This measure  $\Lambda_\alpha$  is called spectral measure of a given sequence  $\alpha$ , and we can describe certain properties of the sequence  $\alpha$  by means of its spectral measure (20).

If a  $q$ -multiplicative sequence is almost periodic, then its spectral measure is atomic.

If a  $q$ -multiplicative sequence is pseudo-random, then its spectral measure is singular.

Back to the sum of digits. Let us define

$$\xi(n) = \exp(2i\pi cs_q(n)),$$

where  $(q-1)c \notin \mathbb{Z}$ . The spectral measure  $\Lambda_\xi$  is continuous but singular with respect to Lebesgue measure. Let  $p$  and  $q$  be two relatively prime integers larger than 2, and

$$\alpha(n) = \exp(2i\pi as_p(n)), \quad \beta(n) = \exp(2i\pi bs_q(n)),$$

where  $a$  and  $b$  are integers satisfying

$$(p-1)a \notin \mathbb{Z}, \quad (q-1)b \notin \mathbb{Z}.$$

Then  $\Lambda_\alpha$  and  $\Lambda_\beta$  are singular to each other, proved by Kamae.

This is a kind of reply to the Steinhaus problem 144 stated in my report "normal numbers and dimension".

Finally I remark that a series of papers on spectral measures of arithmetical sequences have been produced by Coquet.

LIST OF PUBLICATIONS OF MICHEL MENDES FRANCE

- ( 1 ) Nombres normaux et fonctions pseudo-aléatoires, Ann.Inst. Fourier,13,91-104(1963).
- ( 2 ) Representation des réels, C.R.Acad.Sci,Paris,258,4643-4645 (1964).
- ( 3 ) Dimension de Hausdorff. Application aux nombres normaux, Séminaire Delange-Pisot:1963/1964,Théorie des nombres,Exp.6.
- ( 4 ) A set of nonnormal numbers, Pacific J.Math.,1165-1170 (1965).
- ( 5 ) Nombres normaux. Applications aux fonctions pseudo-aléatoires, J.Analyse Math.,20,1-56(1967).
- ( 6 ) Deux remarques concernant l'équirépartition des suites, Acta Arith.,14,163-167(1968).
- ( 7 ) Nombres transcendants et ensembles normaux, Acta Arith.,15, 189-192(1969). (Séminaire Delange-Pisot-Poitou:1967/68, Théorie des Nombres,Exp.16.)
- ( 8 ) La réunion des ensembles normaux, J.Number Th.,2,345-351 (1970).
- ( 9 ) DRESS,F. et MENDES FRANCE,M.: Caractérisation des ensembles dans  $\mathbb{Z}$ , Acta Arith.,17,115-120(1970). (Séminaire Delange-Pisot-Poitou:1968/69,Théorie des nombres,Exp.17. Erratum: Séminaire Delange-Pisot-Poitou:1969/70,Théorie des nombres, Exp.24.)

- (10) LESCA, J., et MENDES FRANCE, M.: Ensembles normaux, Acta Arith., 17, 273-282 (1970).
- (11) Suites et sous-suites équiréparties modulo 1, Journées Arithmétiques Françaises (Univ. Provence, 1971).
- (12) Les suites à spectre vide et la répartition (mod.1), J. Number Th., 5, 1-15 (1973).
- (13) Sur les fractions continues limitées, Acta Arith., 23, 207-215 (1973).
- (14) The depth of a rational number, Coll. Math. Soc. Janos Bolyai, 1974, Debrecen, North-Holland, Amsterdam.
- (15) DABOUSSI, H., and MENDES FRANCE, M.: Spectrum, almost periodicity and equidistribution modulo 1, Stud. Sci. Math. Hungar., 9, 173-180 (1974).
- (16) Fractions continues limitées et théorie des langages, Séminaire Delange-Pisot-Poitou: 1974/75, Théorie des nombres, Exp. 9.
- (17) A characterization of Pisot numbers, Mathematika (1975).
- (18) Hasard et déterministe, Séminaire Théorie des Nombres, Bordeaux: 1975/76.
- (19) COQUET, J., et MENDES FRANCE, M.: Suites à spectre vide et suites pseudo-aléatoires, Acta Arith., 32, 100-106 (1977).
- (20) COQUET, J., KAMAE, T., et MENDES FRANCE, M.: Sur la mesure spectrale de certaines suites arithmétiques, Bull. Soc. Math. France, 105, 369-384 (1977).
- (21) KAMAE, T., and MENDES FRANCE, M.: Van der Corput's difference theorem, Israel J. Math., 31, 335-342 (1978).
- (22) Apéry, irrationalité de  $\zeta(3)$ , Les Recherches (1978).



- (23) Algebraic numbers and automata theory, Proc. Queen's Univ., Number Th. Conf., Kingston Ontario (1979) Ed. Ribenboim, Exp. 54, 79-89.
- (24) CUSICK, T.W., and MENDES FRANCE, M.: The Lagrange spectrum of a set, Acta Arith., 34, 287-293 (1979).
- (25) Nombres algébriques et théorie des automates, l'Enseignement Math., 26, 193-199 (1980).
- (26) CHRISTOL, G., KAMAE, T., MENDES FRANCE, M., et RAUZY, G.: Suites algébriques, automates et substitutions, Bull. Soc. Math. France, 108, 401-419 (1980).
- (27) Principes de la symétrie perturbée, Séminaire Delange-Pisot-Poitou: 1979/80, Théorie des nombres, Exp. 15, Birkhäuser, Boston-Basel-Stuttgart, 77-98 (1981).
- (28) de l'arbre de Leonardo da Vinci à la théorie de la dimension, Revue du Palais de la Découverte, 52-60 (1981).
- (29) MENDES FRANCE, M. et TANENBAUM, G.: Dimension des courbes planes, papiers pliés et suites de Rudin-Shapiro, Bull. Soc. Math. France, 109, 207-219 (1981).
- (30) MENDES FRANCE, M. and van der POORTEN, A.J.: Arithmetic and analytic properties of paper folding sequences, Bull. Austral. Math. Soc., 24, 123-13K (1981).
- (31) DEKKING, F.M., and MENDES FRANCE, M.: Uniform distribution modulo one: a geometrical viewpoint, J. reine und Angew. Math., 329, 143-153 (1981).
- (32) Paper folding, space filling curves and Rudin-Shapiro sequences, Contemporary Math., 9, 85-95 (1982).
- (33) BLANCHARD, A., et MENDES FRANCE, M.: Symétrie et transcendance, Bull. Soc. Math. France, 110 (1982).
- (34) DEKKING, F.M., MENDES FRANCE, M. and van der POORTEN, A.J.: Folds!, Math. Intelligencer, 4, 130-138 (1982). - 9 - FIN.