

PROBLEMS RELATED TO BASES

MICHEL MENDES FRANCE

UNIVERSITE DE BORDEAUX I

I. Let θ and τ be two integers ≥ 2 . Define

$B(\theta)$ = set of real numbers normal to base θ

$B(\tau)$ = set of real numbers normal to base τ .

It is wellknown that

$B(\theta) = \{ x \in \mathbb{R} / (x\theta^n) \text{ is equidistributed (mod 1) } \}$

and that

$$B(\theta) = B(\tau) \Leftrightarrow \frac{\log \theta}{\log \tau} \in \mathbb{Q} .$$

Suppose now θ and τ are real numbers > 1 , not

necessarily integers.

PROBLEM 1: Is it true that

$$B(\theta) = B(\tau) \Leftrightarrow \frac{\log \theta}{\log \tau} \in \mathbb{Q} ?$$

The answer is surely NO! Indeed, it seems very plausible that

$\sqrt{2}$ is normal to base 2: $\sqrt{2} \in B(2)$. On the other hand

obviously $\sqrt{2} \notin B(\sqrt{2})$ because one term out of two in the sequence

$\sqrt{2}(\sqrt{2})^n$ is 0 (mod 1). So

$$B(2) \neq B(\sqrt{2}) \text{ inspite of the fact } \frac{\log 2}{\log \sqrt{2}} \in \mathbb{Q} .$$

I conjecture:

If θ and τ are Pisot numbers, then

$$B(\theta) = B(\tau) \quad \frac{\log \theta}{\log \tau} \in \mathbb{Q} .$$

(See A.Bertrand (2)).

II. Let $\theta > 2$ and consider the Cantor set $C(\theta)$

$$\begin{array}{c} \text{-----} \text{-----} \text{-----} \\ | \quad \quad \quad | \quad \quad \quad | \\ 0 \quad \quad \frac{1}{\theta} \quad \quad 1 - \frac{1}{\theta} \quad \quad 1 \end{array} \quad \begin{array}{l} \text{(the usual Cantor set} \\ \text{is } C(3) \text{) .} \end{array}$$

Then $C(\theta)$ is the set of real numbers

$$x = (\theta - 1) \sum_{n=1}^{\infty} \frac{\epsilon_n}{\theta^n}, \quad \epsilon_n = 0 \text{ or } 1 .$$

Obviously, for all integer $\theta \geq 3$, $B(\theta) \cap C(\theta) = \emptyset$.

PROBLEM 2: Let μ_θ be the "canonical measure" on $C(\theta)$ (I do not suppose that $\theta \in \mathbb{N}$). μ_θ is the measure defined on the set of infinite 0,1-sequences. I can prove that for μ_θ -almost all $x \in C(\theta)$, x is not in $B(\theta)$. In other words

$$\mu_\theta (B(\theta) \cap C(\theta)) = 0 .$$

Is it true that $B(\theta) \cap C(\theta) = \emptyset$?

I can also prove that for all $x \in C(\theta)$ there exists a nonzero algebraic integer $\alpha \in \mathbb{Q}(\theta)$ such that

$$\alpha x \notin B(\theta) .$$

If this integer α was known to be rational then the conjecture

$B(\theta) \wedge C(\theta) = \emptyset$ would be true.

On the other hand, if θ is not assumed to be a Pisot number, results seem different. I can prove that for almost all real $\theta > 1$,

$$\mu_{\theta}(B(\theta) \wedge C(\theta)) = 1.$$

(See my thesis (8), chapter 2).

III. Let $s_2(n)$ be the sum of the binary digits of n and $s_3(n)$ the sum of the digits of n in base 3. Straus and Senge proved (12):

$$\forall A \geq 1 \quad \forall B \geq 1 \quad \{ n / s_2(n) < A \text{ and } s_3(n) < B \}$$

is finite.

Gelfond (6) asked the 3 following questions:

QUESTION 1: Suppose $(b-1, 2) = (b'-1, 3) = 1$. Let a and a' be given. Show that the set

$$\{ n \in \mathbb{N} / s_2(n) \equiv a \pmod{b} \text{ and } s_3(n) \equiv a' \pmod{b'} \}$$

has density equal to $1/bb'$.

QUESTION 2: Show that (p prime)

$$\{ p \leq x / s_2(p) \equiv a \pmod{b} \} \sim \frac{1}{b} \pi(x).$$


QUESTION 3: Show that

$$\{ n \in \mathbb{N} / s_2(n^2) \equiv a \pmod{b} \}$$

has density $1/b$.

Question 1 was answered affirmatively by J.Besineau a student of mine (published in Acta Arith. (3)). Question 2 and 3 are still open. Question 2 should not be too difficult. As for Question 3, I believe it to be very deep for the following reason:

The sequence $s_2(n) \pmod{2}$ is the famous Morse sequence generated by automaton



A_0 is the initial state. The automaton works as follows. Feed in the integer n written as a string of 1's and 0's . And as an output read the index i of A_i .

Because $s_2(n) \pmod{2}$ is autonomous-generated, density type properties are easily established. For example

$$\text{density } \{ n / s_2(n) \equiv 0 \pmod{2} \} = 1/2 .$$

Now, J.P.Allouche (1), another student of mine proved that

$$n \mapsto s_2(n^2) \pmod{2}$$

is not generated by automaton. The sequence must then be quite complex.

IV. The following is a wellknown open problem. Define

$$T(n) = \begin{cases} \frac{1}{2}n & \text{if } n \text{ is even} \\ 3n + 1 & \text{if } n \text{ is odd} \end{cases} .$$

To prove that $\forall n, \exists k$ such that $T^{(k)}(n) = 1$. It seems quite clear (??) that this problem is related to the representation of integers in both bases 2 and 3, or what seems to amount to the same thing, to the distribution (mod 1) of $(\frac{3}{2})^n$.

So we are led to the other famous problem:

Is it true that $(\frac{3}{2})^n$ is dense (mod 1) ?

If so, then the following is obviously true. Define $\psi(x)$ as

the "depth" of the rational number x :

$$x = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\dots + \frac{1}{a_{\psi(x)}}}}}$$

If $(\frac{3}{2})^n$ is dense (mod 1), then

$$\limsup_{n \rightarrow \infty} \psi\left(\left(\frac{3}{2}\right)^n\right) = +\infty .$$

G.Choquet (4) proved this in a series of Comptes Rendus a l'Acad.

Sc. 1981-82 . His method was dynamical systems. Pourchet (11)

had already proved that actually

$$\lim_{n \rightarrow \infty} \psi\left(\left(\frac{3}{2}\right)^n\right) = +\infty .$$

(Details are given in (14)).

Heilbronn (7), Tonkov (13), Porter (10) have proved:

$$\frac{1}{\phi(q)} \sum_{\substack{a \leq q \\ (a,q)=1}} \psi\left(\frac{a}{q}\right) \sim \frac{12}{\pi^2} \log 2 \cdot \log q \quad (q \rightarrow \infty)$$

and J.D.Dixon (5) : for all $a < q \leq x$

$$\left| \psi\left(\frac{a}{q}\right) - \frac{12}{\pi^2} \log 2 \cdot \log q \right| < \sqrt{\log q}$$

with the exception of at most $o(x^2)$ a/q .

This leads to the following conjecture:

$$\lim_{n \rightarrow \infty} \frac{1}{n} \psi\left(\left(\frac{a}{q}\right)^n\right) = \frac{12}{\pi^2} \log 2 \cdot \log b$$

where $b = \min \{a, q\}$.

V. Back to Pisot numbers:

Let $\theta > 1$ be a real algebraic number. Define

$$E(x) = \{ x, x\theta, x\theta^2, x\theta^3, \dots \}$$

and let $E'(x)$ denote the set of limit points of $E(x)$, $E''(x)$

the set of limit points of $E'(x), \dots$ Pisot (9) shows that if

$E''(x) = \emptyset$ then θ is a Pisot number and $x \in \mathbb{Q}(\theta)$.

CONJECTURE: If there exists a $v \geq 2$ such that $E^{(v)}(x) = \emptyset$,

then θ is a Pisot number and $x \in \mathbb{Q}(\theta)$.

If I knew how to prove this, I would then have another proof of

Choquet's result.

REFERENCES

- (1) J.P.ALLOUCHE: Somme des chiffres et transcendance, Bull.Soc. Math.France,110,1982.
- (2) A.BERTRAND: Développements en base de Pisot et répartition modulo 1 de la suite (x_{θ^n}) , Thèse présentée à Université Bordeaux I,1981.
- (3) J.BESINEAU: Indépendance statistique d'ensembles liés à la fonction "somme des chiffres", Acta Arith.,20,1972,401-416.
- (4) G.CHOQUET: Les fermés $(3/2)$ -stables de T ; structures des fermés dénombrables;applications arithmétiques, Copmtes Rendus Acad.Sc.Paris,291,1980,239-244.
- (5) J.D.DIXON: The number of steps in the Euclidean algorithm, Jour.Numb.Th.,2,1970,414-422. (See also Amer.Math.Monthly, 78,1971,374-376).
- (6) A.O.GELFOND: Sur les nombres qui ont des propriétés additives et multiplicatives données, Acta Arith.,13,1968, 259-265.
- (7) H.HEILBRONN: On the average length of a class of finite continued fractions, Abhandlungen aus Zahlentheorie und Analysis, Berlin,1968,87-96,edited by P.Turán.
- (8) M.MENDES FRANCE: Nombres normaux. Applications aux fonctions pseudo-aléatoires, Jour.Analyse Math.,20,1967,1-56.
- (9) CH.PISOT:
- (10) J.W.PORTER: On a theorem of Heilbronn, Mathematika,22,1975, 20-28.
- (11) Y.POURCHET: Private communication c.1970.

- (12) H.G.SENGE,E.G.STRAUS: PV numbers and sets of multiplicity,
Periodica Math.Hungar.,3,1973,93-100.
- (13) T.TONKOV: On the average length of finite continued
fractions, Acta Arith.,26,1974,47-57.
- (14) A.van der POORTEN: Some problems of recurrent interest,
Colloquia Math.Soc.Janos Bolyai,34,Topics in classical
number theory (Budapest 1981),North Holland (to appear).