

Tomographic Reconstruction Problem

in

Hilbert Spaces

ISAMU SUZUKI (鈴木 勇)

Medical Engineering Division

Tokyo Metropolitan Institute of Medical Science

Section 1.

In this note we consider the problem of reconstructing a function from its projections. This problem has arisen in many fields such as medicine, astronomy and electron microscopy. Let B be a closed ball in R^n and we denote $L^2(B)$ a Hilbert space of all square integrable functions on B . Let $f \in L^2(B)$ and let $\{(R_{\omega_i} f)(\cdot)\}$ be a given finite set of hyperplane integrals where $\{\omega_1, \dots, \omega_k\}$ is a finite set of unit vectors in R^n and hyperplane integral $R_{\omega_i} f$ of f is defined as follows:

$$R_{\omega_i} f(p) = \int_{\langle x, \omega_i \rangle = p} f(x) dx$$

Under these notations the two main questions for this problem are stated as follows: [1]

- Q1. Estimate f by $\{(R_{\omega_i} f)(\cdot)\}$ and evaluate $\|f - \tilde{f}\|$, where \tilde{f} is an estimator of f ,
- Q2. for f and k are given, determine $\{\omega_i\}$ which optimizes the estimation.

Section 2.

If we have a complete set of hyperplane integrals, then the problem is nothing but an inversion theorem of Radon transform. In our case we consider an approximate reconstruction from finitely many projections. S.Helgason[2] and D.C.Solmon [3,4] showed several elegant results for Q1. Conversely we show some results for Q2 which give optimal projections. Let μ be the measure of $L^2(B)$ and let S be a covariance operator of μ with respect to f . Then the eigen vectors of S form an optimal coordinates system for approximating f . Furthermore this coordinates system has the entropy minimizing property.

References

- [1] I.SUZUKI, Computed Tomography and Radon transform.
Fourth Seminar on Applied Functional Analysis
1981,67-72.
- [2] S.Helgason, The Radon Transform. Progress in Mathematics
Birkhauser, Cambridge 1980.
- [3] D.C.Solmon, The X-ray Transform, J. Math. Anal. Appl.
31,1976,61-83.
- [4] _____, Practical and Mathematical Aspect of The
Problem of Reconstructing Objects From
Radiographs, Bulletin of the Amer. Math. Soc.
83,6,1977,1227-1268.