

Reidemeister Torsion of a Homology Lens Space

Tsuyoshi Sakai (北大 酒井健)

In my talk, I announced a formula giving the Reidemeister torsion of a homology lens space obtained by Dehn surgery on a knot in the three sphere associated to the universal abelian covering. And we applied it to get a generalization of Fukuhara's result in [2] that is a generalization of the classification of lens spaces.

After that, I found that essentially same result is already obtained by Turaev [4] in a more general setting. Here I will give the precise statement of result. For a proof of Theorem 1, the reader can refer the paper of Turaev [4] or a self-contained treatment in [3].

Statement of Results

Let  $k$  be a knot in the three sphere  $S^3$ . For coprime integers  $p > 0$  and  $q$ , let  $K = L(p, q; k)$  denote a 3-manifold obtained from  $S^3$  by Dehn surgery on  $k$  with coefficient  $p/q$ . Then  $H_1(K, Z)$  is isomorphic to the cyclic group of order  $p$  generated by a meridian loop of  $k$ . Let  $\tilde{K}$  denote the universal abelian covering of  $K$  with the covering transformation group  $\Pi$  and let  $T$  denote a generator of  $\Pi$  corresponding to the meridian loop of  $k$ . We assume that  $K$  is triangulated. Then the integral cellular chain group  $C_q(\tilde{K}; Z)$  can be considered as a  $Z\Pi$ -free module with the standard basis

determined by  $K$  which is well defined up to sign, and up to multiplication, by elements of  $\mathbb{T}$ .

Suppose a homomorphism  $h$  from  $\mathbb{T}$  to the field of complex numbers  $F$  is given that takes  $T$  into a  $p$ -th root of unity  $\tau (\neq 1)$ . Using  $h$ , we can form the chain complex

$$C_* = F \otimes_{\mathbb{T}} C_*(\tilde{K}; Z)$$

over  $F$ . Then  $C_q$  is a finite dimensional vector space over  $F$  with the standard basis determined by the basis for  $C_q(\tilde{K}; Z)$  above (thus determined by  $K$ ). For each  $q$ , let  $v_q$  denote the volume in  $C_q$  determined by this basis. Then

Theorem 1. Suppose that the Alexander polynomial of  $k$  is  $A(t)$ . Then  $C_*$  is acyclic if and only if  $A(\tau) \neq 0$ . Therefore, if  $A(\tau) \neq 0$ , the Reidemeister torsion  $\Delta_h(\tilde{K})$  is defined as  $\pm h(\mathbb{T}) v_0 v_1^{-1} v_2 v_3^{-1}$  and is equal to

$$\pm h(\mathbb{T}) A(\tau) (\tau^r - 1)^{-1} (\tau - 1)^{-1}$$

where  $r$  is determined by the congruence  $qr \equiv 1 \pmod{p}$ .

The following generalizes Theorem 2 in Fukuhara [2], which he proved by using EA-matrix, an invariant for closed orientable 3-manifold defined by Fukuhara and Kanno [1].

Theorem 2. Let  $k$  and  $k'$  be knots in  $S^3$  with trivial Alexander polynomials. Then  $L(p, q; k)$  is homeomorphic to  $L(p, q; k')$  only if  $\pm qq' \equiv 1 \pmod{p}$  or  $\pm q \equiv q' \pmod{p}$ .

The classification of lens spaces is the case that  $k$  and  $k'$

are trivial and Fukuhara's result is the case that  $k$  has the trivial Alexander polynomial and  $k'$  is trivial.

#### References

- [1] Fukuhara, S. and Kanno, J.: Extended Alexander matrices of 3-manifolds. Preprint. Tsuda College. Tokyo, Japan(1983).
- [2] Fukuhara, S.: Homology Lens Spaces obtained by Dehn surgeries on knots. Preprint. Tsuda College. Tokyo, Japan(1983).
- [3] Sakai, T.: Reidemeister Torsion of a Homology Lens Space. To appear in Kobe Journal of Math.
- [4] Turaev, V.G.: Reidemeister torsion and the Alexander polynomial. Math. USSR Sbornik 30, (1976), 221-237.