


A naive example of a proper  $\Pi_4^0$  set of reals

埼玉大理 花沢正純 (Masazumi Hanazawa)  
 法大工 田中尚夫 (Hisao Tanaka)

We show that a subset  $A$  of the product space  $2^\omega$  which is defined by

$A = \{ \alpha \in 2^\omega : \text{there are infinitely many integers } m \in \omega \text{ such that } \alpha \text{ contains infinitely many segments of the form } 10^m 1 \}$   
 is proper  $\Pi_4^0$ .

§1. This is concerned with Landweber's work on finite automaton theory. A finite automaton  $\mathcal{M}$  consists of four elements  $S, 2 = \{0, 1\}, s_0$ , and  $M$ .  $S$  is a finite set, whose elements are called states.  $s_0$  is an element of  $S$  and is called the starting states.  $M$  is a function from  $S \times 2$  into  $S$ , which is called a next state function. When a sequence  $\alpha \in 2^\omega$  was given (which is identified with a data tape), the machine  $\mathcal{M}$  runs as illustrated below.



The sequence  $\langle s_i \rangle_{i \in \omega}$  in the above is determined by  $\alpha$ , we denote it by  $\langle s_i(\alpha) \rangle_{i \in \omega}$ .

§2. Let  $D \subseteq S$ . Then  $\mathcal{M}$  accepts  $\alpha$  with respect to  $D$  if some member of  $D$  occurs in the sequence  $\langle s_i(\alpha) \rangle_{i \in \omega}$ , and rejects  $\alpha$  otherwise. This condition of acceptance of a sequence is of the most standard type. Some other conditions have been studied by Hartmanis, Stearns, Büchi, McNaughton, etc. ([1], [2], [3], [4], [5], [6])

Let  $D \subseteq S$ .

$\mathcal{M}$  1-accepts  $\alpha$  w.r.t.  $D$  if  $\exists i s_i(\alpha) \in D$ ,

$\mathcal{M}$  1'-accepts  $\alpha$  w.r.t.  $D$  if  $\forall i s_i(\alpha) \in D$ ,

$\mathcal{M}$  2-accepts  $\alpha$  w.r.t.  $D$  if  $\exists^\infty i s_i(\alpha) \in D$ ,

where  $\exists^\infty i$  means that there are infinitely many  $i$ 's such that...

Put  $\text{In}(\alpha) = \{s \in S : \exists^\infty i s = s_i(\alpha)\}$ . Let  $\mathcal{D} \subseteq \mathcal{P}(S)$ .

$\mathcal{M}$  2'-accepts  $\alpha$  w.r.t.  $\mathcal{D}$  if  $\exists D \in \mathcal{D} \text{In}(\alpha) \subseteq D$ ,

$\mathcal{M}$  3-accepts  $\alpha$  w.r.t.  $\mathcal{D}$  if  $\exists D \in \mathcal{D} \text{In}(\alpha) = D$ .

Let  $i$  stand for 1, 1', 2, 2', or 3.  $A \subseteq 2^\omega$  is  $i$ -definable if there is  $\mathcal{M}$  and  $D$  (or  $\mathcal{D}$ ) such that  $A = \{\alpha \in 2^\omega : \mathcal{M} \text{ } i\text{-accepts } \alpha \text{ w.r.t. } D \text{ (or } \mathcal{D})\}$ . The following are easily observed:

Proposition (Landweber [6]). (1) Every 1-definable set is  $\Sigma_1^0$ .

(2) Every 1'-definable set is  $\Pi_1^0$ .

(3) Every 2-definable set is  $\Pi_2^0$ .

(4) Every 2'-definable set is  $\Sigma_2^0$ .

(5) Every 3-definable set is  $\Delta_3^0$ .

These estimations are known to be proper by the following examples:

Examples (Landweber [6]). Put  $A^+ = \{\alpha \in 2^\omega : \text{in the sequence } \alpha, \text{ only a finite number of 1's occur}\}$ . Then:

(1)  $A^+$  is 2'-definable and in  $\Sigma_2^0 - \Pi_2^0$ .

(2)  $A^{+c}$  is 2-definable and in  $\Pi_2^0 - \Sigma_2^0$ .

(3)  $A^\# = \{\alpha : \alpha(0) = 0 \ \& \ \alpha \in A^+\} \cup \{\alpha : \alpha(0) = 1 \ \& \ \alpha \in A^{+c}\}$  is 3-definable and in  $\Delta_3^0 - (\Sigma_2^0 \cup \Pi_2^0)$ .

Problem (Landweber [6]). Find a N A T U R A L condition for acceptance which enables finite machines to define sets above  $\Delta_3^0$ .

To answer this question, it needs to find a set of reals in a higher hierarchy which is however easily handled by finite automaton. Unfortunately we could not find satisfactory examples of such sets in the literatures. So we give here such an example. Our answer to Landweber's problem is under improvemant, which will appear elsewhere in a satisfactory form.

§3. When a finite sequence  $x \in 2^{\omega}$  is an initial segment of a sequence  $y \in 2^{\omega} \cup 2^{\omega}$ , then we write  $x \prec y$ . The concatenation of two sequences  $x$  and  $y$  is denoted by  $xy$ . Our results are as follows:

Put  $A_0 = \{\alpha \in 2^{\omega} : \text{There is an } m \in \omega \text{ such that } 0\text{-blocks of length } m \text{ (} 0^m\text{-blocks) occur infinitely often in the sequence } \alpha \}$ , i.e.:

$$\alpha \in A_0 \text{ iff } (\exists m)(\exists^{\infty} x \in 2^{\omega}) x10^{m-1} \prec \alpha.$$

Theorem 1.  $A_0$  is in  $\Sigma_3^0 - \Pi_3^0$ .

Put  $A_1 = \{\alpha \in 2^{\omega} : \text{There are infinitely many integers } m \text{ such that } 0^m\text{-blocks occur infinitely often in the sequence } \alpha \}$ , i.e. :

$$\alpha \in A_1 \text{ iff } (\exists^{\infty} m) \text{ s. t. } (\exists^{\infty} x \in 2^{\omega}) x10^{m-1} \prec \alpha.$$

Theorem 2.  $A_1$  is in  $\Pi_4^0 - \Sigma_4^0$ .

Theorem 1 is essentially a part of Theorem 2. The rest of the paper is entirely devoted to the proof of Theorem 2.

§4. Proof of Theorem 2. Identifying the finite sequence  $0^{m-1}$  with  $m$  for each integer  $m$ , we can translate the subset  $A_1$  of  $2^{\omega}$  into the subset  $A$  of the space  $\omega^{\omega}$  as follows:

$$A = \{\alpha \in \omega^{\omega} : \exists^{\infty} m \text{ s.t. } \exists^{\infty} i \alpha(i) = m \}.$$

Thus we show that  $A$  is in  $\Pi_4^0 - \Sigma_4^0$ . Since  $A$  is clearly  $\Pi_4^0$  by the definition, it needs only to show  $A \notin \Sigma_4^0$ . For the purpose we use the following lemma:

Lemma 1 (Landweber [6], in modified form). For every subset  $X$  of  $\omega^\omega$ ,  $X$  is  $\Pi_2^0$  iff  $\exists B \subseteq \omega^\omega$  s.t.  $X = \widehat{B}$ , where  $\widehat{B}$  is the set of all  $\alpha \in \omega^\omega$  such that infinitely many segments of  $\alpha$  belongs to  $B$ , i.e. :  $\widehat{B} = \{\alpha \in \omega^\omega : (\exists^\infty y \in B) y \prec \alpha\}$ .

To the end of contradiction we suppose  $A \in \Sigma_4^0$ , equivalently  $A^c \in \Pi_4^0$ . Then by the lemma there is an indexed family of sets of finite sequences  $\{B_{1,i}\}_{1,i \in \omega}$  such that  $A^c = \bigcap_{i \in \omega} \bigcup_{j \in \omega} \widehat{B_{1,i}}$ .

Definition. Let  $x \prec y$  and  $J \subseteq \omega$ , and  $z$  be the sequence satisfying  $xz = y$ . If in the sequence  $z$ , only the elements of  $J$  occur, we say that  $y$  is a  $J$ -restricted extension of  $x$ , or shortly  $y$   $J$ -extends  $x$ , and write  $x \xrightarrow{J} y$ .

Notice that by the definition of  $A$ ,  $A^c$  is the set of all  $\alpha$  such that the set  $\{m : m \text{ occurs infinitely often in } \alpha\}$  is finite.

Lemma 2. If a family of sets of finite sequences  $\{C_i\}_{i \in \omega}$  satisfies that

$$A^c \subseteq \bigcup_{i \in \omega} \widehat{C_i}, \quad \text{then}$$

(\*) for every  $x$  in  $\omega^\omega$ , each infinite set  $I \subseteq \omega$ , and each finite set  $F \subseteq \omega$ , we can take  $x_0 \in \omega^\omega$  which  $F \cup I$ -extends  $x$ , a set  $C_i$  in  $\{C_i\}_{i \in \omega}$ , and an infinite set  $I_1 \subseteq I$  such that for every  $x_1$  which  $F \cup I_1$ -extends  $x_0$ , for every infinite set  $J \subseteq I_1$ , there is  $x^*$  which  $F \cup J$ -extends  $x_1$  such that  $x^* \in C_i$ ; in symbols,

$$(\forall x, F, I) \exists x_0 \xrightarrow{FI} x \exists I_1 \subseteq I \exists C_i \in \{C_i\}_{i \in \omega} \text{ such that}$$

$$\forall x_1 \xrightarrow{FI_1} x_0 \forall J \subseteq I_1 \exists x^* \xrightarrow{FJ} x_1 \quad x^* \in C_i.$$

Proof of Lemma 2. If not, we can obtain  $x$ ,  $F$ , and  $I$  such that:

$$(\#) (\forall x_0 \xrightarrow{FI} x \forall I_1 \subseteq I \forall C_i \in \{C_i\}_{i \in \omega}) \exists x_1 \xrightarrow{FI_1} x_0 \exists J \subseteq I_1$$

such that  $\forall x^* \xrightarrow{FJ} x_1 \quad x^* \text{ is not in } C_i$ .

We extend this  $x$  to  $\alpha \in \omega^\omega$  as follows:

$$x \xrightarrow{\text{FI}} x_0 \xrightarrow{\text{FJ}_0} x_1 \xrightarrow{\text{FJ}_1} \dots x_i \xrightarrow{\text{FJ}_i} \dots,$$

where infinite sets  $J_i$ ,  $i \in \omega$ , are chosen so that:

(\\$)  $I \supseteq J_0 \supseteq J_1 \supseteq J_2 \supseteq \dots \supseteq J_i \supseteq \dots$ , and  $J_n \cap \{0, 1, \dots, n\} = \emptyset$  for each  $n$ .

Moreover in addition  $x_i$  and  $J_i$ ,  $i \in \omega$ , are chosen so as to satisfy that: (%)<sub>0</sub> Every  $F \cup J_0$ -extension of  $x_0$  is not in  $C_0$ ,

(%)<sub>1</sub> Every  $F \cup J_1$ -extension of  $x_1$  is not in  $C_1$ ,

...

(%)<sub>i</sub> Every  $F \cup J_i$ -extension of  $x_i$  is not in  $C_i$ ,

...

This is clearly possible by the condition (#). Now put  $\alpha = \bigcup_{i \in \omega} x_i$ .

Then by the condition (%)<sub>i</sub>, any initial segment of  $\alpha$  which belongs to  $C_i$  must be shorter than  $x_i$  for each  $i$ , so  $\alpha \notin \bigcup_{i \in \omega} \widehat{C}_i$ .

But by the condition (\$), for every  $m$  in the co-finite set  $\omega - F$ , all occurrences of  $m$  in the sequence  $\alpha$  are in the initial segment  $x_m$ . This means that  $\alpha$  is in  $A^c$ ; a contradiction. Lemma 2 is thus proved.

In the present case,  $A^c \subseteq \bigcup_{i \in \omega} B_{l,i}$  for every  $l$ . Hence by this lemma, the following holds for every  $l$ :

(@)<sub>l</sub>  $(\forall x, F, I) \exists x_0 \xleftarrow{\text{FI}} x \exists B_{l,i} \in \{B_{l,i}\}_{i \in \omega} \exists I' \subseteq I$

such that  $(\forall x_1 \xleftarrow{\text{FI}} x_0)(\forall J \subseteq I') \exists x^* \xleftarrow{\text{FJ}} x_1$  s.t.  $x^* \in B_{l,i}$ .

Using this, we construct  $\alpha$  which is in  $\bigcap_{l \in \omega} \bigcup_{i \in \omega} B_{l,i}$  but not in  $A^c$ . Starting with null sequence  $\lambda$ , we repeat extensions of finite sequences in the following manner:

$$\lambda (= y_{-1}) \xrightarrow{\omega (= F_{-1} \cup I_{-1})} x_0 \xrightarrow{F_0 I_0} y_0 \xrightarrow{F_0 I_0} x_1 \xrightarrow{F_1 I_1} y_1 \xrightarrow{F_1 I_1} x_2 \dots,$$

where (1)  $F_0 \cup I_0 \supseteq F_1 \cup I_1 \supseteq F_2 \cup I_2 \supseteq \dots$  and

(2)  $F_0 \not\subseteq F_1 \not\subseteq F_2 \not\subseteq \dots$ .

I. Stage to define  $x_l \leftarrow_{F_{l-1} \cup I_{l-1}} y_{l-1}$ .

The extension  $x_l$  is taken together with a set  $I_l \subseteq I_{l-1}$  and a set

$B_{l,i_l} \in \{B_{l,i}\}_{i \in \omega}$  so that :

$$(\#)_l \quad (\forall x \leftarrow_{F_{l-1} \cup I_l} x_l) (\forall J \subseteq I_l) \exists y \in B_{l,i_l} \text{ s.t. } y \leftarrow_{F_{l-1} \cup J} x.$$

This is possible by the condition  $(@)_l$ . We put :

$$F_l = F_{l-1} \cup \{\min(I_l - F_{l-1})\}. \quad \text{Hence:}$$

$$(3) \quad F_{l-1} \cup I_l = F_l \cup I_l \supseteq \dots \supseteq F_m \cup I_m \supseteq \dots$$

Notice by the above conditions  $(\#)_l$ , (1), and (3) that:

$(\#)_l^m$  Every  $x_m$  with  $m > 1$  or its any  $F_m \cup I_m$ -extension can be  $F_m \cup I_m$ -extended to a member of  $B_{l,i_l}$ .

II. Stage to define  $y_l \leftarrow_{F_l \cup I_l} x_l$ .

By the help of the conditions  $(\#)_0^l, (\#)_1^l, \dots, (\#)_l^l$ , we repeat extensions  $l$  times starting from  $x_l$  as follows:

$$x_l \xrightarrow{F_l \cup I_l} y_{l,0} \xrightarrow{F_l \cup I_l} y_{l,1} \xrightarrow{F_l \cup I_l} \dots \xrightarrow{F_l \cup I_l} y_{l,l},$$

where  $y_{l,j}$  is a member of  $B_{j,i_j}$  for each  $j \leq l$ . Now we  $F_l \cup I_l$ -extend  $y_{l,l}$  to  $y_l$  so that all members of  $F_l$  occur in the newly added part; for example, put  $y_l = y_{l,l} k_0 k_1 \dots k_l$ , the concatenation of  $y_{l,l}$  and a list of all members of  $F_l = \{k_0, k_1, \dots, k_l\}$ .

The construction is completed. Put  $\alpha = \bigcup_{l \in \omega} x_l$ . Let  $j \in \omega$ .

Then for every  $l$  larger than  $j$ , the sequence  $y_{l,j}$  belongs to  $B_{j,i_j}$ ; this shows  $\alpha \in \widehat{B_{j,i_j}}$ . Thus  $\alpha \in \bigcap_{l \in \omega} \bigcup_{i \in \omega} \widehat{B_{l,i}}$ .

On the other hand, for each  $l \in \omega$ , the segment  $z_l$  that satisfies  $x_l z_l = y_l$  has an occurrence of each element of  $F_l$ . Since  $\{F_l\}_{l \in \omega}$  is strictly increasing, every element of infinite set  $\bigcup_{l \in \omega} F_l$  occurs infinitely often in the sequence  $\alpha$ , that means  $\alpha \notin A^c$ ; a contradiction. Theorem is thus proved.

## References

- [1] J. R. Büchi, On a decision method in restricted order arithmetic, Logic, Methodology and Philosophy of Science (Proc. 1960 Internat. Congr.); Stanford Univ. Press, Stanford, Cal., 1962.
- [2] J. R. Büchi and L. H. Landweber, Solving sequential conditions by finite state strategies, Trans. Amer. Math. Soc. 138 (1969), 295-311.
- [3] J. R. Büchi and L. H. Landweber, Definability in the monadic second order theory of successor, J. Symbolic Logic 34 (1969), 166-170.
- [4] J. Hartmanis and R. E. Stearns, Sets of numbers defined by finite automata, Amer. Math. Monthly 74 (1967), 539-542.
- [5] R. McNaughton, Testing and generating infinite sequences by a finite automaton, Information and Control 9 (1966), 521-530.
- [6] L. H. Landweber, Decision Problem for  $\omega$ -Automata, Math. Systems Theory 3 (1969) 376-384