

A REMARK ON α -CONVEX FUNCTIONS

M. NUNOKAWA AND S. HOSHINO

(布川 護, 星野晋一, 群馬大学)

ABSTRACT. Let α be real and suppose that $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ is analytic in the unit disk Δ . If $\operatorname{Re}[(1-\alpha)zf'(z)/f(z) + \alpha(1+zf''(z)/f'(z))] > 0$ for $z \in \Delta$, then $f(z)$ is said to be α -convex function. In this paper, we will show that if an α -convex function $f(z)$ satisfies certain conditions, then $f(z)$ is starlike of order at least $1/2$.

1. Introduction. Let A be the class of functions of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n$$

which are analytic in $\Delta = \{z : |z| < 1\}$.

A function $f(z)$ in A is said to be starlike iff

$$\operatorname{Re} \frac{zf'(z)}{f(z)} > 0 \quad \text{in } \Delta.$$

Further, a function $f(z)$ in A is said to be convex iff

$$1 + \operatorname{Re} \frac{zf''(z)}{f'(z)} > 0 \quad \text{in } \Delta.$$

It is well known that all convex functions are starlike of order at least $1/2$ [2,5].

On the other hand, a function $f(z)$ in A is said to be α -convex iff

$$\operatorname{Re} \left[(1-\alpha) \frac{zf'(z)}{f(z)} + \alpha \left(1 + \frac{zf''(z)}{f'(z)} \right) \right] > 0 \quad \text{in } \Delta.$$

Miller, Mocanu and Reade [4] proved the following theorem.

THEOREM A. If $f(z) \in A$ is α -convex in Δ , then $f(z)$ is starlike in Δ . Moreover, if $\alpha \geq 1$, then $f(z)$ is convex in Δ , and if $\alpha \leq -1$, then $1/f(1/z)$ is convex for $|z| > 1$.

It is the purpose of the present paper to partly improve THEOREM A.

2. Main theorem. We need the following lemma.

LEMMA 1. Let $w(z)$ be analytic in Δ , $w(0) = 0$. If $|w(z)|$ attains its maximum value on the circle $|z|=r < 1$ at a point z_0 , then we can write

$$z_0 w'(z_0) = kw(z_0)$$

where k is a real number and $k \geq 1$.

We owe this lemma to Jack [1] (also, by Miller and Mocanu [3]).

LEMMA 2. Let $p(z)$ be analytic in Δ , $p(0) = 1$ and suppose that

$$(1) \quad \operatorname{Re} \left(p(z) + \alpha \frac{zp'(z)}{p(z)} \right) > \frac{1-\alpha}{2} \quad \text{in } \Delta,$$

when α is a positive real number, or

$$(2) \quad \operatorname{Re} \left(p(z) + \alpha \frac{zp'(z)}{p(z)} \right) < \frac{1-\alpha}{2} \quad \text{in } \Delta,$$

when $\alpha < -1$.

Then we have

$$\left| \frac{p(z) - 1}{p(z)} \right| < 1 \quad \text{in } \Delta,$$

or

$$\operatorname{Re} p(z) > \frac{1}{2} \quad \text{in } \Delta.$$

PROOF. From the assumptions (1) and (2), we have $p(z) \neq 0$ in Δ , because if there exists a point $\beta \in \Delta$ such that $p(\beta) = 0$ and $\beta \neq 0$, then we can write

$$p(z) = (z - \beta)^s p_1(z),$$

where s is a positive integer and $p_1(\beta) \neq 0$, then we have

$$(3) \quad \operatorname{Re} \left[p(z) + \alpha \frac{zp'(z)}{p(z)} \right] \\ = \operatorname{Re} \left[(z - \beta)^s p_1(z) + \frac{\alpha sz}{z - \beta} + \frac{\alpha zp_1'(z)}{p_1(z)} \right].$$

Letting $z \rightarrow \beta$ on the straight line which pass through the origin and β , then the right hand side of (3) become positive and negative infinite.

This contradicts (1) and (2).

Therefore we have

$$p(z) \neq 0 \quad \text{in } 0 < |z| < 1.$$

On the other hand, from the assumption $p(0) = 1$, this shows that

$$p(z) \neq 0 \quad \text{in } \Delta.$$

Let us put

$$p(z) = \frac{1}{1 - w(z)}$$

or

$$w(z) = 1 - \frac{1}{p(z)}.$$

Then $w(z)$ is analytic in Δ and $w(0) = 0$, since $p(z) \neq 0$ in Δ . If there exists a point z_0 such that $|w(z)| < 1$ for $|z| < |z_0| < 1$, $|w(z_0)| = 1$, $w(z_0) = e^{i\theta}$, then from LEMMA 1, we have

$$z_0 w'(z_0) = kw(z_0).$$

Then we have

$$\begin{aligned} \operatorname{Re} \left(p(z_0) + \frac{z_0 p'(z_0)}{p(z_0)} \right) &= \operatorname{Re} \left(\frac{1}{1 - w(z_0)} - \frac{\alpha z_0 w'(z_0)}{1 - w(z_0)} \right) \\ &= \operatorname{Re} \left(\frac{1}{1 - e^{i\theta}} + \frac{\alpha k e^{i\theta}}{1 - e^{i\theta}} \right) \\ &= \frac{1}{2} - \frac{\alpha k}{2} \quad \left\{ \begin{array}{l} \leq \frac{1 - \alpha}{2} \quad \text{for the case } \alpha > 0 \\ \geq \frac{1 - \alpha}{2} \quad \text{for the case } \alpha < -1. \end{array} \right. \end{aligned}$$

This contradicts (1) and (2). Therefore we have $|w(z)| < 1$ in Δ .

This shows that

$$\left| \frac{p(z) - 1}{p(z)} \right| < 1 \quad \text{in } \Delta$$

or

$$\operatorname{Re} p(z) > \frac{1}{2} \quad \text{in } \Delta.$$

From LEMMA 2, we easily have the following theorem.

MAIN THEOREM. Let $f(z) \in A$ and suppose that

$$\begin{aligned} \operatorname{Re} \left[(1 - \alpha) \frac{zf'(z)}{f(z)} + \alpha \left(1 + \frac{zf''(z)}{f'(z)} \right) \right] \\ \left\{ \begin{array}{l} \geq \frac{1 - \alpha}{2} \quad \text{for the case } \alpha > 0 \\ \leq \frac{1 - \alpha}{2} \quad \text{for the case } \alpha < -1. \end{array} \right. \end{aligned}$$

Then $f(z)$ is starlike of order at least $1/2$ and

$$\frac{|zf'(z) - f(z)|}{|zf'(z)|} < 1 \quad \text{in } \Delta.$$

REMARK. It is trivial that MAIN THEOREM partly improves THEOREM A.

REFERENCES

1. I. S. Jack, Functions starlike and convex of order α , J. London Math. Soc. 3 (1971), 469-474.
2. A. Marx, Untersuchungen über schlicht Abbildungen, Math. Ann. 107 (1932), 40-67.
3. S. S. Miller and P. T. Mocanu, Second order differential inequalities in the complex plane, J. Math. Anal. Appl. 65 (1978), 289-305.
4. S. S. Miller, P. T. Mocanu and M. O. Reade, All α -convex functions are univalent and starlike, Proc. Amer. Math. Soc. 37 (1973), 553-554.
5. E. Strohäcker, Beiträge zur Theorie der schlichten Funktionen, Math. Z. 37 (1933), 356-380.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GUNMA, ARAMAKI, MAEBASHI 371, JAPAN

1980 Mathematics Subject Classification (1985 Revision). Primary 30C45.