

# Moduli spaces of maps with two critical points

Masayo FUJIMURA

Dept. of Math., College of Sci. and Tech., Nihon Univ.

## Abstract

We give directly a defining equation of the symmetry locus, a singular part of the moduli space of the quadratic rational maps. We show a characterization of this locus. We can expand analogous discussion for the cubic polynomials and give a “chart” making a comparison between properties of these moduli spaces in Appendix A. Moreover, we apply these method to the polynomials of degree  $n$ , and give some conjectures.

## 1 Quadratic rational maps

### 1.1 Moduli space of quadratic rational maps

Let  $\overline{\mathbf{C}}$  be the Riemann sphere and  $\text{Rat}_2(\mathbf{C})$  the space of all quadratic rational maps from  $\overline{\mathbf{C}}$  to itself. The group  $\text{PSL}_2(\mathbf{C})$  of Möbius transformations acts on the space  $\text{Rat}_2(\mathbf{C})$  by conjugation,

$$g \circ f \circ g^{-1} \in \text{Rat}_2(\mathbf{C}) \quad \text{for } g \in \text{PSL}_2(\mathbf{C}), f \in \text{Rat}_2(\mathbf{C}).$$

Two maps  $f_1, f_2 \in \text{Rat}_2(\mathbf{C})$  are **holomorphically conjugate**, denoted by  $f_1 \sim f_2$ , if and only if there exists  $g \in \text{PSL}_2(\mathbf{C})$  with  $g \circ f_1 \circ g^{-1} = f_2$ . The quotient space of  $\text{Rat}_2(\mathbf{C})$  under this action will be denoted by  $\mathcal{M}_2(\mathbf{C})$ , and called the **moduli space** of holomorphic conjugacy classes  $\langle f \rangle$  of quadratic rational maps  $f$ .

Milnor introduced coordinates in  $\mathcal{M}_2(\mathbf{C})$  as follows; for each  $f \in \text{Rat}_2(\mathbf{C})$ , let  $z_1, z_2, z_3$  be the fixed points of  $f$  and  $\mu_i$  the multipliers of  $z_i$ ;  $\mu_i = f'(z_i)$  ( $1 \leq i \leq 3$ ). Consider the elementary symmetric functions of the three multipliers,

$$\sigma_1 = \mu_1 + \mu_2 + \mu_3, \quad \sigma_2 = \mu_1\mu_2 + \mu_2\mu_3 + \mu_3\mu_1, \quad \sigma_3 = \mu_1\mu_2\mu_3.$$

These three multipliers determine  $f$  up to holomorphic conjugacy, and are subject only to the restriction that

$$\sigma_3 = \sigma_1 - 2.$$

Hence the moduli space  $\mathcal{M}_2(\mathbf{C})$  is canonically isomorphic to  $\mathbf{C}^2$  with coordinates  $\sigma_1$  and  $\sigma_2$  (Lemma 3.1 in [Mil93]).

For each  $\mu \in \mathbf{C}$  let  $\text{Per}_n(\mu)$  be the set of all conjugacy classes  $\langle f \rangle$  of maps  $f$  which having a periodic point of period  $n$  and multiplier  $\mu$ .

Each of  $\text{Per}_1(\mu)$  and  $\text{Per}_2(\mu)$  forms a straight lines as follows:

$$\begin{aligned}\text{Per}_1(\mu) &= \{ \langle f \rangle \in \mathcal{M}_2(\mathbf{C}); \sigma_2 = (\mu + \mu^{-1})\sigma_1 - (\mu^2 + 2\mu^{-1}) \} \\ \text{Per}_2(\mu) &= \{ \langle f \rangle \in \mathcal{M}_2(\mathbf{C}); \sigma_2 = -2\sigma_1 + \mu \},\end{aligned}$$

(Lemmas 3.4 and 3.6 in [Mil93]).

**Remark**  $\text{Per}_1(-1) \subseteq \text{Per}_2(1)$  by definition. But, in the case of  $\mathcal{M}_2(\mathbf{C})$ , it is clear that two families coincide.

## 1.2 Symmetry locus

By an automorphism of a quadratic rational map  $f$ , we will mean  $g \in \text{PSL}_2(\mathbf{C})$  which commutes with  $f$ . The collection  $\text{Aut}(f)$  of all automorphisms of  $f$  forms a finite group. It is clear that  $\text{Aut}(\tilde{f})$  is isomorphic to  $\text{Aut}(f)$  for any  $\tilde{f} \in \langle f \rangle$ .

The set

$$\mathcal{S} = \{ \langle f \rangle; \text{Aut}(f) \text{ is non-trivial} \} \subset \mathcal{M}_2(\mathbf{C})$$

is called the **symmetry locus**.

**Corollary 1** *The symmetry locus  $\mathcal{S}$  of quadratic rational maps forms an irreducible algebraic curve as follows;*

$$S(\sigma_1, \sigma_2) = 2\sigma_1^3 + \sigma_1^2\sigma_2 - \sigma_1^2 - 4\sigma_2^2 - 8\sigma_1\sigma_2 + 12\sigma_1 + 12\sigma_2 - 36 = 0. \quad (1)$$



### Proof of Corollary 1.

$\text{Aut}(f)$  coincides with the group consisting of all permutations of the fixed points which preserve the multipliers. In the case of  $f$  has the three distinct fixed points,  $\text{Aut}(f)$  has order 1, 2, or 6 according as three multipliers are distinct, two are equal, or all the three are equal, respectively, while, if  $f$  has multiple fixed points then  $\text{Aut}(f)$  is non-trivial if and only if  $f$  has a triple fixed point. The multipliers  $\mu_i$  are the roots of the equation:

$$\mu^3 - \sigma_1\mu^2 + \sigma_2\mu - \sigma_1 + 2 = 0. \quad (2)$$

The equation (2) has multiple roots if and only if its discriminant is equal to zero. Hence we have

$$(\sigma_2 - 2\sigma_1 + 3)(2\sigma_1^3 + \sigma_1^2\sigma_2 - \sigma_1^2 - 4\sigma_2^2 - 8\sigma_1\sigma_2 + 12\sigma_1 + 12\sigma_2 - 36) = 0.$$

Figure 1:  $\mathcal{M}_2(\mathbf{R})$  with the real cut of  $\mathcal{S}$ .Figure 2: Lines  $\{\text{Per}_1(\mu)\}$  in the real cut of the moduli space  $\mathcal{M}_2(\mathbf{C})$ 

The first factor corresponds with  $\text{Per}_1(1)$ . Considering the line of the first factor ( $\text{Per}_1(1)$ ) tangent to the curve of the second factor ( $S$ ) with tangency of degree three, the second factor is the required equation. ■

The following result is obtained immediately by the definition of the envelope of the family of curves.

**Corollary 2** *The envelope of  $\{\text{Per}_1(\mu)\}_\mu$  coincides with the symmetry locus.*

**Remark** (Theorem 5.1. of [Mil93]) A quadratic rational map has a non-trivial automorphism if and only if it is conjugate to a map in the unique normal form  $f(z) = k(z + \frac{1}{z})$  with  $k \in \mathbf{C} \setminus \{0\}$ .

### 1.3 Real moduli space

Let  $\text{Rat}_2(\mathbf{R})$  be the set of real quadratic rational maps. Then the parameters  $\sigma_i$  ( $1 \leq i \leq 3$ ) are all real, because the three fixed points and the corresponding multipliers are either all real or one real and a pair of complex conjugate numbers. According to J. Milnor, we define the real moduli space  $\mathcal{M}_2(\mathbf{R})$  for  $\text{Rat}_2(\mathbf{R})$  to be simply the real  $(\sigma_1, \sigma_2)$ -plane. This notation needs some care when used: if we put  $\mathcal{S}_{\mathbf{R}} = \mathcal{S} \cap \mathcal{M}_2(\mathbf{R})$ , and denote by  $\langle \rangle_{\mathbf{R}}$  the real conjugacy class, then  $(\text{Rat}_2(\mathbf{R})/\text{PGL}_2(\mathbf{R})) \setminus \{\langle a(x + \frac{1}{x}) \rangle_{\mathbf{R}}, \langle a(x - \frac{1}{x}) \rangle_{\mathbf{R}}\}_{a \in \mathbf{R}^\times}$  is canonically isomorphic to  $\mathbf{R}^2 \setminus \mathcal{S}_{\mathbf{R}}$ , whereas there is a canonical two-to-one correspondence between  $\{\langle a(x \pm \frac{1}{x}) \rangle_{a \in \mathbf{R}^\times}\}$  and  $\mathcal{S}_{\mathbf{R}}$ .

## 2 Cubic polynomials

### 2.1 Moduli space of cubic polynomials

Let be  $\text{Poly}_3(\mathbf{C})$  the space of all cubic polynomials from  $\mathbf{C}$  to itself. The group  $\text{Poly}_3(\mathbf{C})$  of affine transformations acts on the space  $\text{Poly}_3(\mathbf{C})$  by conjugation,

$$g \circ p \circ g^{-1} \in \text{Poly}_3(\mathbf{C}) \quad \text{for} \quad g \in \text{Poly}_1(\mathbf{C}), \quad p \in \text{Poly}_3(\mathbf{C}).$$

Two maps  $p_1, p_2 \in \text{Poly}_3(\mathbf{C})$  are **holomorphically conjugate**, denoted by  $p_1 \sim p_2$ , if and only if there exists  $g \in \text{Poly}_1(\mathbf{C})$  with  $g \circ p_1 \circ g^{-1} = p_2$ . The quotient space of  $\text{Poly}_3(\mathbf{C})$  under this action will be denoted by  $M_3(\mathbf{C})$ , and called the **moduli space** of holomorphic conjugacy classes  $\langle p \rangle$  of cubic polynomials  $p$ .

Doing the same as the case of quadratic rational maps, we introduce coordinates in  $M_3(\mathbf{C})$  as follows; for each  $p \in \text{Poly}_3(\mathbf{C})$ , let  $z_1, z_2, z_3, z_4 (= \infty)$  be the fixed points of  $p$  and  $\mu_i$  the multipliers of  $z_i$ ;  $\mu_i = p'(z_i)$  ( $1 \leq i \leq 3$ ), and  $\mu_4 = 0$ . Consider the elementary symmetric functions of the four multipliers,

$$\begin{aligned} \sigma_1 &= \mu_1 + \mu_2 + \mu_3 + \mu_4 = \mu_1 + \mu_2 + \mu_3 \\ \sigma_2 &= \mu_1\mu_2 + \mu_1\mu_3 + \mu_1\mu_4 + \mu_2\mu_3 + \mu_2\mu_4 + \mu_3\mu_4 = \mu_1\mu_2 + \mu_1\mu_3 + \mu_2\mu_3 \\ \sigma_3 &= \mu_1\mu_2\mu_3 + \mu_1\mu_2\mu_4 + \mu_1\mu_3\mu_4 + \mu_2\mu_3\mu_4 = \mu_1\mu_2\mu_3 \\ \sigma_4 &= \mu_1\mu_2\mu_3\mu_4 = 0. \end{aligned}$$

These multipliers determine uniquely  $p$  up to holomorphic conjugacy, and are subject only to the restriction that

$$3 - 2\sigma_1 + \sigma_2 = 0.$$

Hence the moduli space  $M_3(\mathbf{C})$  is canonically isomorphic to  $\mathbf{C}^2$  with coordinates  $\sigma_1$  and  $\sigma_3$ .

**Proposition 1**    *The locus  $\text{Per}_1(\mu)$  forms a straight lines as follows:*

$$\text{Per}_1(\mu) = \{ \langle f \rangle \in M_3(\mathbf{C}); \sigma_3 = (-\mu^2 + 2\mu)\sigma_1 + \mu^3 - 3\mu \}.$$

*The locus  $\text{Per}_2(\mu)$  forms an algebraic curve of degree three as follows:*

$$\begin{aligned} \text{Per}_2(\mu) &= \{ \langle f \rangle \in \mathcal{M}_2(\mathbf{C}); \sigma_3^2 + (4\sigma_1^2 - (\mu + 57)\sigma_1 + 252)\sigma_3 - (4\mu - 16)\sigma_1^3 \\ &\quad + (61\mu - 252)\sigma_1^2 - (4\mu^2 + 246\mu - 1134)\sigma_1 - \mu^3 + 51\mu^2 \\ &\quad - 99\mu - 459 = 0 \}. \end{aligned}$$

*Note that this curve is irreducible if and only if  $\mu \neq 1$ . In the case of  $\mu = 1$ ,*

$$\text{Per}_2(1) = \{ \text{Per}_1(-1) \} \cup \{ \langle f \rangle \in \mathcal{M}_2(\mathbf{C}); \sigma_3 + 4\sigma_1^2 - 61\sigma_1 + 254 = 0 \}.$$

## 2.2 Symmetry locus

Using conjugation described in above, we can define symmetry locus of this moduli space as one in  $\mathcal{M}_2(\mathbf{C})$ , and we obtain next results.

**Theorem 1** *The symmetry locus  $\mathcal{S}$  of cubic polynomials forms an irreducible algebraic curve:*

$$S(\sigma_1, \sigma_3) = 27\sigma_3 + (\sigma_1 - 6)(2\sigma_1 - 3)^2 = 0. \quad (3)$$

The following result is obtained immediately by the definition of the envelope of the family of curves.

**Corollary 3** *The envelope of  $\{\text{Per}_1(\mu)\}_\mu$  coincides with the symmetry locus.*




Figure 3:  $M_3(\mathbf{R})$  with the real cut of  $\mathcal{S}$ .




Figure 4: Lines  $\{\text{Per}_1(\mu)\}$  in the real cut of the moduli space  $M_3(\mathbf{C})$ .

**Remark** A cubic polynomial has non-trivial automorphism if and only if it is conjugate to a map in the unique normal form  $p(z) = z^3 + az$ .

## 2.3 Real moduli space

Let  $\text{Poly}_3(\mathbf{R})$  be the set of real cubic polynomials. By the same reason for the case of  $\mathcal{M}_2$ , we define the real moduli space  $M_3(\mathbf{R})$  for  $\text{Poly}_3(\mathbf{R})$  to be simply the real  $(\sigma_1, \sigma_3)$ -plane. This notation needs some care when used: if we put  $\mathcal{S}_{\mathbf{R}} = \mathcal{S} \cap M_3(\mathbf{R})$ , and denote by  $\langle \rangle_{\mathbf{R}}$  the real conjugacy class, then  $(\text{Poly}_3(\mathbf{R})/\text{Poly}_1(\mathbf{R})) \setminus \{\langle x^3 + ax \rangle_{\mathbf{R}}, \langle -x^3 + ax \rangle_{\mathbf{R}}\}_{a \in \mathbf{R}^\times}$  is canonically isomorphic to  $\mathbf{R}^2 \setminus \mathcal{S}_{\mathbf{R}}$ , whereas there is a canonical two-to-one correspondence between  $\{\langle \pm x^3 + ax \rangle\}_{a \in \mathbf{R}^\times}$  and  $\mathcal{S}_{\mathbf{R}}$ .

### 3 Polynomials of degree $n$

#### 3.1 Moduli space of polynomials of degree $n$

Now we discuss about the moduli space  $M_n(\mathbf{C})$  for the space,  $\text{Poly}_n(\mathbf{C})$ , of polynomials of degree  $n$ .

Doing the same as the case of cubic polynomials, we try introducing coordinates in  $M_n(\mathbf{C})$  as follows; for each  $p(z) \in \text{Poly}_n(\mathbf{C})$ , let  $z_1, \dots, z_n, z_{n+1}(= \infty)$  be the fixed points of  $p$  and  $\mu_i$  the multipliers of  $z_i$ ;  $\mu_i = p'(z_i)$  ( $1 \leq i \leq n$ ), and  $\mu_{n+1} = 0$ . Consider the elementary symmetric functions of the  $n$  multipliers,

$$\begin{aligned}\sigma_{n,1} &= \mu_1 + \dots + \mu_n, \\ \sigma_{n,2} &= \mu_1\mu_2 + \dots + \mu_{n-1}\mu_n = \sum_{i=1}^{n-1} \mu_i \sum_{j>i}^n \mu_j, \\ &\dots \\ \sigma_{n,n} &= \mu_1\mu_2 \dots \mu_n, \\ \sigma_{n,n+1} &= 0.\end{aligned}$$

**Example 1** For example, we assume  $p(z) \in \text{Poly}_4(\mathbf{C})$ ;

- fixed points:  $z_1, z_2, z_3, z_4, \infty$
- multiplier:  $\mu_1, \mu_2, \mu_3, \mu_4, 0$
- elementary symmetric functions:

$$\begin{cases} \sigma_{4,1} = \mu_1 + \mu_2 + \mu_3 + \mu_4 \\ \sigma_{4,2} = \mu_1\mu_2 + \mu_1\mu_3 + \mu_1\mu_4 + \mu_2\mu_3 + \mu_2\mu_4 + \mu_3\mu_4 \\ \sigma_{4,3} = \mu_1\mu_2\mu_3 + \mu_1\mu_2\mu_4 + \mu_1\mu_3\mu_4 + \mu_2\mu_3\mu_4 \\ \sigma_{4,4} = \mu_1\mu_2\mu_3\mu_4 \\ \sigma_{4,5} = 0 \end{cases}$$

Applying Fatou-index theorem to these fixed points;

$$\frac{1}{1-\mu_1} + \frac{1}{1-\mu_2} + \frac{1}{1-\mu_3} + \frac{1}{1-\mu_4} + \frac{1}{1-0} = 1, \quad (4)$$

where  $\mu_i \neq 1$  ( $1 < i < n$ ). Arranging this equation for the form of elementary symmetric functions;

$$4 - 3(\mu_1 + \mu_2 + \mu_3 + \mu_4) + 2(\mu_1\mu_2 + \mu_1\mu_3 + \mu_1\mu_4 + \mu_2\mu_3 + \mu_2\mu_4 + \mu_3\mu_4) - (\mu_1\mu_2\mu_3 + \mu_1\mu_2\mu_4 + \mu_1\mu_3\mu_4 + \mu_2\mu_3\mu_4) = 0.$$

Hence we have

$$4 - 3\sigma_{4,1} + 2\sigma_{4,2} - \sigma_{4,3} = 0. \quad (5)$$

For the equation (5), the cases  $\mu_i = 1$  are also allowable.

Now we consider a polynomial  $p(z) = a_4z^4 + a_3z^3 + a_2z^2 + a_1z + a_0 \in \text{Poly}_4(\mathbf{C})$  that has at least two fixed points. After affine conjugation, we can assume they are 0 and 1. Then, we will solve the following question: “Do the four multipliers

$$\mu_0 = p'(0), \mu_1 = p'(1), \mu_2 = p'(z_2), \mu_3 = p'(z_3),$$

where  $z_1, z_2$  are fixed points of  $p(z)$ , determine the five coefficients  $a_4, a_3, a_2, a_1, a_0$  of  $p(z)$ ?”

In fact, the following equations hold;

$$\begin{aligned} a_0 &= 0 && \text{because of } f(0) = 0, \\ a_1 &= \mu_0 && \text{because of } f'(0) = \mu_0, \\ a_2 &= a_4 + 3 - 2\mu_0 - \mu_1 && \text{because of } f'(1) = \mu_1, \\ a_3 &= 1 - a_4 - a_2 - \mu_0 && \text{because of } f(1) = 1, \end{aligned}$$

and  $a_4$  is a common root of the following two equations;

$$\begin{aligned} A_1 &= (\mu_2^2 - 2\mu_3\mu_2 + \mu_3^2 - \mu_0^2 + 2\mu_1\mu_0 - \mu_1^2)a_4^4 + (-4\mu_0^3 + (4\mu_1 + 8)\mu_0^2 + (-4\mu_1^2 - 8)\mu_0 + 4\mu_1^3 - 8\mu_1^2 + 8\mu_1)a_4^3 + (-6\mu_0^4 + (-4\mu_1 + 28)\mu_0^3 + (4\mu_1^2 + 4\mu_1 - 44)\mu_0^2 + (-4\mu_1^3 + 4\mu_1^2 - 8\mu_1 + 32)\mu_0 - 6\mu_1^4 + 28\mu_1^3 - 44\mu_1^2 + 32\mu_1 - 16)a_4^2 + (-4\mu_0^5 + (-12\mu_1 + 32)\mu_0^4 + (-8\mu_1^2 + 64\mu_1 - 96)\mu_0^3 + (8\mu_1^3 - 96\mu_1 + 128)\mu_0^2 + (12\mu_1^4 - 64\mu_1^3 + 96\mu_1^2 - 64)\mu_0 + 4\mu_1^5 - 32\mu_1^4 + 96\mu_1^3 - 128\mu_1^2 + 64\mu_1)a_4 - \mu_0^6 + (-6\mu_1 + 12)\mu_0^5 + (-15\mu_1^2 + 60\mu_1 - 60)\mu_0^4 + (-20\mu_1^3 + 120\mu_1^2 - 240\mu_1 + 160)\mu_0^3 + (-15\mu_1^4 + 120\mu_1^3 - 360\mu_1^2 + 480\mu_1 - 240)\mu_0^2 + (-6\mu_1^5 + 60\mu_1^4 - 240\mu_1^3 + 480\mu_1^2 - 480\mu_1 + 192)\mu_0 - \mu_1^6 + 12\mu_1^5 - 60\mu_1^4 + 160\mu_1^3 - 240\mu_1^2 + 192\mu_1 - 64 = 0, \\ A_2 &= (\mu_2 + \mu_3 + \mu_0 + \mu_1 - 4)a_4^2 + (2\mu_0^2 - 4\mu_0 - 2\mu_1^2 + 4\mu_1)a_4 + \mu_0^3 + (3\mu_1 - 6)\mu_0^2 + (3\mu_1^2 - 12\mu_1 + 12)\mu_0 + \mu_1^3 - 6\mu_1^2 + 12\mu_1 - 8 = 0. \end{aligned}$$

Above two equations have common roots if and only if  $\mu_0, \mu_1, \mu_2, \mu_3$  satisfy the equation (5). Since  $\mu_0, \mu_1, \mu_2, \mu_3$  are the four multipliers of  $p(z)$  and they should satisfy the equation (5), the two equations always have common roots. Hence five coefficients of  $p(z)$  are calculated by its four multipliers, however, this calculation is not decisive when they have distinct two common roots.

For the case of  $\text{Poly}_n(\mathbf{C})$ , it is clear from (4) that the equation corresponds to (5) cannot have the term of  $\sigma_{n,n}$ . Hence we can put

$$c_0 + c_1\sigma_{n,1} + c_2\sigma_{n,2} + \cdots + c_{n-1}\sigma_{n,n-1} = 0$$

where  $c_k$  ( $0 \leq k \leq n-1$ ) are functions of  $n$  variable.

Paying attention to the form of elementary symmetric functions, we obtain the following equation;

$$c_k = (-1)^k \frac{\binom{n-1}{k} n}{\binom{n}{k}} = n - k.$$

where  $\binom{n}{k}$  means binomial coefficient. For convenience, put  $\sigma_{n,0} = 1$ . we have

$$\sum_{k=0}^{n-1} (-1)^k (n-k) \sigma_{n,k} = 0. \quad (6)$$

**Question** Is the moduli space  $M_n(\mathbf{C})$  for polynomials of degree  $n$  canonically isomorphic to  $\mathbf{C}^{n-1}$  with coordinates  $\sigma_1, \sigma_2, \dots, \sigma_{n-2}$ , and  $\sigma_n$ ?

### 3.2 Symmetry locus

**Proposition 2** *A polynomial of degree four has a non-trivial automorphism if and only if it is conjugate to a map in the unique normal form*

$$\{z^4 + az\}, \quad a \in \mathbf{C}.$$

*For a map  $p(z)$  in this normal form,  $\text{Aut}(p)$  is a cyclic group of order three.*

**Outline of proof.** Let  $p(z) \in \text{Poly}_4(\mathbf{C})$ .

1. In the case of a map  $p(z)$  with multiple fixed points.
  - (a) The case of  $p(z)$  with a fixed point of order four:  $\text{Aut}(p)$  is non-trivial.
  - (b) The case of  $p(z)$  with a fixed point of order three:  $\text{Aut}(p)$  is trivial.
  - (c) The case of  $p(z)$  with two fixed points of order two: there is not such  $p(z)$ .
  - (d) The case of  $p(z)$  with a fixed point of order two:  $\text{Aut}(p)$  is trivial.
2. In the case of a map  $p(z)$  with four distinct fixed points.
  - (a) The case of four distinct multipliers:  $\text{Aut}(p)$  is trivial.
  - (b) The case that only two of multipliers are coincide:  $\text{Aut}(p)$  is trivial.
  - (c) The case of two pair of same multipliers: there is not such  $p(z)$ .
  - (d) The case of three same multipliers: By an affine conjugation, if three fixed points (whose multipliers are same) are mapped on the vertices of a regular triangle whose barycenter is the origin and the other fixed point on the origin, then  $\text{Aut}(p)$  is non-trivial. Otherwise  $\text{Aut}(p)$  is trivial.
  - (e) The case of four same multipliers: there is not such  $p(z)$ .



Therefore a map  $p(z)$  has non-trivial automorphisms if and only if  $p(z)$  is in the case 1-(a) and the first part of 2-(d). We can check easily that these maps coincide with the normal form  $\{z^4 + az\}$ . ■

**Conjecture** A polynomial of degree  $n$  has a non-trivial automorphism if and only if it is conjugate to a map in the unique normal form

$$\left\{ z^n + \sum_{k|(n-1), k \neq n-1} A(k)z^k \right\}$$

where  $A(k)$  are parameters in  $\mathbf{C}$ .

## References

- [FN] M. Fujimura and K. Nishizawa. Bifurcations and Algebraic Curves for Quadratic Rational Functions  $\{mf(x)\}_m$ . submitted.
- [FN96] M. Fujimura and K. Nishizawa. Bifurcations of Nusse-Yorke's family in the quadratic rational functions. In T. Sugawa, editor, *RIMS Kokyuroku 959: Complex Dynamics and Related Problems*, pages 90–102. Kyoto Univ., 1996.
- [Mil92] J. Milnor. Remarks on iterated cubic maps. *Experimental Mathematics*, 1:5–24, 1992.
- [Mil93] J. Milnor. Geometry and Dynamics of Quadratic Rational Maps. *Experimental Mathematics*, 2(1):37–83, 1993.

## A Comparison between the quadratic rational maps and cubic polynomials

	Quadratic rational maps	Cubic polynomials
<i>Moduli Space</i>	$\mathcal{M}_2(\mathbf{C}) \simeq \mathbf{C}^2$	$\mathcal{M}_3(\mathbf{C}) \simeq \mathbf{C}^2$
<i>Real Moduli Space</i>	$\mathcal{M}_2(\mathbf{R}) \simeq \mathbf{R}^2$ <i>excepts on the symm. locus</i>	$\mathcal{M}_3(\mathbf{R}) \simeq \mathbf{R}^2$ <i>excepts on the symm. locus</i>
<i>Coordinates</i>	$(\sigma_1, \sigma_2), \quad \sigma_3 = \sigma_1 - 2$	$(\sigma_1, \sigma_3), \quad 3 - 2\sigma_1 + \sigma_2 = 0$
<i>Normal Forms</i>	<i>Fixed Point Normal Form, etc.</i>	$\{f(z) = z^3 + az + b\}_{(a,b)}$
<i>Periodic Orbits</i>	$\text{Per}_1(\mu) :$ $\sigma_2 = (\mu + \frac{1}{\mu})\sigma_1 - (\mu^2 + \frac{2}{\mu})$ $\text{Per}_2(\mu) :$ $2\sigma_1 + \sigma_2 = \mu$ $\text{Per}_1(-1) = \text{Per}_2(1)$	$\text{Per}_1(\mu) :$ $\sigma_3 = (-\mu^2 + 2\mu)\sigma_1 + \mu^3 - 3\mu$ $\text{Per}_2(\mu) :$ <i>cubic algebraic curve</i> $\text{Per}_1(-1) \subset \text{Per}_2(1)$
<i>Symmetry Locus</i>	<i>the envelope of</i> $\{\text{Per}_1(\mu)\}$ <i>normal form</i> : $\{k(z + \frac{1}{z})\}$	<i>the envelope of</i> $\{\text{Per}_1(\mu)\}$ <i>normal form</i> : $\{z^3 + az\}$
<i>Topological Partition</i>	<i>degree</i> $\pm 2$ , <i>monotone</i> , <i>unimodal, bimodal</i>	$\mathcal{R}_0, \mathcal{R}_1, \mathcal{R}_2, \mathcal{R}_3$
<i>Hyp. Components</i>	B, C, D, E	A, B, C, D

Masayo FUJIMURA

Dept. of Math., College of Sci. and Tech., Nihon Univ.  
 1-8, Kanda-Surugadai, Chiyoda-ku, Tokyo, 101 JAPAN  
 e-mail: masayo@math.cst.nihon-u.ac.jp