

# A Real Inversion Formula for the Laplace Transform in a Sobolev Space (preliminary report)

群馬大工 天野 一男 (Kazuo Amano)  
群馬大工 斎藤 三郎 (Saburou Saitoh)  
群馬大工 アドミ シャリフ (Admi Syarif)

For the real-valued Sobolev Hilbert space on  $[0, \infty)$  comprising absolutely continuous functions  $F(t)$  normalized by  $F(0) = 0$  and equipped with the inner product

$$(F_1, F_2) = \int_0^\infty (F_1(t)F_2(t) + F_1'(t)F_2'(t)) dt,$$

we shall establish the real inversion formula and its error estimate for the Laplace transform of the Sobolev Hilbert space.

**Key words:** Laplace transform, real inversion formula, Sobolev space, reproducing kernel, Mellin transform, Szegő space.

AMS subject classification: 44A10, 30C40

## 1 Introduction and results

The real inversion formulas for the Laplace transform are important in mathematical sciences, but the formulas are, in general, very involved. See, for example [7, 11]. In [3, 10], new real inversion formulas for some general situations were given by a new method for integral transforms in the framework of Hilbert spaces. In some special cases, their error estimates were given in [2]. In the new method, inversion formulas for integral transforms will be, in general, given in terms of strong convergence. For some practical purposes, we wish to obtain inversion formulas in terms of pointwise convergence. For this purpose, we shall establish a real inversion formula for the Laplace transform of a simple Sobolev space, which will be given in terms of pointwise convergence.

Let  $S$  be the Sobolev Hilbert space on  $t \geq 0$  comprising absolutely continuous real-valued functions  $F(t)$  normalized by  $F(0) = 0$  and equipped with the

inner product

$$(F_1, F_2)_S = \int_0^{\infty} (F_1(t)F_2(t) + F_1'(t)F_2'(t)) dt.$$

We consider the Laplace transform of  $F \in S$

$$f(x) = [LF](x) = \int_0^{\infty} F(t)e^{-xt} dt, \quad x > 0. \quad (1)$$

In connection with some general real inversion formulas [3, 10], we would like to consider a more general Sobolev space such that for any positive  $q$  the following inner product is given by

$$(F_1, F_2)_{S,q} = \int_0^{\infty} (F_1(t)F_2(t) + F_1'(t)F_2'(t)) t^{1-2q} dt.$$

However in this general case, its reproducing kernel will be very involved. So, we shall consider the simple Sobolev space  $S$ . For more general order Sobolev spaces, the circumstances are similar. That is, the Sobolev space  $S$  will be a reasonable space for the Laplace transform for our purposes. See Lemmas 1 and 3 for this comment.

Then, we obtain

**Theorem.** *For the Laplace transform (1) of the Sobolev Hilbert space  $S$ , we have the real inversion formula*

$$F(t) = \lim_{N \rightarrow \infty} \int_0^{\infty} f(x) \int_0^{\infty} e^{-x\tau} K(\tau, t) (P_N(x, \tau) + Q_N(x, \tau)) d\tau dx \quad (2)$$

where

$$\begin{aligned} K(\tau, t) &= \frac{1}{2} (e^{-|\tau-t|} - e^{-\tau} e^{-t}), \\ P_N(x, \tau) &= \sum_{n=0}^N \sum_{\nu=n}^{2n} (-1)^{\nu-n+1} \binom{2n}{\nu} \binom{\nu}{n} \frac{1}{(n+1)(\nu+1)!} (\tau x)^\nu \\ &\quad \times ((2n+1)(\tau x)^2 - (2+5n+\nu+3n\nu)\tau x + n(\nu+1)^2), \\ Q_N(x, \tau) &= \frac{1}{\tau^2} \sum_{n=0}^N \sum_{\nu=n}^{2n} (-1)^{\nu-n+1} \binom{2n}{\nu} \binom{\nu}{n} \frac{1}{(n+1)(\nu+2)!} (\tau x)^{\nu+1} \end{aligned} \quad (3)$$

$$\times \left( (4n^2 + 6n + 2)(\tau x)^3 - (8 + 3\nu + 26n + 10n\nu + 20n^2 + 8n^2\nu)(\tau x)^2 + (\nu + 2)(3 + \nu + 10n + 4n\nu + 9n^2 + 5n^2\nu)(\tau x) - n^2(\nu + 1)^2(\nu + 2) \right). \quad (4)$$

In the real inversion formula (2), for any  $t \geq 0$  the right hand side converges and its convergence is uniform on  $[0, \infty)$ .

We introduce the differential operator

$$D_n = x^n \partial_x^n x \partial_x$$

for any nonnegative integer  $n$ .

## 2 Preliminaries for Theorem

At first we note

**Lemma 1.** *The reproducing kernel  $K(t, \hat{t})$  for the Sobolev Hilbert space  $S$  is given by*

$$\begin{aligned} K(t, \hat{t}) &= \frac{2}{\pi} \int_0^\infty \frac{\sin(t\xi) \sin(\hat{t}\xi)}{\xi^2 + 1} d\xi \\ &= \frac{1}{2} \left( e^{-|t-\hat{t}|} - e^{-t} e^{-\hat{t}} \right). \end{aligned} \quad (5)$$

**Proof.** For the positive matrix  $K(t, \hat{t})$  defined by (5) we shall show that the reproducing kernel Hilbert space  $H_K$  admitting the reproducing kernel  $K(t, \hat{t})$  coincides with  $S$ .

From (5), we see that any member  $F$  of  $H_K$  is expressible in the form

$$F(t) = \frac{2}{\pi} \int_0^\infty \frac{H(\xi) \sin(t\xi)}{\xi^2 + 1} d\xi \quad (6)$$

for a (of course, uniquely determined) function  $H$  satisfying

$$\frac{2}{\pi} \int_0^\infty \frac{H(\xi)^2}{\xi^2 + 1} d\xi < \infty \quad (7)$$

and we have the isometrical identity

$$\| F \|_{H_K}^2 = \frac{2}{\pi} \int_0^\infty \frac{H(\xi)^2}{\xi^2 + 1} d\xi. \quad (8)$$

For this argument see [8, 9, or 10]. From (6)

$$H(\xi) = (\xi^2 + 1) \int_0^\infty F(t) \sin(t\xi) dt, \quad (9)$$

in the  $L_2$  space and so, from (9) we obtain

$$\|F\|_{H_K}^2 = \int_0^\infty (F(t)^2 + F'(t)^2) dt. \quad (10)$$

From the uniqueness of reproducing kernels, we have the desired result.

**Lemma 2.** *In the Laplace transform (1) of  $S$ , we have the isometrical identity*

$$\|F\|_S^2 = \sum_{n=0}^{\infty} \frac{1}{n!(n+1)!} \int_0^\infty \{(D_n f(x))^2 + (D_n(xf(x)))^2\} dx. \quad (11)$$

**Proof.** In general, for  $F \in L_2(0, \infty)$  we have the isometrical identity

$$\int_0^\infty F(t)^2 dt = \sum_{n=0}^{\infty} \frac{1}{n!(n+1)!} \int_0^\infty (D_n f(x))^2 dx \quad (12)$$

([10, Chapter 5]). Since  $F(0) = 0$  and by integration by parts we have

$$\int_0^\infty F'(t) e^{-xt} dt = xf(x). \quad (13)$$

Hence, from (13) we have the desired isometrical identity (11).

**Lemma 3.** *In the Laplace transform (1) of  $S$ , we have the real inversion formula*

$$\begin{aligned} F(t) &= \sum_{n=0}^{\infty} \frac{1}{n!(n+1)!} \int_0^\infty \left[ D_n f(x) \cdot D_n \int_0^\infty e^{-\tau x} K(\tau, t) d\tau \right. \\ &\quad \left. + D_n(xf(x)) \cdot D_n \left( x \int_0^\infty e^{-\tau x} K(\tau, t) d\tau \right) \right] dx \\ &= \sum_{n=0}^{\infty} \frac{1}{n!(n+1)!} \int_0^\infty \left[ D_n f(x) \cdot D_n \left( \frac{e^{-t} - e^{-xt}}{x^2 - 1} \right) \right. \\ &\quad \left. + D_n(xf(x)) \cdot D_n \left( x \left( \frac{e^{-t} - e^{-xt}}{x^2 - 1} \right) \right) \right] dx. \end{aligned} \quad (14)$$

*The convergence of this series is uniform on  $[0, \infty)$ .*

**Proof.** First we have

$$\begin{aligned}
 (LK(\cdot, t))(x) &= \int_0^\infty e^{-x\tau} K(\tau, t) d\tau \\
 &= \int_0^\infty e^{-x\tau} \left( \frac{2}{\pi} \int_0^\infty \frac{\sin(\tau\xi) \sin(t'\xi)}{\xi^2 + 1} d\xi \right) d\tau \\
 &= \frac{2}{\pi} \int_0^\infty \frac{\xi \sin(t\xi)}{(\xi^2 + 1)(\xi^2 + x^2)} d\xi \\
 &= \frac{e^{-t} - e^{-tx}}{x^2 - 1}
 \end{aligned}$$

(see [1], page 410). Hence, by using the reproducing property of  $K(\cdot, t)$  in  $S$

$$F(t) = (F(\cdot), K(\cdot, t))_S \quad (15)$$

and from the isometrical identity (11) we have the desired result (14). The uniform convergence of (14) on  $[0, \infty)$  follows from the general property of reproducing kernel Hilbert spaces (see, [10], page 35, Theorem 1) and the boundedness of the reproducing kernel (5) for  $S$  on  $[0, \infty)$ .

For the property of  $f(x)$  satisfying (12) we note

**Proposition 1** ([10, Chapter 5]). *For a function  $f$  satisfying (12), we have the isometrical identity*

$$\begin{aligned}
 &\sum_{n=0}^{\infty} \frac{1}{n!(n+1)!} \int_0^\infty (D_n f(x))^2 dx \\
 &= \lim_{x \rightarrow 0^+} \frac{1}{2\pi} \int_{-\infty}^{\infty} |f(x+iy)|^2 dy,
 \end{aligned} \quad (16)$$

where  $f(z)$  is analytic on the right half complex plane  $R^+ = \{Re z > 0\}$  and belongs to the Szegő space on  $R^+$  with a finite norm (16). Furthermore, then we have, for  $n \geq 1$ ,  $0 \leq m \leq n-1$ ,

$$\partial_x^m [x f'(x)] x^{n+m+1} = o(1), \text{ as } x \rightarrow 0^+,$$

$$f(x) x^{\frac{1}{2}} = O(1), \text{ as } x \rightarrow 0^+,$$

and for  $n \geq 0$ ,

$$\partial_x^n f(x) \rightarrow 0 \text{ as } x \rightarrow \infty.$$

### 3 Proof of Theorem

For  $n \geq 1$ , by integration by parts and by using Proposition 1, we have

$$\begin{aligned}
& \int_0^\infty D_n f(x) \cdot D_n e^{-x\tau} dx \\
&= \tau^n \int_0^\infty (x f'(x)) \partial_x^n ((nx^{2n} - \tau x^{2n+1}) e^{-x\tau}) dx \\
&= -\tau^n \int_0^\infty f(x) \partial_x x \partial_x^n ((nx^{2n} - \tau x^{2n+1}) e^{-x\tau}) dx \\
&= -\tau^n \int_0^\infty f(x) \left( \partial_x^n ((nx^{2n} - \tau x^{2n+1}) e^{-x\tau}) \right. \\
&\quad \left. + x \partial_x^{n+1} ((nx^{2n} - \tau x^{2n+1}) e^{-x\tau}) \right) dx \\
&= -\tau^n \int_0^\infty f(x) \left( \partial_x^n ((nx^{2n} - \tau x^{2n+1}) e^{-x\tau}) \right. \\
&\quad \left. - x\tau \partial_x^n ((nx^{2n} - \tau x^{2n+1}) e^{-x\tau}) + x \partial_x^n ((2n^2)x^{2n-1} - (2n)\tau x^{2n}) e^{-x\tau} \right) dx \\
&= -(\tau)^n \int_0^\infty f(x) e^{-\tau x} \left( \sum_{\nu=0}^n \binom{n}{\nu} (-\tau)^\nu (n \partial_x^{n-\nu} x^{2n} - \tau \partial_x^{n-\nu} x^{2n+1}) \right. \\
&\quad \left. - \tau x \sum_{\nu=0}^n \binom{n}{\nu} (-\tau)^\nu (n \partial_x^{n-\nu} x^{2n} - \tau \partial_x^{n-\nu} x^{2n+1}) \right. \\
&\quad \left. + x \sum_{\nu=0}^n \binom{n}{\nu} (-\tau)^\nu ((2n^2) \partial_x^{n-\nu} x^{2n-1} - \tau(2n+1) \partial_x^{n-\nu} x^{2n}) \right) dx \\
&= \int_0^\infty f(x) \sum_{\nu=0}^n (-1)^{\nu+1} \binom{n}{\nu} e^{-\tau x} (\tau x)^{(n+\nu)} \frac{\Gamma(2n+1)}{\Gamma(n+\nu+1)} \\
&\quad \times \left( \frac{(2n+1)}{(n+\nu+1)} (\tau x)^2 - \left( \frac{(2n+1)}{(n+\nu+1)} + 3n+1 \right) \tau x + n(n+\nu+1) \right) dx \\
&= \int_0^\infty f(x) e^{-\tau x} P_N(x, \tau) dx.
\end{aligned} \tag{17}$$

Similarly, we have

$$\begin{aligned}
& \int_0^\infty D_n(xf(x)) \cdot D_n(xe^{-x\tau}) dx \\
&= (-\tau)^{(n-1)} \int_0^\infty \partial_x^n (x(xf(x)))' (x^2\tau^2 - (2n+1)x\tau + n^2) e^{-x\tau} x^{2n} dx
\end{aligned}$$

$$\begin{aligned}
&= -\tau^{(n-1)} \int_0^\infty (xf(x) + x^2 f'(x)) \\
&\quad \partial_x^n ((x^{2n+2}\tau^2 - (2n+1)x^{2n+1}\tau + n^2x^{2n}) e^{-x\tau}) dx \\
&= -\tau^{(n-1)} \int_0^\infty f(x) \left( x \partial_x^n (x^{2n+2}\tau^2 - (2n+1)x^{2n+1}\tau + n^2x^{2n}) e^{-x\tau} - \right. \\
&\quad \left. \partial_x (x^2 \partial_x^n (x^{2n+2}\tau^2 - (2n+1)x^{2n+1}\tau + n^2x^{2n}) e^{-x\tau}) \right) dx \\
&= \tau^{(n-1)} \int_0^\infty f(x) \left( x \partial_x^n ((x^{2n+2}\tau^2 - (2n+1)x^{2n+1}\tau + n^2x^{2n}) e^{-x\tau}) + \right. \\
&\quad \left. x^2 \partial_x^n \partial_x ((x^{2n+2}\tau^2 - (2n+1)x^{2n+1}\tau + n^2x^{2n}) e^{-x\tau} x^{2n}) \right) dx \\
&= \tau^{(n-1)} \int_0^\infty f(x) \left( x \partial_x^n ((x^{2n+2}\tau^2 - (2n+1)x^{2n+1}\tau + n^2x^{2n}) e^{-x\tau}) - \right. \\
&\quad \left. x^2 \partial_x^n (\tau^3 x^{2n+2} - (4n+3)\tau^2 x^{2n+1} + (5n^2+4n+1)\tau x^{2n} - (2n^3)x^{2n-1}) e^{-x\tau} \right) dx \\
&= \tau^{(n-1)} \int_0^\infty f(x) \sum_{\nu=0}^n \binom{n}{\nu} (-\tau)^\nu \frac{\Gamma(2n+1)}{\Gamma(n+\nu+1)} e^{-x\tau} \\
&\quad \left( x \partial_x^{n-\nu} (x^{2n+2}\tau^2 - (2n+1)x^{2n+1}\tau + n^2x^{2n}) - x^2 \partial_x^{n-\nu} (\tau^3 x^{2n+2} \right. \\
&\quad \left. - (4n+3)\tau^2 x^{2n+1} + (5n^2+4n+1)\tau x^{2n} - (2n^3)x^{2n-1}) \right) dx \\
&= \int_0^\infty f(x) \sum_{\nu=0}^n (-1)^{\nu+1} \binom{n}{\nu} \frac{\Gamma(2n+1)}{\Gamma(n+\nu+1)} e^{-x\tau} x^2 (x\tau)^{n+\nu-1} \\
&\quad \times \left( \frac{(4n^2+6n+2)}{(n+\nu+1)(n+\nu+2)} (\tau x)^3 - \right. \\
&\quad \frac{(8+3\nu+29n+10n\nu+30n^2+8n^2\nu+8n^3)}{(n+\nu+1)(n+\nu+2)} (\tau x)^2 + \\
&\quad \left. \frac{(3+\nu+11n+4n\nu+13n^2+5n^2\nu+5n^3)}{(n+\nu+1)} (\tau x) - n^2(n+\nu+1) \right) dx \\
&= \int_0^\infty f(x) e^{-\tau x} Q_N(x, \tau) dx. \tag{18}
\end{aligned}$$

Therefore, from Lemma 3 we have the desired real inversion formula (2).

## 4 Concluding Remark

The integrals (11) are effectively computable by using the Mellin transform

$$(Mf)(q-it) = \int_0^\infty f(x) x^{q-it-1} dx.$$

Indeed, note the identity

$$\begin{aligned} & 2\pi \int_0^\infty |D_n f(x)|^2 x^{2q-1} dx \\ &= \int_{-\infty}^\infty |(Mf)(q-it)|^2 (q^2+t^2)^2 \{(q+1)^2+t^2\} \\ & \quad \cdots \{(q+n-1)^2+t^2\} dt \quad (q > 0) \end{aligned}$$

([10], page 207, (28)). Hence,

$$\begin{aligned} & 2\pi \int_0^\infty |D_n f(x)|^2 dx \\ &= \int_{-\infty}^\infty |(Mf)\left(\frac{1}{2}-it\right)|^2 \left\{\left(\frac{1}{2}\right)^2+t^2\right\}^2 \left\{\left(\frac{1}{2}+1\right)^2+t^2\right\} \\ & \quad \cdots \left\{\left(\frac{1}{2}+n-1\right)^2+t^2\right\} dt, \end{aligned}$$

and so, the first part of (5) is

$$\begin{aligned} & \sum_{n=0}^\infty \frac{1}{n!(n+1)!} \int_0^\infty |D_n f(x)|^2 dx \\ &= \frac{1}{2\pi} \sum_{n=0}^\infty \frac{1}{n!(n+1)!} \int_{-\infty}^\infty |(Mf)\left(\frac{1}{2}-it\right)|^2 \\ & \quad \times \frac{|\Gamma(\frac{1}{2}+n+it)|^2}{|\Gamma(\frac{1}{2}+it)|^2} dt. \end{aligned} \tag{19}$$

The second part of (11) can be handled similarly by using the transformation rule in the Mellin transform

$$M(xf(x))(q-it) = (Mf)(q+1-it).$$

The series in (19) is estimated by the behavior of the Mellin transform  $(Mf)(\frac{1}{2}-it)$  at infinity, in some cases by using the formulas

$$\int_0^\infty |\Gamma(a+ix)|^2 dx = \frac{\pi}{2^{2a}} \Gamma(2a) \quad (a > 0)$$

and

$$\begin{aligned} & \int_0^\infty |\Gamma(a+ix)\Gamma(b+ix)|^2 dx \\ &= \frac{\sqrt{\pi}\Gamma(a)\Gamma(a+\frac{1}{2})\Gamma(b)\Gamma(b+\frac{1}{2})\Gamma(a+b)}{2\Gamma(a+b+\frac{1}{2})} \quad (a, b > 0) \end{aligned}$$

([1], page 655).



## References

- [1] ABRAMOWITZ, M., STEGUN, I.A., *Handbook of Mathematical Functions with Formulas, Graphs and Mathematical Tables*, U.S. Government Printing Office, Washington D.C., 1972.
- [2] AMANO, K., SAITOH, S., YAMAMOTO, M., *Error Estimates of the Real Inversion Formulas of the Laplace Transform*, Preprint Series, Graduate School of Mathematical Sciences, The University of Tokyo, 98-29.
- [3] BYUN, D.-W., SAITOH, S., *A Real Inversion Formula for the Laplace Transform*, *Zeitschrift für Analysis und ihre Anwendungen* 12(1993), 597-603.
- [4] KAJIWARA, J., TSUJI, M., *Program for the numerical analysis of inverse formula for the Laplace Transform*, Proceedings of the Second Korean-Japanese Colloquium on Finite and Infinite Dimensional Complex Analysis (1994), 93-107.
- [5] KAJIWARA, J., TSUJI, M., *Inverse Formula for Laplace Transform*, Proceedings of the 5th International Colloquium on Differential Equation, pp.163-172, VSP-Holland, 1995.
- [6] GRADSHTEYN, I. S., RYZHIK, I. M., *Table of Integrals, Series, and Products*, Academic Press, New York, 1980.
- [7] RAMM, A. G., *Multidimensional Inverse Scattering Problems*, Pitman Monographs and Surveys in Pure and Applied Mathematics, 51, Longman Scientific and Technical, UK (1992).
- [8] SAITOH, S., *Hilbert Spaces Induced by Hilbert Space Valued Functions*, Proc. Amer Math Soc., 89(1983), 74-78.
- [9] SAITOH, S., *One Approach to Some General Integral Transforms and its Applications*, *Integral Transforms and Special Functions* 3(1995), 49-84.
- [10] SAITOH, S., *Integral Transforms, Reproducing Kernels and their Applications*, Pitman Research Notes in Mathematical Series 369 (1997), Addison Wesley Longman, UK.
- [11] SAITOH, S., YAMAMOTO, M., *Stability of Lipschitz Type in Determination of Initial Heat Distribution*, *Journal of Inequalities and Applications* 1 (1997), 73-83