

## UNIFORMLY SHADOWING PROPERTY

都立大理 花井 博史 (Hirofumi Hanai)

東京理大理工 岡 正俊 (Masatoshi Oka)

Let  $M$  be a closed  $C^\infty$  manifold and  $C^r(M)$  be the set of all  $C^r$ -differentiable maps endowed with the  $C^r$ -topology ( $r \geq 1$ ).  $D_x f$  is the derivative of  $f$  at  $x$ . Denote as  $\tilde{M}$  the topological product space  $\prod_{-\infty}^{\infty} M$  and define a compatible metric  $\tilde{d}$  on  $\tilde{M}$  by  $\tilde{d}((x_n), (y_n)) = \sum_{-\infty}^{\infty} d(x_n, y_n)/2^{|n|}$  for  $(x_n), (y_n) \in \tilde{M}$ , where  $d$  is a metric on  $M$  induced by a Riemannian metric. We define a continuous map  $\tilde{f} : \tilde{M} \rightarrow \tilde{M}$  by

$$\tilde{f}((x_n)) = (f(x_n)).$$

Then the projection  $P^0 : \tilde{M} \rightarrow M$  defined by  $P^0((x_n)) = x_0$  satisfies  $P^0 \circ \tilde{f} = f \circ P^0$ . For a subset  $\Lambda$  an  $\tilde{f}$ -invariant set  $\Lambda_f$  is defined by

$$\Lambda_f = \{(x_n) \in \tilde{M} : x_n \in \Lambda, f(x_n) = x_{n+1}, n \in \mathbb{Z}\}.$$

If  $\Lambda_f \neq \emptyset$  then  $\tilde{f}|_{\Lambda_f} : \Lambda_f \rightarrow \Lambda_f$  is a surjective homeomorphism. Remark that  $\Lambda_f = M_f \neq \emptyset$  when  $\Lambda = M$ . We say that each element of  $M_f$  is an orbit of  $f$ .

For  $\delta \geq 0$  a sequence  $\{x^i\}_{i \in \mathbb{Z}} \subset M$  is called a  $\delta$ -pseudo-orbit of  $f$  if  $d(f(x^i), x^{i+1}) \leq \delta$  for every  $i \in \mathbb{Z}$ . A sequence  $\{x^i\}_{i \in \mathbb{Z}} \subset M$  is said to be  $\varepsilon$ -traced by an orbit  $(y_i) \in M_f$  if  $d(x^i, y_i) < \varepsilon$  for every  $i \in \mathbb{Z}$ . We say that  $f$  has the *shadowing property* if for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that every  $\delta$ -pseudo-orbit of  $f$  can be  $\varepsilon$ -traced by an orbit of  $f$ .

A sequence  $\{\tilde{x}^i\}_{i \in \mathbb{Z}} \subset M_f$  is called an orbit of  $\tilde{f}$  if  $\tilde{f}(\tilde{x}^i) = \tilde{x}^{i+1}$ . For  $\delta \geq 0$  a sequence  $\{\tilde{x}^i\}_{i \in \mathbb{Z}} \subset M_f$  is a  $\delta$ -pseudo-orbit of  $\tilde{f}$  if  $\tilde{d}(\tilde{f}(\tilde{x}^i), \tilde{x}^{i+1}) \leq \delta$  for every  $i \in \mathbb{Z}$ . A sequence  $\{\tilde{x}^i\}_{i \in \mathbb{Z}} \subset \tilde{M}$  is said to be  $\varepsilon$ -traced by an orbit  $(\tilde{y}^i)$  of  $\tilde{f}$  if  $\tilde{d}(\tilde{x}^i, \tilde{y}^i) < \varepsilon$  for every  $i \in \mathbb{Z}$ . We say that  $\tilde{f}$  satisfies the *shadowing property* if for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that every  $\delta$ -pseudo-orbit of  $\tilde{f}$  can be  $\varepsilon$ -traced by an orbit of  $\tilde{f}$ .

We say that  $\tilde{f}$  satisfies  *$C^r$  uniformly shadowing property* if there is a neighborhood  $\mathcal{U}(f)$  of  $f$  in  $C^r(M)$  with the property that for  $\varepsilon > 0$  there is  $\delta > 0$  such that for  $g \in \mathcal{U}(f)$  every  $\delta$ -pseudo-orbit of  $\tilde{g}$  is  $\varepsilon$ -traced by an orbit of  $\tilde{g}$ .

Let  $\pi : TM \rightarrow M$  be a tangent bundle of  $M$  and  $\|\cdot\|$  be a Riemannian metric on  $TM$ . Define a subset of the product topological space  $\tilde{M} \times TM$  by

$$T\tilde{M} = \{(\tilde{x}, v) \in \tilde{M} \times TM : P^0(\tilde{x}) = \pi(v)\}$$

and define a Finsler  $\|\cdot\|$  on  $T\tilde{M}$  by  $\|(\tilde{x}, v)\| = \|v\|$ . Then  $\tilde{\pi} : T\tilde{M} \rightarrow \tilde{M}$  defined by  $\tilde{\pi}(\tilde{x}, v) = \tilde{x}$  is a  $C^0$ -vector bundle over  $\tilde{M}$ . Define the projection  $\bar{P}^0 : T\tilde{M} \rightarrow TM$  by  $\bar{P}^0(\tilde{x}, v) = v$ . Then,

$$\bar{P}^0|_{T_{\tilde{x}}\tilde{M}} : T_{\tilde{x}}\tilde{M} \rightarrow T_{P^0(\tilde{x})}M$$

is a linear isomorphism where  $T_{\tilde{x}}\tilde{M} = \tilde{\pi}^{-1}(\tilde{x})$ . A linear bundle map  $D\tilde{f} : T\tilde{M} \rightarrow T\tilde{M}$  covering  $\tilde{f}$  is defined by

$$D\tilde{f}(\tilde{x}, v) = (\tilde{f}(\tilde{x}), D_{P^0(\tilde{x})}f(v)).$$

Then we have  $D\tilde{f}(T_{\tilde{x}}\tilde{M}) \subset T_{\tilde{f}(\tilde{x})}\tilde{M}$  and  $\bar{P}^0 \circ D\tilde{f} = Df \circ \bar{P}^0$ . To simplify the notation we write  $D_{\tilde{x}}\tilde{f} = D\tilde{f}|_{T_{\tilde{x}}\tilde{M}}$ . For a subset  $\tilde{\Lambda}$  define

$$T\tilde{M}|_{\tilde{\Lambda}} = \bigcup_{\tilde{x} \in \tilde{\Lambda}} T_{\tilde{x}}\tilde{M}.$$

A closed  $f$ -invariant set  $\Lambda$  ( $f(\Lambda) = \Lambda$ ) is said to be *hyperbolic* if  $T\tilde{M}|_{\Lambda_f}$  splits into the Whitney sum  $T\tilde{M}|_{\Lambda_f} = \mathbb{E}^s \oplus \mathbb{E}^u$  of subbundles  $\mathbb{E}^s$  and  $\mathbb{E}^u$ , and there are  $C > 0$  and  $0 < \lambda < 1$  such that

- (i)  $D\tilde{f}(\mathbb{E}^s) \subset \mathbb{E}^s$  and  $D\tilde{f}(\mathbb{E}^u) = \mathbb{E}^u$ ,
- (ii)  $D\tilde{f}|_{\mathbb{E}^u} : \mathbb{E}^u \rightarrow \mathbb{E}^u$  is invertible,
- (iii)  $\|D\tilde{f}^n|_{\mathbb{E}^s}\| \leq C\lambda^n$  and  $\|(D\tilde{f}|_{\mathbb{E}^u})^{-n}\| \leq C\lambda^n$  for  $n \geq 0$ ,

where  $\|T\|$  denotes the supremum norm of a linear bundle map  $T$ . The number  $\lambda$  is called the *skewness* of the hyperbolic set  $\Lambda$ . For  $\varepsilon > 0$  and

$\tilde{x} \in M_f$  the *local stable* and the *local unstable manifolds* are defined by

$$W_\varepsilon^s(\tilde{x}, f) = \{y \in M : d(x_n, f^n(y)) \leq \varepsilon \text{ for } n \geq 0\},$$

$$W_\varepsilon^u(\tilde{x}, f) = \left\{ y \in M \left| \begin{array}{l} \text{there exists } \tilde{y} \in M_f \text{ such that } y_0 = y \\ \text{and } d(x_{-n}, y_{-n}) \leq \varepsilon \text{ for } n \geq 0 \end{array} \right. \right\}.$$

Then,  $W_\varepsilon^s(\tilde{x}, f) = W_\varepsilon^s(\tilde{y}, f)$  for  $\tilde{x}, \tilde{y} \in M_f$  with  $x_0 = y_0$ .

For  $\tilde{x} \in M_f$  the *stable* and the *unstable sets* are defined by

$$W^s(\tilde{x}, f) = \{y \in M : \lim_{n \rightarrow \infty} d(x_n, f^n(y)) = 0\},$$

$$W^u(\tilde{x}, f) = \left\{ y \in M \left| \begin{array}{l} \text{there is } \tilde{y} \in M_f \text{ satisfying } y_0 = y \\ \text{and } \lim_{n \rightarrow \infty} d(x_{-n}, y_{-n}) = 0 \end{array} \right. \right\}.$$

Then,  $W^s(\tilde{x}, f) = W^s(\tilde{y}, f)$  for  $\tilde{x}, \tilde{y} \in M_f$  with  $x_0 = y_0$ . If  $\Lambda$  is a hyperbolic set, for  $\tilde{x} \in \Lambda_f$  we have

$$W^s(\tilde{x}, f) = \bigcup_{n=0}^{\infty} f^{-n}(W_\varepsilon^s(\tilde{f}^n(\tilde{x}), f)), \quad W^u(\tilde{x}, f) = \bigcup_{n=0}^{\infty} f^n(W_\varepsilon^u(\tilde{f}^{-n}(\tilde{x}), f)).$$

Remark that  $W^\sigma(\tilde{x}, f)$  ( $\sigma = s, u$ ) are not always the manifolds like the stable and unstable manifolds given by diffeomorphisms. However we can define the transversality condition between  $W^s(\tilde{x}, f)$  and  $W^u(\tilde{y}, f)$  as follows.

Let  $\tilde{y}$  and  $\tilde{z}$  be points in  $\Lambda_f$ . We say that  $W^s(\tilde{y}, f)$  is *transversal* to  $W^u(\tilde{z}, f)$  if  $f^{n+m} | W_\varepsilon^u(\tilde{f}^{-m}(\tilde{z}), f)$  is transversal to  $W_\varepsilon^s(\tilde{f}^n(\tilde{y}), f)$  for  $\varepsilon > 0$  small enough and  $n, m \geq 0$ .

The non-wandering set  $\Omega(f)$  is defined by

$$\Omega(f) = \left\{ x \in M \left| \begin{array}{l} \text{for any neighborhood } U \text{ of } x \text{ there is } n > 0 \\ \text{satisfying } U \cap f^n(U) \neq \emptyset \end{array} \right. \right\}.$$

Obviously,  $\Omega(f)$  is closed and satisfies that  $f(\Omega(f)) \subset \Omega(f)$  and  $Per(f) \subset \Omega(f)$ , where  $Per(f)$  denotes the set of all periodic points of  $f$ . A differentiable map  $f$  is said to satisfy *Axiom A* if

- (i)  $Per(f)$  is dense in  $\Omega(f)$ ,
- (ii)  $\Omega(f)$  is hyperbolic.

We say that an Axiom A differentiable map  $f$  satisfies the *strong transversality* if  $W^s(\tilde{y}, f)$  is transversal to  $W^u(\tilde{z}, f)$  for  $\tilde{y}, \tilde{z} \in \Omega(f)_f$ .

**Theorem.** If  $C^1$ -differentiable map  $f$  satisfies both Axiom A and the strong transversality, then  $\tilde{f}$  satisfies  $C^1$  uniformly shadowing property.

This result was proved by Sakai for the class of  $C^1$ -diffeomorphisms. The full proof of our theorem will appear elsewhere.

#### REFERENCES

- [1] N.Aoki, K.Moriyasu and M.Oka, Differentiable maps having hyperbolic sets, *Topology Appl.* 82(1998) 15 – 48.
- [2] N.Aoki, K.Moriyasu and N.Sumii,  $C^1$  maps having hyperbolic periodic points, to appear in *Fund. Math.*
- [3] J.Guckenheimer, J.Moser and S.Newhouse, *Dynamical Systems* (Birkhäuser, Boston, 1980).

- [4] M.Hirsch, J.Palis, C.Pugh and M.Shub, Neighborhoods of hyperbolic sets, *Inv. Math.* 9 (1970) 121–134.
- [5] M.Hirsch, C.Pugh and M.Shub, *Invariant manifolds*, Lecture Note in Math. 583 (Springer, Berlin, 1977).
- [6] K.Moriyasu, The topological stability of diffeomorphisms, *Nagoya Math. J.*123 (1991) 91–102.
- [7] K.Moriyasu, Axiom A endomorphisms having no cycle, in: N.Aoki, K.Shiraiwa, Y.Takahashi, ed., *Proceedings of the International Conference on Dynamical Systems and Chaos 1* (World Scientific, Singapore, 1995) 182–186.
- [8] K.Moriyasu and M.Oka, The creation of homoclinic points of  $C^1$ -maps, *Topology Appl.* 54 (1993) 47–64.
- [9] K.Moriyasu and M.Oka, Pseudo orbit tracing property and strong transversality of  $C^1$  maps, in: N.Aoki, K.Shiraiwa, Y.Takahashi, ed., *Proceedings of the International Conference on Dynamical Systems and Chaos 1* (World Scientific, Singapore, 1995) 187–191.
- [10] F.Przytycki, Anosov endomorphisms, *Studia Math.* 58 (1976) 249–285.
- [11] F.Przytycki, On  $\Omega$ -stability and structural stability of endomorphisms satisfying Axiom A, *Studia Math.* 60 (1977) 61–77.
- [12] C.Robinson, Stability theorem and hyperbolicity in dynamical systems, *Rocky Mountain J. Math.* 7 (1977) 425–437.
- [13] C.Robinson, Structural stability of  $C^1$  diffeomorphisms, *J. Diff. Equat.* 22 (1976) 28–73.
- [14] K.Sakai, The  $C^1$  uniform pseudo-orbit tracing property, *Tokyo J. Math.* 15 (1992) 99–109.
- [15] K.Sakai,  $C^r$  Axiom A maps having strong transversality, *Topology Appl.* 54 (1993) 173–182.
- [16] K.Sakai, Pseudo orbit tracing property and strong transversality, *Osaka J. Math.* 31 (1994) 373–386.