

## A SOLUTION TO A PROBLEM OF TEODOR PRZYMUSIŃSKI

VALENTIN GUTEV

A subset  $A$  of a space  $X$  is  $C^*$ -embedded in  $X$  if every bounded real-valued continuous function on  $A$  is continuously extendable to the whole of  $X$ . If this holds for all real-valued continuous functions on  $A$ , then  $A$  is  $C$ -embedded in  $X$ .

The present note provides detailed suggestions to the solution of the following problem. For a *non-discrete metric* space  $M$  and a subset  $A$  of a space  $X$ , does the  $C^*$ -embedding of  $A \times M$  in  $X \times M$  imply that it is also  $C$ -embedded in  $X \times M$ , i.e.

$$A \times M \overset{C^*}{\hookrightarrow} X \times M \quad \implies \quad A \times M \overset{C}{\hookrightarrow} X \times M \quad ?$$

The problem was stated as Problem 3 of [T. Przymusiński, *Notes on extendability of continuous functions from products with a metric factor*, unpublished note, May 1983], later on as Problem 4.14 of [T. Hoshina, *Extensions of mappings II*, Topics in General Topology (K. Morita and J. Nagata, eds.), North-Holland, Amsterdam, 1989, pp. 41–80] and Problem 3.1 of [T. Hoshina, *Extensions of mappings*, Recent Progress in General Topology (M. Hušek and J. van Mill, eds.), North-Holland, Amsterdam, 1992, pp. 405–416].

### THE SOLUTION

To state the main result we call in use also the following imbedding-like properties. Let  $\lambda$  be an infinite cardinal number.

$P^\lambda$ -embedding: A subset  $A$  of a space  $X$  is  $P^\lambda$ -embedded in  $X$ , or briefly  $A \overset{P^\lambda}{\hookrightarrow} X$ , if every continuous  $f : A \rightarrow Y$  in a Banach space  $Y$  of  $w(Y) \leq \lambda$  is continuously extendable to the whole of  $X$ .

$U^\omega$ -embedding: A subset  $A$  of a space  $X$  is  $U^\omega$ -embedded in  $X$ , or briefly  $A \overset{U^\omega}{\hookrightarrow} X$ , if for every continuous  $f : A \rightarrow \mathbb{R}$  there exists a continuous  $g : X \rightarrow \mathbb{R}$  with  $f(x) \leq g(x)$  whenever  $x \in A$ .

It should be mentioned that  $A$  is  $C$ -embedded in  $X$  if and only if it is  $P^\omega$ -embedded in  $X$ , while  $A$  is  $P^\omega$ -embedded in  $X$  if and only if it is both  $U^\omega$ - and  $C^*$ -embedded in  $X$ . That is, always

$$C = P^\omega = U^\omega + C^*.$$

The following recent result was obtained together with Haruto Ohta.

**Theorem.** *For a  $P^\lambda$ -embedded subset  $A$  of a space  $X$  and a metric space  $M$ , the following conditions are equivalent*

- (a)  $A \times M \xrightarrow{P^\lambda} X \times M$
- (b)  $A \times M \xrightarrow{C^*} X \times M$
- (c)  $A \times M \xrightarrow{U^\omega} X \times M$

Note that  $A \times M \xrightarrow{C^*} X \times M$  implies  $A \xrightarrow{C} X$  provided  $M$  is non-discrete because, in this case,  $M$  contains an infinite compact subset. Hence, the above result provides a complete positive solution to the problem of interest. For the proper understanding of this theorem, a word should be said also about the last condition (c). The statement that it is equivalent to the previous ones should be compared with Rudin-Starbird's result that, for a non-discrete metric space  $M$ , the normality of  $X \times M$  implies the countable paracompactness of  $X \times M$ . Namely, the  $U^\omega$ -embedding has a quite nice and useful reading just in terms of Ishikawa's characterization of countable paracompactness.

#### ON THE WAY TO THE PROOF

Special cases of (a)  $\Leftrightarrow$  (b):  $X \times M$  an  $M$ -independent product and  $\lambda = \omega$  (Przymusiński, 1983);  $M = \mathbb{P}$  the space of irrational numbers and  $\lambda = \omega$  (Ohta, 1993);  $M$   $\sigma$ -locally compact (Yamazaki, 1997);  $M^2$  homeomorphic to  $M$  (Hoshina and Yamazaki, 199?).

#### FIRST STEP: A reduction to "nice" metric factors

For a space  $Y$ , let  $\mathcal{P}(Y)$  be the set of all closed subsets of  $Y$ . Let  $A$ ,  $X$  and  $M$  be as in our theorem. To  $M$  we associate the family of *all solutions*, or the *Przymusiński* family for  $M$ , by

$$\mathfrak{B} = \{S \subset M : A \times S \xrightarrow{P^\lambda} X \times S\}.$$

The following important fact will play a central role in this part of the proof.

**Fact 1** (Michael).  $S \in \mathfrak{B} \implies \mathcal{P}(S) \subset \mathfrak{B}$ .

It will be useful to illustrate the idea first on a partial case. For the purpose, let  $M^{(\mathcal{K},0)} = M$ , and, for every ordinal  $\alpha > 0$ , let

$$M^{(\mathcal{K},\alpha)} = X \setminus \bigcup \{K \subset M \text{ compact} : K \subset M^{(\mathcal{K},\beta)} \text{ is open for some } \beta < \alpha\}.$$

Take an ordinal  $\gamma$  with  $M^{(\mathcal{K},\gamma)} = M^{(\mathcal{K},\gamma+1)}$ . Then,

1.  $M^{(\mathcal{K},\gamma)} \in \mathcal{P}(M)$  is nowhere locally compact;
2.  $M \setminus M^{(\mathcal{K},\gamma)}$  is  $\sigma$ -locally compact.

Now, suppose that  $M$  is a Polish space with  $\dim(M) = 0$ . Then, relying on the known partial solution and Fact 1, we get the following series of implications.

$$M^{(\mathcal{K},\gamma)} = \emptyset \implies M \text{ is } \sigma\text{-locally compact} \implies M \in \mathfrak{P}.$$

On the other hand,

$$\begin{aligned} M^{(\mathcal{K},\gamma)} \neq \emptyset &\implies M^{(\mathcal{K},\gamma)} = \mathbb{P} \\ &\Downarrow \\ &M^{(\mathcal{K},\gamma)} \in \mathfrak{P} \\ &\Downarrow \\ M \in \mathcal{P}(\mathbb{P}) = \mathcal{P}(M^{(\mathcal{K},\gamma)}) &\subset \mathfrak{P}. \end{aligned}$$

That is, always  $M \in \mathfrak{P}$ .

Let  $\mathcal{K} = \{S \in \mathcal{P}(M) : S \text{ is compact}\}$ . Then, by the known results,  $\mathcal{K} \subset \mathfrak{P}$ . On the other hand,  $M^{(\mathcal{K},\gamma)}$  is a resulting set by a  $\mathcal{K}$ -scattered procedure and, hence, a procedure that is *scattered* also with respect to a part of the members of  $\mathfrak{P}$ . This arguments suggest that, for a better result, we need to call in use all members of  $\mathfrak{P}$ , i.e. to arrange a  $\mathfrak{P}$ -scattered procedure on  $M$ .

Turning to this case, we change our definition as follows. Let  $S \subset M$ , and let  $S^{(\mathfrak{P},0)} = S$ . Next, for any ordinal  $\alpha > 0$ , we consider the set

$$S^{(\mathfrak{P},\alpha)} = S \setminus \bigcup \{U \subset S : U \text{ is open and } \text{cl}_S(U) \cap S^{(\mathfrak{P},\beta)} \in \mathfrak{P} \text{ for some } \beta < \alpha\}.$$

Suppose that  $M \notin \mathfrak{P}$ , and let  $S \in \mathcal{P}(M) \setminus \mathfrak{P}$  be such that

$$w(S) = \min\{w(F) : F \in \mathcal{P}(M) \setminus \mathfrak{P}\}.$$

Then, as before, take an ordinal  $\gamma$  with  $S^{(\mathfrak{P},\gamma)} = S^{(\mathfrak{P},\gamma+1)}$ . As a result, we get that

1.  $S^{(\mathfrak{P},\gamma)} \in \mathcal{P}(S)$  is **weight-homogeneous**, that is,  $w(U) = w(S)$  for every non-empty open  $U \subset S$ ;
2.  $S \setminus S^{(\mathfrak{P},\gamma)}$  has a  $\sigma$ -discrete closed cover  $\Sigma \subset \mathfrak{P}$ .

On the other hand, for the members of  $\mathfrak{P}$ , we have that

**Fact 2.**  $\mathcal{D} \subset \mathfrak{P}$  discrete in  $\bigcup \mathcal{D} \implies \bigcup \mathcal{D} \in \mathfrak{P}$ .

In view of our next arguments, let us make the following

**Assumption.**  $S^{(\mathfrak{P}, \gamma)} \in \mathfrak{P}$ .

As a result, we now get that

**Conclusion 3.** There exists a countable cover  $\mathcal{F}$  of  $S$  with  $\mathcal{F} \subset \mathcal{P}(S) \cap \mathfrak{P}$ .

**Conclusion 4.**  $A \times S \xrightarrow{\text{well}} X \times S$ .

Here,  $A \times S \xrightarrow{\text{well}} X \times S$  if  $A \times S$  is completely separated from any zero-set of  $X \times S$  which doesn't meet  $A \times S$ . To involve Conclusion 4, we also need the following *weak embedding* properties:

$C_1$ -embedding: A subset  $B$  of  $Y$  is  $C_1$ -embedded in  $Y$ , or briefly  $B \xrightarrow{C_1} Y$ , if  $F \xrightarrow{\text{well}} Y$  for every zero-set  $F$  of  $B$ . That is, for any zero-set  $F$  of  $B$  and any zero-set  $Z$  of  $Y$ , with  $Z \cap F = \emptyset$ , there exists a zero-set  $Z_F$  of  $Y$  such that  $F \subset Z_F$  and  $Z_F \cap Z = \emptyset$ .

$CU$ -embedding: A subset  $B$  of  $Y$  is  $CU$ -embedded in  $Y$ , or briefly  $B \xrightarrow{CU} Y$ , if for any zero-set  $F$  of  $B$  and any zero-set  $Z$  of  $Y$ , with  $Z \cap F = \emptyset$ , there exists a zero-set  $Z_F$  of  $Y$  such that  $F \subset Z_F$  and  $Z_F \cap Z \cap B = \emptyset$ .

The relations between our weak-embedding properties could be now summarized into the following diagram.

**Observation 5.**

$$\begin{array}{ccc} C^* & & U^\omega \\ & \searrow & \swarrow \\ C_1 & = & CU + \text{well} \end{array}$$

Then, by Conclusion 4, we have

**Conclusion 6.**  $A \times S \xrightarrow{C_1} X \times S$ .

According to Conclusion 3, this implies

**Final Conclusion.**  $S \in \mathfrak{P}$ .

The so obtained contradiction provides the following result which accomplishes the first step of the proof of our theorem.

**Theorem A.**  $M \in \mathfrak{P}$  provided  $S \in \mathfrak{P}$  for any weight-homogeneous and nowhere locally compact  $S \in \mathcal{P}(M)$ .

**SECOND STEP: Separating the factors**

NOTATIONS: For sets  $D$  and  $R$ , let  $R^D$  denote all maps from  $D$  to  $R$ , and  $2^R$  — all subsets of  $R$ . For cardinals  $\kappa$  and  $\mu$ , let  $\kappa^{<\mu} = \bigcup\{\kappa^\delta : \delta < \mu\}$ . For reasons of convenience, we regard  $\kappa^0$  as the singleton  $\{\emptyset\}$ . To every  $\sigma \in \kappa^\delta$  and  $\alpha < \kappa$  we associate another map  $\sigma \hat{\alpha} \in \kappa^{\delta+1}$  defined by  $\sigma \hat{\alpha}|_\delta = \sigma$  and  $\sigma \hat{\alpha}(\delta) = \alpha$ . Also, to every  $\mathcal{H} : T \rightarrow (2^R)^D$  we associate another map  $\langle \mathcal{H}, D \rangle : T \rightarrow 2^R$  defined by

$$\langle \mathcal{H}, D \rangle(t) = \bigcup \mathcal{H}[t](D) \quad \forall t \in T.$$

Finally, for a space  $Y$ , we shall use  $\text{coz}(Y)$  to denote the collection of all *cozero-sets* of  $Y$  and  $\text{zero}(Y)$  for that of all *zero-sets* of  $Y$ .

**CONCEPTS:**

Monotone decreasing map:  $\mathcal{H} : \kappa^{<\omega} \rightarrow (2^R)^D$  if  $\mathcal{H}[\sigma \hat{\alpha}](D)$  refines  $\mathcal{H}[\sigma](D)$  for every  $\sigma \in \kappa^{<\omega}$  and  $\alpha < \kappa$ .

Sieve:  $\mathcal{S} : \kappa^{<\omega} \rightarrow \text{coz}(Y)$  if  $\mathcal{S}(\emptyset) = Y$  and  $\mathcal{S}(\sigma) = \bigcup\{\mathcal{S}(\sigma \hat{\alpha}) : \alpha < \kappa\}$  for every  $\sigma \in \kappa^{<\omega}$ .

Strong Sieve:  $\mathcal{S} : \kappa^{<\omega} \rightarrow \text{coz}(Y)$  if  $\mathcal{S}$  is a sieve such that  $\emptyset \notin \mathcal{S}(\kappa^{<\omega})$ , each family  $\mathcal{S}(\kappa^n)$ ,  $n < \omega$ , is a locally finite in  $Y$  and, whenever  $y \in \bigcap\{\mathcal{S}(t|n) : n < \omega\}$  for some  $t \in \kappa^\omega$ , the collection  $\mathcal{S}(t|n)$ ,  $n < \omega$ , stands for a local base at  $y$  in  $Y$ .

$\mathcal{S}$ -free map:  $\mathcal{G} : \kappa^{<\omega} \rightarrow (2^Y)^\kappa$ , where  $\mathcal{S}$  is a map  $\mathcal{S} : \kappa^{<\omega} \rightarrow \text{coz}(M)$ , if for every  $t \in \kappa^\omega$  we have that  $\bigcap\{\text{cl}_Y((\langle \mathcal{G}, \kappa \rangle)(t|n)) \times \mathcal{S}(t|n) : n < \omega\} = \emptyset$ .

Expansion:  $\mathcal{H} : \kappa^{<\omega} \rightarrow (2^X)^\kappa$  of  $\mathcal{G} : \kappa^{<\omega} \rightarrow (2^Y)^\kappa$ , where  $Y \subset X$ , if  $\mathcal{G}[\sigma](\alpha) = \mathcal{H}[\sigma](\alpha) \cap Y$  whenever  $\sigma \in \kappa^{<\omega}$  and  $\alpha < \kappa$ .

The second step of the proof of our theorem reads now as follows.

**Theorem B.** *Under the conditions of the main theorem, let, in addition,  $M$  be weight homogeneous and nowhere locally compact. Also, let  $w(M) = \kappa$ . Then, the following conditions are equivalent.*

- (a)  $A \times M \xrightarrow{C^*} X \times M$
- (b) Whenever  $\mathcal{S} : \kappa^{<\omega} \rightarrow \text{coz}(M)$  is a strong sieve, every monotone decreasing and  $\mathcal{S}$ -free map  $\mathcal{G} : \kappa^{<\omega} \rightarrow \text{coz}(A)^\kappa$  has a monotone decreasing and  $\mathcal{S}$ -free expansion  $\mathcal{G} : \kappa^{<\omega} \rightarrow \text{coz}(X)^\kappa$ .
- (c)  $A \times M \xrightarrow{P^\lambda} X \times M$

Here is a brief scheme of (a)  $\implies$  (b). Suppose that  $\mathcal{G} : \kappa^{<\omega} \rightarrow \text{coz}(A)^\kappa$  and  $\mathcal{S} : \kappa^{<\omega} \rightarrow \text{coz}(M)$  are as in (b). Then, the statement that  $\mathcal{G}$  is an  $\mathcal{S}$ -free map becomes equivalent to the statement that the family  $\{(\mathcal{G}, \kappa)(\sigma) \times \mathcal{S}(\sigma) : \sigma \in \kappa^{<\omega}\}$  is locally finite in  $A \times M$ . The last becomes “almost” equivalent to the existence of  $F_{(\mathcal{G}, \mathcal{S})}^0, F_{(\mathcal{G}, \mathcal{S})}^1 \in \text{zero}(A \times M)$  such that  $F_{(\mathcal{G}, \mathcal{S})}^0 \cap F_{(\mathcal{G}, \mathcal{S})}^1 = \emptyset$ . However, by (a),  $A \times M \xrightarrow{C^*} X \times M$ . Hence, there are  $Z_{(\mathcal{H}, \mathcal{S})}^0, Z_{(\mathcal{H}, \mathcal{S})}^1 \in \text{zero}(X \times M)$  such that

$$F_{(\mathcal{G}, \mathcal{S})}^i \subset Z_{(\mathcal{H}, \mathcal{S})}^i, \quad i < 2, \quad \text{and} \quad Z_{(\mathcal{H}, \mathcal{S})}^0 \cap Z_{(\mathcal{H}, \mathcal{S})}^1 = \emptyset.$$

Relying on the “almost” equivalence mentioned above, these two zero-sets of  $X \times M$  yield a monotone decreasing and  $\mathcal{S}$ -free expansion  $\mathcal{H} : \kappa^{<\omega} \rightarrow \text{coz}(X)^\kappa$  of  $\mathcal{G}$ .

Here is also a brief scheme of (b)  $\implies$  (c). This implication is based on the following chain of arguments.

**Fact 1.** There exists a strong sieve  $\mathcal{S} : \kappa^{<\omega} \rightarrow \text{coz}(M)$  on  $M$  such that

$$\mathcal{S}_n(z) = \bigcup \{ \mathcal{S}(\sigma) : \sigma \in \kappa^n \ \& \ z \in \text{cl}_M(\mathcal{S}(\sigma)) \}, \quad n < \omega,$$

constitute a local base at  $z$  for every  $z \in M$ .

A CONCEPT MORE: Let  $\mathbb{I} = [0, 1]$ .

Sieve partition of unity:  $\xi : \kappa^{<\omega} \rightarrow C(M, \mathbb{I})$ , or a function version of strong sieve, if  $\xi[\emptyset]$  is the constant function on  $M$  with the value of 1, and  $\xi[\sigma] = \sum \{ \xi[\sigma \hat{\ } \alpha] : \alpha < \kappa \}$  for every  $\sigma \in \kappa^{<\omega}$ .

**Fact 2.** For every strong sieve  $\mathcal{S} : \kappa^{<\omega} \rightarrow \text{coz}(M)$  there exists a sieve-partition of unity  $\xi : \kappa^{<\omega} \rightarrow C(M, \mathbb{I})$  such that  $\text{supp}(\xi[\sigma]) \subset \mathcal{S}(\sigma)$  for every  $\sigma \in \kappa^{<\omega}$ .

Let  $(Y, \|\cdot\|)$  be a Banach space, and let  $f : A \times M \rightarrow Y$  be a continuous map. The statement of (c) becomes now equivalent to the existence of a continuous map  $g : X \times M \rightarrow Y$  with  $g|_{A \times M} = f$ . Towards this end, for every space  $T$  we shall associate a map  $\Delta_T$

$$T \quad \longrightarrow \quad \Delta_T : C(T \times M, Y) \rightarrow C(T, Y)^{\kappa^{<\omega}}$$

that defines into the following manner. Let  $\mathcal{S} : \kappa^{<\omega} \rightarrow \text{coz}(M)$  be a strong sieve on  $M$  as in Fact 1. Take a dense  $D \subset M$  with  $|D| = \kappa$ , and then define a map  $\theta : \kappa^{<\omega} \rightarrow M$  by  $\theta(\alpha) \in D \cap \mathcal{S}(\alpha)$  for every  $\alpha \in \kappa^{<\omega}$ . Finally, our  $\Delta_T$  is defined by  $\Delta_T(h)[\sigma](x) = h(x, \theta(\sigma))$  whenever  $h \in C(T \times M, Y)$ ,  $\sigma \in \kappa^{<\omega}$  and  $x \in T$ .

The correspondence  $\Delta_T$  is “nice” invertible on the image of  $C(T \times M, Y)$  under  $\Delta_T$ . That is, one could restore in full  $h \in C(T \times M, Y)$  relying only on  $\Delta_T(h)$ . Namely,

let  $\xi : \kappa^{<\omega} \rightarrow C(M, \mathbb{I})$  be a sieve partition of unity on  $M$  as in Fact 2 applied to  $\mathcal{S}$ . Then,

$$(*) \quad h = \lim_{n \rightarrow \infty} \sum \{ \xi[\sigma] \cdot \Delta_T(h)[\sigma] : \sigma \in \kappa^n \}.$$

The idea of (b)  $\Rightarrow$  (c) could be now stated in the following abstract setting. To the map  $f$  we associate the corresponding one  $\Phi = \Delta_A(f) : \kappa^{<\omega} \rightarrow C(A, Y)$ . In this way, the correspondence  $\Delta_T$  transforms our extension problem to an extension problem for  $\Phi$ . Namely, it is now sufficient to find  $\Gamma : \kappa^{<\omega} \rightarrow C(X, Y)$  subject to the following

**Extension Condition:**

$$(EC) \quad \Gamma[\sigma] \Big|_A = \Phi[\sigma], \quad \text{for every } \sigma \in \kappa^{<\omega};$$

**Continuity Condition:**

$$(CC) \quad \Gamma \in \Delta_X(C(X \times M, Y)).$$

If one could deal with this last problem, then merely  $g = \Delta_X^{\leftarrow}(\Gamma) \in C(X \times M, Y)$  will be the required extension of  $f$ . Turning to this, let us observe that

$$A \xrightarrow{P^\lambda} X \quad \Longrightarrow \quad \text{“many” solutions of (EC)}$$

$$???????? \quad \Longrightarrow \quad \text{at least one solution of (CC)}$$

To discover the nature of (CC) we call in use (\*) and thus we get the following its more concrete setting:

$$(CC)^* \quad \lim_{n \rightarrow \infty} \sum \{ \xi[\sigma] \cdot \Gamma[\sigma] : \sigma \in \kappa^n \} \in C(X \times M, Y).$$

We are now ready for the final realization of this implication. Namely, the hidden property “????????” becomes the **controlled** extending of monotone decreasing  $\mathcal{S}$ -free maps. That is, just these maps will take care about the control on (CC). Briefly, to the map  $\Phi$  we associate a sequence  $\{\mathcal{F}_\ell : \ell < \omega\}$  of monotone decreasing and  $\mathcal{S}$ -free maps  $\mathcal{F}_\ell : \kappa^{<\omega} \rightarrow \text{coz}(A)^\kappa$ . According to (b), each  $\mathcal{F}_\ell$  admits a monotone decreasing and  $\mathcal{S}$ -free expansion  $\mathcal{G}_\ell : \kappa^{<\omega} \rightarrow \text{coz}(X)^\kappa$ .

The fact that  $\Phi = \Delta_A(f)$  could be now stated as

$$\begin{aligned} & \ell < \omega, \quad m \leq n < \omega \quad \& \quad \sigma \in \kappa^n \\ & \Downarrow \\ & \|\Phi[\sigma](x) - \Phi[\sigma|m](x)\| \leq \frac{1}{2^{\ell+1}} \quad \forall x \in A \setminus \langle \mathcal{F}_\ell, \kappa \rangle(\sigma|m) \end{aligned}$$

Relying on this, we finally construct  $\Gamma$  just satisfying the same condition, i.e. such that

$$\begin{aligned}
 (\text{CC})^{**} \quad & \ell \leq m \leq n < \omega \quad \& \quad \sigma \in \kappa^n \\
 & \Downarrow \\
 & \|\Gamma[\sigma](x) - \Gamma[\sigma|_m](x)\| \leq \frac{1}{2^{\ell+1}} \quad \forall x \in X \setminus (\mathcal{G}_\ell, \kappa)(\sigma|_m)
 \end{aligned}$$