

THE COMPLETIONS OF METRIC ANR'S AND UNIFORM ANR'S

KATSURO SAKAI

Institute of Mathematics
University of Tsukuba

A subset Y of a space X is said to be *homotopy dense* in X if there exists a homotopy $h: X \times [0, 1] \rightarrow X$ such that $h_0 = \text{id}$ and $h_t(X) \subset Y$ for $t > 0$. This concept is very important in ANR Theory and Infinite-Dimensional Topology. When X is an ANR, the concept of the homotopy denseness is dual to the one of local homotopy negligibility introduced by Toruńczyk in [To₃], that is, $Y \subset X$ is homotopy dense in X if and only if the complement $X \setminus Y$ is locally homotopy negligible in X (cf. [To₃, Theorem 2.4]). The following fact is well-known:

Fact. *Every homotopy dense subset of an ANR is also an ANR and a metrizable space is an ANR if it contains an ANR as a homotopy dense subset.*

The lack of the homotopy denseness of a metric ANR in its completion often destroys the ANR property of the completion. For instance, the $\sin \frac{1}{x}$ -curve in the plane \mathbb{R}^2 is an ANR but the completion of this curve (= the closure in \mathbb{R}^2) is not an ANR. Moreover, even if the completion is an ANR, it is very different from the original ANR. The circle \mathbf{S}^1 is the completion of the space $\mathbf{S}^1 \setminus \{\text{pt}\}$ and the both spaces are ANR but they are topologically very different from each other. It should be remarked that $\mathbf{S}^1 \setminus \{\text{pt}\}$ is not homotopy dense in \mathbf{S}^1 . In this note,¹ we consider the following interesting problem:

Problem. *When is a metric ANR homotopy dense in the metric completion?*

1. A CHARACTERIZATION OF METRIC ANR'S

The nerve of an open cover \mathcal{V} of a space X is denoted by $N(\mathcal{V})$. A sequence $\mathcal{U} = (\mathcal{U}_n)_{n \in \mathbb{N}}$ of open covers of a metric space X is called a *zero-sequence* if $\lim_{n \rightarrow \infty} \text{mesh} \mathcal{U}_n = 0$. For such a sequence, we define the simplicial complex

$$TN(\mathcal{U}) = \bigcup_{n \in \mathbb{N}} N(\mathcal{U}_n \cup \mathcal{U}_{n+1}),$$

¹The results mentioned in this note were obtained in [Sa]. Then, for details, one can refer to the paper [Sa].

where we regard $\mathcal{U}_n \cap \mathcal{U}_m = \emptyset$ ($n \neq m$) as sets of vertices of $TN(\mathcal{U})$ even if $\mathcal{U}_n \cap \mathcal{U}_m \neq \emptyset$ as collections of open sets,² whence

$$N(\mathcal{U}_n \cup \mathcal{U}_{n+1}) \cap N(\mathcal{U}_{n+1} \cup \mathcal{U}_{n+2}) = N(\mathcal{U}_{n+1}).$$

Theorem 1. *A metric space $X = (X, d)$ is an ANR if and only if X has a zero-sequence $\mathcal{U} = (\mathcal{U}_n)_{n \in \mathbb{N}}$ of open covers with a map $f: |TN(\mathcal{U})| \rightarrow X$ satisfying the following conditions:*

- (i) $f(U) \in U$ for each $U \in TN(\mathcal{U})^{(0)} = \bigcup_{n \in \mathbb{N}} \mathcal{U}_n$, and
- (ii) $\lim_{n \rightarrow \infty} \text{mesh}\{f(\sigma) \mid \sigma \in N(\mathcal{U}_n \cup \mathcal{U}_{n+1})\} = 0$.

Under the above circumstances, if the image $f(|TN(\mathcal{U})|)$ is always contained in $Y \subset X$, then Y is homotopy dense in X .

This characterization of ANR's is due to Nguyen To Nhu [N] (cf. [NS]). By the alternative proof given in [Sa], the additional assertion was obtained. As a corollary, we have the following:

Corollary 1. *Let X be an ANR (resp. an AR) contained in a metric space M . Then, there exists a G_δ -set $Z \subset M$ such that Z is an ANR (resp. an AR) and X is homotopy dense in Z .*

We can also apply Theorem 1 to find conditions such that the metric completion of a metric space X is an ANR with X a homotopy dense subset. A subset D of a metric space X is said to be δ -dense in X if $\text{dist}(x, D) < \delta$ for every $x \in X$.

Corollary 2. *Let X be a metric space which has a zero-sequence $\mathcal{U} = (\mathcal{U}_n)_{n \in \mathbb{N}}$ of open covers with a map $f: |TN(\mathcal{U})| \rightarrow X$ satisfying the conditions (i) and (ii) of Theorem 1, where suppose $\mathcal{U}_n = \{B_X(x, \gamma_n) \mid x \in D_n\}$ for some δ_n -dense subset $D_n \subset X$ and $0 < \delta_n < \gamma_n$. Then, any metric space Z containing X isometrically as a dense subset is an ANR and X is homotopy dense in Z . In particular, the metric completion \tilde{X} of X is an ANR and X is homotopy dense in \tilde{X} .*

In the above, note that the γ_n -dense subset D_n of X may not be δ_n -dense in Z . For example, $D_n = \{i/n \mid 1 \leq i < n\}$ is $1/n$ -dense in $(0, 1)$ but it is not $1/n$ -dense in $[0, 1]$.

Now, we consider the following extension property:

- (e)_k There exist constants $\alpha > 0$ and $\beta > 1$ such that every map $f: |K^{(k)}| \rightarrow X$ of the k -skeleton of an arbitrary simplicial complex K with $\text{mesh}\{f(\sigma^{(k)}) \mid \sigma \in K\} < \alpha$ extends to a map $\tilde{f}: |K| \rightarrow X$ such that $\text{diam } \tilde{f}(\sigma) \leq \beta \text{diam } f(\sigma^{(k)})$ for each $\sigma \in K$.

²In [NS], we did not regard like this. Considering the set $\bigcup_{n \in \mathbb{N}} \mathcal{U}_n \times \{n\}$ as the set of vertices of $NT(\mathcal{U})$, this is reasonable.

The following corollary is motivated by the proof of AR property of the hyperspaces (cf. [vM, §5.3]).

Corollary 3. *Every LC^{k-1} metric space X with the property $(e)_k$ is an ANR. In particular, a metric space X with $(e)_0$ is an ANR (cf. Theorem 3).*

Remark. In Theorem 1, X is an AR when $\mathcal{U}_1 = \{X\}$. Every C^{k-1} and LC^{k-1} metric space X is an AR if it has the following:

$(\tilde{e})_k$ there exists a constant $\beta > 1$ such that every map $f: |K^{(k)}| \rightarrow X$ of the k -skeleton of an arbitrary simplicial complex K extends to a map $\tilde{f}: |K| \rightarrow X$ such that $\text{diam } \tilde{f}(\sigma) \leq \beta \text{diam } f(\sigma^{(k)})$ for each $\sigma \in K$.

2. UNIFORM ANR'S

In [Mi₂], E. Michael introduced uniform AR's and uniform ANR's, and studied them. Let $X = (X, d_X)$ and $Y = (Y, d_Y)$ be metric spaces and $A \subset X$. A map $f: X \rightarrow Y$ is said to be *uniformly continuous* at A if, for any $\varepsilon > 0$, there exists $\delta > 0$ such that if $a \in A$, $x \in X$ and $d_X(a, x) < \delta$ then $d_Y(f(a), f(x)) < \varepsilon$. A neighborhood U of A in X is called a *uniform neighborhood* if $\bigcup_{a \in A} B_X(a, \delta) \subset U$ for some $\delta > 0$. A metric space Y is called a *uniform ANR* if, for an arbitrary metric space X and a closed set $A \subset X$, every uniformly continuous map $f: A \rightarrow Y$ extends to a map $\tilde{f}: U \rightarrow Y$ from some uniform neighborhood U of A in X which is uniformly continuous at A . When f always extends over X (i.e., $U = X$), Y is a *uniform AR*. By virtue of [Mi₂, Theorem 1.2], a metric space Y is a uniform ANR (resp. a uniform AR) if and only if, for an arbitrary metric space Z which contains Y isometrically as a closed subset, there exists a retraction $r: U \rightarrow Y$ for some uniform neighborhood U in Y in Z (resp. $r: Z \rightarrow Y$) which is uniformly continuous at Y .³ The concept of uniform ANR's is useful since the metric completion of every uniform ANR is also a uniform ANR.

By using a zero-sequence of open covers in §1, we can prove the following version of Proposition 1.4 in [Mi₂]:

Theorem 2. *For an arbitrary metric space X , the following are equivalent:*

- (a) X is a uniform ANR;
- (b) Every metric space Z containing X isometrically as a dense subset is a uniform ANR and X is homotopy dense in Z ;
- (c) X is isometrically embedded in some uniform ANR Z as a homotopy dense subset.

³Such a retraction is called a *regular retraction* by H. Toruńczyk in [To₂].

Theorem 2 above means that a metric space X is a uniform ANR if and only if the metric completion of X is a uniform ANR with X a homotopy dense subset. However, in order that the metric completion of a metric ANR X is an ANR with X a homotopy dense subset, it is not necessary that X is a uniform ANR. In case X is totally bounded, X is a uniform ANR if and only if the metric completion of X is an ANR with X a homotopy dense subset.

A metric space Y is said to be *uniformly LC^k* if, for each $\varepsilon > 0$, there exists $\delta > 0$ such that any map $f: \mathbf{S}^i \rightarrow Y$ with $\text{diam } f(\mathbf{S}^i) < \delta$ extends to a map $\tilde{f}: \mathbf{B}^{i+1} \rightarrow Y$ with $\text{diam } \tilde{f}(\mathbf{B}^{i+1}) < \varepsilon$ for every $i \leq k$. In stead of “uniformly LC^0 ”, we also say “uniformly locally path-connected”. The subspace of \mathbb{R}^2 in the example above is not uniformly locally path-connected.

Theorem 3. *Every uniformly LC^{k-1} metric space Y with the property $(e)_k$ is a uniform ANR. In particular, a metric space X with $(e)_0$ is a uniform ANR.*

By Theorems 2 and 3, we have the following variation of Corollary 3 (cf. [SU, Lemma 2]):

Corollary 4. *Let X be a metric space and Y a dense subset of X . If Y is uniformly LC^{k-1} and has the property $(e)_k$, then X and Y are uniformly ANR's and Y is homotopy dense in X .*

Remark. In Theorem 3 and Corollary 4, by replacing the property $(e)_k$ with $(\tilde{e})_k$ and adding the condition that Y is C^{k-1} , “uniform ANR” can be “uniform AR”. In particular, a metric space X with $(\tilde{e})_0$ is a uniform AR.

3. DENSE (OR UNIFORM) LOCAL HYPER-CONNECTEDNESS

By using the notion of (local) hyper-connectedness, C.R. Borges [Bo] and R. Cauty [Ca] characterized AR's and ANR's, respectively. Here is introduced a little weaker notion. By Δ^{n-1} , we denote the standard $(n-1)$ -simplex in \mathbb{R}^n , that is,

$$\Delta^{n-1} = \{(t_1, \dots, t_n) \in \mathbb{R}^n \mid t_i \geq 0, \sum_{i=1}^{n+1} t_i = 1\}.$$

For an open cover \mathcal{U} of a space X and $Y \subset X$, we denote

$$Y^n(\mathcal{U}) = \{(y_1, \dots, y_n) \in Y^n \mid \exists U \in \mathcal{U} \text{ such that } \{y_1, \dots, y_n\} \subset U\}.$$

It is said that a space X is *densely locally hyper-connected* if X has an open cover \mathcal{W} , a dense subset D and functions $h_n: D^n(\mathcal{W}) \times \Delta^{n-1} \rightarrow X$, $n \in \mathbb{N}$, which satisfy the following conditions:

(i) if $t_i = 0$ then

$$\begin{aligned} h_n(y_1, \dots, y_n; t_1, \dots, t_n) \\ = h_{n-1}(y_1, \dots, y_{i-1}, y_{i+1}, \dots, y_n; t_1, \dots, t_{i-1}, t_{i+1}, \dots, t_n); \end{aligned}$$

- (ii) $\Delta^{n-1} \ni (t_1, \dots, t_n) \mapsto h_n(y_1, \dots, y_n; t_1, \dots, t_n) \in X$ is continuous for each $(y_1, \dots, y_n) \in D^n(\mathcal{W})$;
- (iii) every open cover \mathcal{U} of X has an open refinement \mathcal{V} such that $\mathcal{V} \prec \mathcal{W}$ (hence $D^n(\mathcal{V}) \subset D^n(\mathcal{W})$) and

$$\{h_n((D \cap V)^n \times \Delta^{n-1}) \mid V \in \mathcal{V}\} \prec \mathcal{U} \quad \text{for each } n \in \mathbb{N}.$$

It should be noticed that each h_n need not be continuous. If \mathcal{W} can be taken as $\mathcal{W} = \{X\}$ (i.e., $D^n(\mathcal{W}) = D^n$), we say that X is *densely hyper-connected*. In case $D = X$ (resp. $D = X$ and $\mathcal{W} = \{X\}$), X is *locally hyper-connected*⁴ (resp. *hyper-connected*). This concept is very similar to Michael's convex structure in [Mi₁]. In [Bo] and [Ca], AR's and ANR's are characterized by the hyper-connectedness and the local hyper-connectedness, respectively. A similar characterization was obtained by Himmelberg [Hi] (cf. Curtis [Cu]). These characterizations can be generalized in terms of the dense hyper-connectedness as follows:

Theorem 4. *A metrizable space X is an ANR if and only if X is densely locally hyper-connected. Moreover, X is an AR if and only if X is densely hyper-connected.*

Remark. In the definition of densely local hyper-connectedness, if the images of functions h_n are contained in Y , then Y is homotopy dense in X . In fact, if the images of functions h_n are contained in Y , then $f(|TN(\mathcal{U})|) \subset Y$, hence Y is homotopy dense in X by the additional statement of Theorem 1.

For a metric space X and $\eta > 0$, we denote

$$X^n(\eta) = \{(x_1, \dots, x_n) \in X^n \mid \text{diam}\{x_1, \dots, x_n\} < \eta\}.$$

A metric space X is said to be *uniformly locally hyper-connected* if there are $\eta > 0$ and functions $h_n: X^n(\eta) \times \Delta^{n-1} \rightarrow X$, $n \in \mathbb{N}$, which satisfy the same conditions as (i) and (ii) above, and the following (iii') instead of (iii):

- (iii') for each $\varepsilon > 0$, there is $0 < \delta < \varepsilon$ such that

$$\text{diam } h_n(\{x\} \times \Delta^{n-1}) < \varepsilon \quad \text{for every } n \in \mathbb{N} \text{ and } x \in X^n(\delta).$$

When every h_n is defined on the whole space $X^n \times \Delta^{n-1}$, it is said that X is *uniformly hyper-connected*.

Now, we give a characterization of uniform ANR's and uniform AR's.

⁴The local hyper-connectedness is in the sense of [Ca] but not in the sense of [Bo].

Theorem 5. *A metric space $X = (X, d)$ is a uniform ANR if and only if X is uniformly locally hyper-connected. Moreover, X is a uniform AR if and only if X is uniformly hyper-connected.*

The following is a combination of Theorems 2 and 5:

Corollary 5. *Let X be a uniformly (locally) hyper-connected metric space and Z a metric space which contains X isometrically as a dense subset. Then, X and Z are uniform AR's (uniform ANR's) and X is homotopy dense in Z . In particular, the metric completion \tilde{X} of X is a uniform AR (uniform ANR) and X is homotopy dense in \tilde{X} .*

REFERENCES

- [AE] R. Arens and J. Eells, *On embedding uniform and topological spaces*, Pacific J. Math. **6** (1956), 397–403.
- [Bo] C.R. Borges, *A study of absolute extensor spaces*, Pacific J. Math. **31** (1969), 609–617; *A correction and an answer*, *ibid.* **50** (1974), 29–30.
- [Ca] R. Cauty, *Rétraction dans les espaces stratifiables*, Bull. Soc. Math. France **102** (1974), 129–149.
- [Cu] D.W. Curtis, *Some theorem and examples on local equiconnectedness and its specializations*, Fund. Math. **72** (1971), 101–113.
- [Hi] C.J. Himmelberg, *Some theorems on equiconnected and locally equiconnected spaces*, Trans. Amer. Math. Soc. **115** (1965), 43–53.
- [Mi₁] E.A. Michael, *Convex structures and continuous selections*, Canad. J. Math. **11** (1959), 556–575.
- [Mi₂] ———, *Uniform AR's and ANR's*, Compositio Math. **39** (1979), 129–139.
- [vM] J. van Mill, *Infinite-Dimensional Topology, Prerequisites and Introduction*, North-Holland Math. Library **43**, Elsevier Sci. Publ. B.V., Amsterdam, 1989.
- [N] Nguyen To Nhu, *Investigating the ANR-property of metric spaces*, Fund. Math. **124** (1984), 243–254; *Corrections*, *ibid.* **141** (1992), 297.
- [NS] ——— and K. Sakai, *The compact neighborhood extension property and local equi-connectedness*, Proc. Amer. Math. Soc. **121** (1994), 259–265.
- [Sa] K. Sakai, *The completions of metric ANR's and homotopy dense subsets*, J. Math. Soc. Japan (to appear).
- [SU] ——— and S. Uehara, *A Hilbert cube compactification of the Banach space of continuous functions*, Topology Appl. **92** (1999), 107–118.
- [To₁] H. Toruńczyk, *A short proof of Hausdorff's theorem on extending metrics*, Fund. Math. **77** (1972), 191–193.
- [To₂] ———, *Absolute retracts as factors of normed linear spaces*, Fund. Math. **86** (1974), 53–67.
- [To₃] ———, *Concerning locally homotopy negligible sets and characterization of ℓ_2 -manifolds*, Fund. Math. **101** (1978), 93–110.