

GL_l のスーパーカスピダル表現の指標公式 (CHARACTER FORMULA FOR THE SUPERCUSPIDAL REPRESENTATIONS OF GL_l)

大阪府立大総合科学部 高橋 哲也 (TETSUYA TAKAHASHI)

Introduction

Let l be a prime, A a central simple algebra of dimension l^2 over a non-archimedean local field F and E/F an extension of degree l in A . As is well-known ([15], [5]), any irreducible supercuspidal representation of A^\times is obtained from a quasi-character of E^\times . The aim of this paper is to get a character formula for the irreducible supercuspidal representations of A^\times on the set of elliptic regular elements. To calculate the character on the split torus is another problem. See Murnaghan's papers ([16], [17] and [18]) for this topic. We only remark that the character value on the elliptic regular conjugacy classes determines uniquely the supercuspidal representation.

When E/F is unramified, a character formula was obtained in [20]. Therefore we treat the case E/F is ramified. When the residual characteristic p of F equals to l , the ramification is wild. This case is very hard to treat (see e.g. [21]). Thus we assume $p \neq l$. Since the case $l = 2$ is already solved in [13], we also assume l is odd. We note that A is isomorphic to a division algebra $D = D_l$ of dimension l^2 over F or the algebra $M_l(F)$ of $l \times l$ matrices over F .

Let D_n be a division algebra of dimension n^2 over F . Deligne-Kazhdan-Vignéras [8] and Rogawski [19] proved an abstract matching theorem: there is a bijection between the set of equivalence classes of irreducible representations of D_n^\times and that of essentially square-integrable representations of $GL_n(F)$ which preserves the characters up to $(-1)^{n-1}$. In the tame case, i.e., when n is prime to the residual characteristic of F , Moy [15] proved that there is a bijection between the same sets as above using the concrete construction of the representations given by Howe [12]. Henniart ([10]) has shown that two correspondences coincide when $n = l \neq p$. Thus we only treat the GL_l case. Our main result is Theorem 3.12. As in the unramified case, the analogue of Weyl's character formula holds for our character formula. This does not hold when $l = p$ (cf. [22]).

Section 1 is devoted to the review of the construction of an irreducible supercuspidal representation π_θ (resp. π'_θ) of $GL_l(F)$ (resp. D_l^\times) from a generic quasi-character θ of E^\times and the known results about the representation. We note that π_θ is not always monomial, i.e., induced from a one-dimensional representation, but it can be written as a \mathbb{Q} -linear combination of monomial representations. In fact π_θ is written as \mathbb{Q} -linear combination of the forms $\text{ind}_H^{GL_l(F)} \rho_\theta$ where H is a compact mod center subgroup of $GL_l(F)$ and ρ_θ is quasi-characters of H .

In section 2, we compute the character of π_θ up to some root numbers. Let $G = GL_l(F)$, B the normalizer of an Iwahori subgroup of G containing H and $\eta_\theta = \text{ind}_H^B \rho_\theta$. Since we treat only elliptic regular conjugacy classes, we consider the character χ_{π_θ} on L^\times where L/F are extensions of fields of degree l . Moreover the case $L = E$ is essential. By the

Frobenius formula and the result of Kutzko ([14]), we have only to calculate the sum

$$\chi_{\eta_\theta}(x) = \sum_{a \in H \setminus B} \rho_\theta(axa^{-1})$$

for $x \in E$ in order to get the character formula of π_θ . Therefore it is essential to know when $axa^{-1} \in H$, which is determined in Lemma 2.1. From this, we get the character formula of η_θ except near the conductor (Proposition 2.2). But this formula contains the Gauss sum part $G(y, j)$, which is calculated later. The exceptional part can be calculated directly by taking the explicit matrix form of E^\times (Lemma 2.4). Except this lemma, there is no new result in this section. But since the proofs are short and simple and we use the property “intertwining implies conjugacy” of E/F -minimal element (very cuspidal in the terminology of Carayol [4]) as the key tool, the result may be extended to GL_n , at least when n is prime to p . Section 3 is devoted to the calculation of the Gauss sum part $G(y, j)$. It appears in the character formula on E^\times . For this purpose, it is the point that we have only to treat the character of π_θ on $U_1^* = F^\times(1 + P_E) - F^\times(1 + P_E^2)$. For this calculation, we use the E^\times -structure of various objects. We first assume E/F is a Galois extension since E^\times -module structure can be described easily for this case. This part is analogous to section 1 of [20], but everything becomes easier since we have only to treat U_1^* . When E/F is non-Galois, we use the base change lift. Let L/F be an unramified extension of degree $l - 1$. In L , there exists a l -th root of unity and EL/L is Galois. Therefore we can use the tools in Galois case for $GL_l(L)$. Let $\text{Gal}(L/F) = \langle \tau \rangle$. By the result of Bushnell-Henniart [3], there is a base change lift η_L of η_θ to H_L^1 such that the twisted trace of η_L by τ gives the trace of η_θ . (See Proposition 3.7 and Lemma 3.8). We remark that we need not assume the characteristic of F is 0 since we do not use the Arthur-Clozel base change lift [1]. The method to calculate the twisted trace of η_L is similar to that of Galois case. The complete character formula is stated as Theorem 3.12.

Closing this introduction, we compare our formula with the known results. The same type of character formula for the division algebra case was given by Corwin, Moy and Sally, Jřin [6] and for GL_l case by Debacker in [7]. Their formulas agree with the result given in section 2. It contains some root numbers associated with a quadratic form. In this paper, we have determined it completely in section 3. Moreover we find the Kloosterman sum appears in the character formula. These are new results of this paper. In [22], the author gave the character formula of π_θ for GL_3 by using the decomposition of π_θ as E^\times -module. But this need the explicit matrix form of an inverse matrix which is hard to treat for large l . We can simplify the proof of the main theorem, although we treat a general prime l .

Notation

Let F be a non-archimedean local field. We denote by \mathcal{O}_F , P_F , ϖ_F , k_F and v_F the maximal order of F , the maximal ideal of \mathcal{O}_F , a prime element of P_F , the residue field of F and the valuation of F normalized by $v_F(\varpi_F) = 1$. We set q to be the number of elements in k_F . Hereafter we fix an additive character ψ of F whose conductor is P_F , i.e., ψ is trivial on P_F and not trivial on \mathcal{O}_F . For an extension E over F , we denote by tr_E , n_E the trace and norm to F respectively. We set $\psi_E = \psi \circ \text{tr}_E$. The trace of matrix is denoted by Tr . For an irreducible admissible representation π of $GL_l(F)$, the conductor exponent of π is defined to be the integer $f(\pi)$ such that the local constant $\varepsilon(s, \pi, \psi)$ of Godement-Jacquet [9] is the form $aq^{-s(f(\pi)-l)}$.

We call π *minimal* if

$$f(\pi) = \min_{\eta} f(\pi \otimes (\eta \circ \text{Nr})),$$

where η runs through the quasi-characters of F^\times . Let G be a totally disconnected, locally compact group. We denote by \widehat{G} the set of (equivalence classes of) irreducible admissible representations of G . For a closed subgroup H of G and a representation ρ of H , we denote by $\text{Ind}_H^G \rho$ (resp. $\text{ind}_H^G \rho$) the induced representation (resp. compactly induced representation) of ρ to G . For a representation π of G , we denote by $\pi|_H$ the restriction of π to H .

1. CONSTRUCTION OF THE REPRESENTATION

Let E be a ramified extension of F of degree l . Then E can be embedded into $M_l(F)$ and, up to conjugacy, the embedding is unique. Let $G = \text{GL}_l(F)$. In this section, we review the construction of supercuspidal representations of G which are parameterized by the quasi-characters of E^\times . Of course, this construction is well-known ([4], [15]).

Definition 1.1. Let θ be a quasi-character of E^\times and $f(\theta) = \min\{n \mid \text{Ker } \theta \subset 1 + P_E^n\}$. Then θ is called *generic* if $f(\theta) \not\equiv 1 \pmod{l}$. For a generic character θ of E^\times , $\gamma_\theta \in P_E^{1-f(\theta)} - P_E^{2-f(\theta)}$ is defined by

$$(1.1) \quad \theta(1+x) = \psi_E(\gamma_\theta x) \quad \text{for } x \in P_E^{[(f(\theta)+1)/2]}.$$

Then $F(\gamma_\theta) = E$. We denote by \widehat{E}_{gen}^\times the set of generic quasi-characters of E^\times .

We construct an irreducible supercuspidal representation of $G = \text{GL}_l(F)$ from $\theta \in \widehat{E}_{gen}^\times$. For simplicity, we set $\gamma = \gamma_\theta$. Since E/F is tamely ramified, there exists a prime element ϖ_E of \mathcal{O}_E satisfying $\varpi_E^l \in F$. Put $\varpi_F = \varpi_E^l$. We identify $M_l(F)$ with $\text{End}_F E$ and G with $\text{Aut}_F E$ by the F -basis $\{\varpi_E^{l-1}, \varpi_E^{l-2}, \dots, \varpi_E, 1\}$ of E , which is also an \mathcal{O}_F -basis of \mathcal{O}_E . By the lattice flag $\{P_E^i\}_{i \in \mathbb{Z}}$, we construct a maximal compact modulo center subgroup. The construction of the representation is well-known. For details, see [15].

Definition 1.2. For $i \in \mathbb{Z}$, set

$$A^i = \{f \in M_l(F) \mid f(P_E^j) \subset P_E^{j+i} \text{ for all } j \in \mathbb{Z}\}.$$

Put $K = (A^0)^\times$, $B = E^\times K$ and $K^i = 1 + A^i$ for $i \geq 1$.

Then K is an Iwahori subgroup of G and B is a normalizer of K . At first we construct an irreducible representation of B from a generic quasi-character of E^\times .

Let θ be a generic quasi-character of E^\times , i.e., $f(\theta) = n \not\equiv 1 \pmod{l}$. There exists an element $\gamma \in P_E^{1-n}$ such that $\theta(1+x) = \psi_E(\gamma x)$ for $x \in P_E^m$ where $m = [(n+1)/2]$. Define ψ_γ on K^m by $\psi_\gamma(1+x) = \psi(\text{Tr}(\gamma x))$ for $x \in A^m$. Then ψ_γ is a quasi-character of K^m . Put $H = E^\times K^m$ and define a quasi-character ρ_θ of H by

$$(1.2) \quad \rho_\theta(h \cdot g) = \theta(h) \psi_\gamma(g) \quad \text{for } h \in E^\times, \quad g \in K^m.$$

Let J be the normalizer of ψ_γ in B , i.e.,

$$J = \{a \in B \mid \psi_\gamma^a = \psi_\gamma\},$$

where $\psi_\gamma^a(x) = \psi_\gamma(a^{-1}xa)$ for $x \in K^m$. Then $J = E^\times K^{m'}$ where $m' = [n/2]$. Put $\eta_\theta = \text{Ind}_H^B \rho_\theta$.

When n is even, i.e., $n = 2m$, then $J = H = E^\times K^m$. By the Clifford theory, η_θ is an irreducible representation of B . We put

$$(1.3) \quad \kappa_\theta = \eta_\theta.$$

When n is odd, i.e., $n = 2m - 1$, then $J = E^\times K^{m-1}$. Thus η_θ is not irreducible. In this case, we put

$$(1.4) \quad \kappa_\theta = \frac{1 - \left(\frac{q}{l}\right) q^{(l-1)/2}}{l q^{(l-1)/2}} \sum_{\chi \in (E^\times / F^\times (1 + P_E))^\wedge} \eta_{\theta \otimes \chi} + \left(\frac{q}{l}\right) \eta_\theta,$$

where $\left(\frac{q}{l}\right)$ is the Legendre symbol. The following result is well-known (see [15]).

Theorem 1.3. *Let the notation be as above. Then κ_θ is an irreducible representation of B . Put $\pi_\theta = \text{ind}_B^G \kappa_\theta$. Then π_θ is an irreducible supercuspidal representation of G such that*

1. *the L -function of π_θ is 1;*
2. *$\varepsilon(\pi_\theta, \psi) = \varepsilon(\theta, \psi_E)$; in particular $f(\pi_\theta) = f(\theta) + l$;*
3. *$\bigcup_E \{\pi_\theta \mid \theta \in \widehat{E}_{gen}^\times\} = \{\pi \in A_0(G) \mid f_{\min}(\pi) \not\equiv 0 \pmod{l}\}$, where E runs through isomorphism classes of ramified extensions of degree l over F and $A_0(G)$ be the set of equivalent classes of the supercuspidal representations of G .*

Remark. If $\pi \in A_0(G)$ and $f_{\min}(\pi) \equiv 0 \pmod{l}$, π can be constructed from a regular quasi-characters of L^\times , where L is an unramified extension of F of degree l . The character formula for such a representation was given in [20].

Next we construct an irreducible representation of $D^\times = D_l^\times$ from $\theta \in E_{gen}^\times$. Let $f(\theta) = n$. We recall $n \not\equiv 1 \pmod{l}$. We define a function ψ_γ on $1 + P_D^m$ by $\psi_\gamma(1 + x) = \psi(\text{Tr}(\gamma x))$ for $x \in P_D^m$. Then ψ_γ is a quasi-character of $1 + P_D^m$. $H' = E^\times (1 + P_D^m) \subset D^\times$ and define a quasi-character ρ'_θ of H' by

$$(1.5) \quad \rho'_\theta(h \cdot g) = \theta(h) \psi_\gamma(g) \quad \text{for } h \in E^\times, \quad g \in 1 + P_D^m.$$

When n is even, i.e., $n = 2m$, we set

$$(1.6) \quad \pi'_\theta = \text{Ind}_{H'}^{D^\times} \rho'_\theta.$$

When n is odd, i.e., $n = 2m - 1$, we set

$$(1.7) \quad \pi'_\theta = \frac{1 - \left(\frac{q}{l}\right) q^{(l-1)/2}}{l q^{(l-1)/2}} \sum_{\chi \in (E^\times / F^\times (1 + P_E))^\wedge} \text{Ind}_{H'}^{D^\times} \rho'_{\theta \otimes \chi} + \left(\frac{q}{l}\right) \text{Ind}_{H'}^{D^\times} \rho'_\theta,$$

where $\left(\frac{q}{l}\right)$ is the Legendre symbol. The following result is essentially well-known. (See [2], [15]).

Theorem 1.4. *Let the notation be as above. Then π'_θ is an irreducible minimal representation of D^\times such that*

1. *the degree of π'_θ is $q^{(n-2)(l-1)/2} \frac{(q^l-1)}{q-1}$;*
2. *$\varepsilon(\pi'_\theta, \psi) = \varepsilon(\theta, \psi_E)$; in particular $f(\pi'_\theta) = f(\theta) + l$;*

3. $\bigcup_E \{\pi'_\theta | \theta \in \widehat{E}_{reg}^\times\} = \{\pi' \in \widehat{D}^\times | f_{\min}(\pi') \not\equiv 0 \pmod{l}\}$, where E runs through the isomorphism classes of ramified extensions of degree l over F .
4. The correspondence $\pi'_\theta \leftrightarrow \pi_\theta$ by way of generic quasi-characters of E^\times is a bijection and preserves ε -factors and conductoral exponents. (This correspondence is a special case of Howe's bijection (see [15]).)

On the other hand, there exists an abstract matching theorem, which is called the Deligne-Kazhdan correspondence ([8], [19]).

Theorem 1.5. *There is a bijection between the set of irreducible representations of D^\times and the set of essentially square-integrable representations of G which preserves the characters on elliptic regular elements. In particular, it preserves ε -factors and conductoral exponents.*

By the result of Henniart ([10] Theorem 8.1), these two correspondences coincide.

Theorem 1.6. *If $l \neq p$ is a prime, Howe's bijection (1.4) coincides with Deligne-Kazhdan correspondence (1.5) between the set of essentially square-integrable representations of GL_l and the set of irreducible representations of D_l^\times .*

At the end of this section, we quote the result of Kutzko ([14]) in the form that the character formula of π_θ on elliptic regular elements is essentially given by the one of κ_θ .

Theorem 1.7. *Let x be an elliptic regular element of G .*

1. If $F(x)/F$ is ramified and $x \notin F^\times(1 + P_{F(x)}^n)$,

$$\chi_{\pi_\theta}(x) = \chi_{\kappa_\theta}(x).$$

2. If $F(x)/F$ is unramified and $x \notin F^\times(1 + P_{F(x)}^{[n/l]+1})$,

$$\chi_{\pi_\theta}(x) = 0.$$

Proof. These are obtained by applying Proposition 5.5 in [14] to our case. □

Remark. Since

$$(1.8) \quad \chi_{\kappa_\theta}(x) = \begin{cases} \left(\frac{q}{l}\right) \chi_{\eta_\theta}(x) & x \in E^\times - F^\times(1 + P_E), \\ \frac{1}{q^{(l-1)/2}} \chi_{\eta_\theta}(x) & x \in F^\times(1 + P_E), \end{cases}$$

we have only to calculate χ_{η_θ} .

2. CALCULATION OF THE CHARACTER

Now we begin to calculate the characters of the representations constructed in the previous section. In this section, we shall get a character formula up to some root numbers. These root numbers are calculated explicitly in the next section.

Hereafter we fix a generic character θ and put $\rho = \rho_\theta$, $\eta = \eta_\theta$ and so on. Since E/F is a totally tamely ramified extension, there exists a prime element ϖ_E of \mathcal{O}_E such that $\varpi_E^l \in P_F - P_F^2$. Put $\varpi_E^l = \varpi_F$. As in the previous section, we identify $M_l(F)$ with $\text{End}_F(E)$ by the F -basis $\{\varpi_E^{l-1}, \varpi_E^{l-2}, \dots, \varpi_E, 1\}$, which is an \mathcal{O}_F -basis of \mathcal{O}_E . Thus we get the explicit matrix forms of various objects:

$$(2.1) \quad \varpi_E = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \\ \varpi_F & 0 & \cdots & \cdots & 0 \end{pmatrix},$$

$$(2.2) \quad K = \left\{ \begin{pmatrix} a_{11} & \cdots & a_{1l} \\ \cdots & \cdots & \cdots \\ a_{l1} & \cdots & a_{ll} \end{pmatrix} \middle| \begin{array}{l} a_{ij} \in \mathcal{O}_F \quad \text{if } i < j \\ a_{ii} \in \mathcal{O}_F^\times \\ a_{ij} \in P_F \quad \text{if } i > j \end{array} \right\},$$

$$(2.3) \quad A^0 = \left\{ \begin{pmatrix} a_{11} & \cdots & a_{1l} \\ \cdots & \cdots & \cdots \\ a_{l1} & \cdots & a_{ll} \end{pmatrix} \middle| \begin{array}{l} a_{ij} \in \mathcal{O}_F \quad \text{if } i \leq j \\ a_{ij} \in P_F \quad \text{if } i > j \end{array} \right\},$$

$$(2.4) \quad A^1 = \left\{ \begin{pmatrix} a_{11} & \cdots & a_{1l} \\ \cdots & \cdots & \cdots \\ a_{l1} & \cdots & a_{ll} \end{pmatrix} \middle| \begin{array}{l} a_{ij} \in \mathcal{O}_F \quad \text{if } i < j \\ a_{ij} \in P_F \quad \text{if } i \geq j \end{array} \right\}.$$

If $q \equiv 1 \pmod{l}$, F has a primitive l -th root of unity ζ and E/F is a Galois extension. Let σ be a generator of $\text{Gal}(E/F)$ determined by $\sigma\varpi_E = \varpi_E\zeta$. We denote the diagonal matrix $\text{diag}(1, \zeta^{l-1}, \zeta^{l-2}, \dots, \zeta)$ by ξ . Then ξ satisfies $\xi^l = 1$ and $\xi x \xi^{-1} = \sigma x$ for $x \in E$.

Define a natural ring morphism R from A^0 to k_F^l by the identification of A^0/A^1 with k_F^l . We note that if $R(a) = (\alpha_0, \alpha_1, \dots, \alpha_{l-1})$, $R(\varpi_E a \varpi_E^{-1}) = (\alpha_1, \alpha_2, \dots, \alpha_0)$. For convenience, we extend the suffix to \mathbb{Z} by putting $\alpha_i = \alpha_{i \bmod l}$. The next lemma is the key tool for the character calculation.

Lemma 2.1. *Let $x \in P_E^i - (F + P_E^{i+1})$, $g \in B$ and j a positive integer. If $g x g^{-1} \in E^\times(1 + A^j)$, then*

$$g \in \begin{cases} E^\times(1 + A^j) & \text{if } q \not\equiv 1 \pmod{l}, \\ \bigcup_{k=0}^{l-1} E^\times(1 + A^j)\xi^k & \text{if } q \equiv 1 \pmod{l}. \end{cases}$$

Proof. We may assume $g \in A_0$ by replacing g by $\varpi_E^{-k}g$ if $g \in A_k$. Let $x = \varpi_E^i x_0$ for $x_0 \in \mathcal{O}_E^\times$ and $R(g) = (\alpha_0, \alpha_1, \dots, \alpha_{l-1})$. Then

$$R(g x g^{-1} x^{-1}) = (\alpha_0 \alpha_i^{-1}, \alpha_1 \alpha_{i+1}^{-1}, \dots, \alpha_{l-1} \alpha_{l-1+i}^{-1}),$$

where $\alpha_s = \alpha_{s \bmod l}$ for $s \in \mathbb{Z}$. Since $x \notin F + P_E^{i+1}$, $i \not\equiv 0 \pmod{l}$. Therefore $g x g^{-1} x^{-1} \in E^\times$ implies

$$\begin{cases} \alpha_0 = \alpha_1 = \cdots = \alpha_{l-1} & \text{if } q \not\equiv 1 \pmod{l}, \\ \alpha_k = \zeta^j \alpha_0 \quad (0 \leq k \leq l-1) & \text{otherwise,} \end{cases}$$

for some integer j . Since $\xi \varpi_E \xi^{-1} = \zeta \varpi_E$, we get:

$$g \in \begin{cases} E^\times(1 + A^1) & \text{if } q \equiv 1 \pmod{l}, \\ \bigcup_{k=0}^{l-1} E^\times(1 + A^1)\xi^k & \text{otherwise.} \end{cases}$$

Thus we may assume $g - 1 \in A^k - (P_E^{k+1} + A^{k+1})$ for $k \geq 1$. Put $g - 1 = \varpi_E^k g_0$ and $R(g_0) = (\beta_0, \beta_1, \dots, \beta_{l-1})$. Since

$$\begin{aligned} g x g^{-1} x^{-1} &\equiv 1 + (g - 1) - x(g - 1)x^{-1} \pmod{A^{k+1}} \\ &\equiv 1 + \varpi_E^k (g_0 - x g_0 x^{-1}) \pmod{A^{k+1}}, \end{aligned}$$

$R(g_0 - x g_0 x^{-1}) = (\beta_0 - \beta_k, \beta_1 - \beta_{1+k}, \dots, \beta_{l-1} - \beta_{l-1+k})$. Therefore $g x g^{-1} x^{-1} \in E^\times K^{k+1}$ contradicts $g - 1 \in A^k - (P_E^{k+1} + A^{k+1})$. It implies that if $g x g^{-1} x^{-1} \in E^\times K^j$,

$$g \in \begin{cases} E^\times(1 + A^j) & \text{if } q \not\equiv 1 \pmod{l}, \\ \bigcup_{k=0}^{l-1} E^\times(1 + A^j)\xi^k & \text{if } q \equiv 1 \pmod{l}. \end{cases}$$

□

Put $U_{-1} = E^\times, U_0 = F^\times \mathcal{O}_E^\times, U_i = F^\times(1 + P_E^i)$ for $i \geq 1$ and $U_i^* = U_i - U_{i+1}$ for $j \geq -1$. The previous lemma gives the character of η_θ on $E^\times - U_{n-1}$. We remark $\text{Aut}_F E = \{1\}$ if $q \not\equiv 1 \pmod{l}$.

Proposition 2.2. *Let $x \in U_i^*$ for $-1 \leq i < n - 1$. If $i \not\equiv 0 \pmod{l}$, x is written in the form $x = c(1 + y)$ for $c \in F$ and $y = \varpi_E^i y_0 \in \varpi_E^i \mathcal{O}_E^\times$. For $u \in k_F^\times$ and $j \in (\mathbb{Z}/l\mathbb{Z})^\times$, we define the Gauss sum part $G(u, j)$ by*

$$(2.5) \quad G(u, j) = \sum_{(\alpha_0, \dots, \alpha_{l-1}) \in k_F^l / \Delta} \psi \left(\sum_{k=0}^{l-1} u(\alpha_{k+1} - \alpha_k) \alpha_{j+k} \right),$$

where $\Delta = \{(\alpha, \dots, \alpha) \mid \alpha \in k_F\}$ is the image of the diagonal embedding of k_F into k_F^l . Then χ_{η_θ} on U_i^* is given as follows:

$$\chi_{\eta_\theta}(x) = \begin{cases} \sum_{\sigma \in \text{Aut}_F E} \theta(\sigma x) & i = -1, \\ q^{\lfloor (i+1)/2 \rfloor (l-1)} \sum_{\sigma \in \text{Aut}_F E} \theta(\sigma x) & i > 0 \text{ and } n - i \text{ even}, \\ q^{\lfloor i/2 \rfloor (l-1)} \sum_{\sigma \in \text{Aut}_F E} \theta(\sigma x) & \\ G(\gamma \varpi_E^{n-1} y_0 (\sigma \varpi_E / \varpi_E)^i, c) & i > 0 \text{ and } n - i \text{ odd}, \end{cases}$$

where $c = i^{-1}(n + i - 1)/2 \in (\mathbb{Z}/l\mathbb{Z})^\times$.

Proof. Put ${}^a x = axa^{-1}$ for $a, x \in G$. At first we treat the case $x \in U_{-1}^* = E^\times - F^\times \mathcal{O}_E^\times$. Since

$$\chi_{\eta_\theta}(x) = \sum_{a \in H \setminus B} \rho_\theta({}^a x),$$

we have only to show that if ${}^a x \in H$ for $a \in B$, then

$$a \in \begin{cases} H & \text{if } q \not\equiv 1 \pmod{l}, \\ \bigcup_{k=0}^{l-1} H \xi^k & \text{if } q \equiv 1 \pmod{l}. \end{cases}$$

This follows immediately from Lemma 2.1.

Now we treat the case $x = c(1+y)$ for $c \in F$ and $y \in P_E^i - (F + P_E^{i+1})$. We may assume $c = 1$ since F^\times is the center of B . For $1+k \in K^{[(n-i+1)/2]}$ and $a \in B$, we have

$$\begin{aligned}\chi_{\eta_\theta}(1+y) &= \sum_{a \in H \setminus B} \rho_\theta({}^a(1+y)) \\ &= C \sum_{1+k \in K^{n-i} \setminus K^{[(n-i+1)/2]}} \sum_{a \in H \setminus B} \rho_\theta({}^{a(1+k)}(1+y)),\end{aligned}$$

where $C = \frac{1}{K^{n-i} \setminus K^{[(n-i+1)/2]}}$. In the above expression,

$$\begin{aligned}\rho_\theta({}^{a(1+k)}(1+y)) &= \rho_\theta(1 + {}^a y + {}^a(ky - yk)) \\ &= \rho_\theta(1 + {}^a y) \rho_\theta(1 + {}^a((1+y)^{-1}(ky - yk))) \\ &= \rho_\theta(1 + {}^a y) \psi(\text{Tr } \gamma^a((1+y)^{-1}(ky - yk))) \\ &= \rho_\theta(1 + {}^a y) \psi(\text{Tr } a^{-1} \gamma(1+y)^{-1}(ky - yk)) \\ &= \rho_\theta(1 + {}^a y) \psi(\text{Tr}(y^{a^{-1}} \gamma - a^{-1} \gamma y)(1+y)^{-1}k)\end{aligned}$$

since $yk^2 \in A^n$ and $a(1+y)^{-1}(ky - yk)a^{-1} \in A^m$. If $y^{a^{-1}} \gamma - a^{-1} \gamma y \notin A^{1-[(n-i+1)/2]}$, the map $k \mapsto \psi(\text{Tr}(y^{a^{-1}} \gamma - a^{-1} \gamma y)(1+y)^{-1}k)$ is a non-trivial character of $A^{n-i} \setminus A^{[(n-i+1)/2]}$; thus

$$\sum_{k \in A^{n-i} \setminus A^{[(n-i+1)/2]}} \psi(\text{Tr}(y^{a^{-1}} \gamma - a^{-1} \gamma y)(1+y)^{-1}k) = 0.$$

By Lemma 3.3 in [4], $y^{a^{-1}} \gamma - a^{-1} \gamma y \in A^{1-[(n-i+1)/2]}$ is equivalent to $a^{-1} \gamma \in E^\times K^{n-i-[(n-i+1)/2]}$. Thus it follows from Lemma 2.1 that

$$\chi_{\eta_\theta}(1+y) = \sum_{\sigma \in \text{Aut}_F E} \sum_{1+a \in H \setminus E^\times K^{[(n-i)/2]}} \rho_\theta(1 + (1+a)^\sigma y(1+a)^{-1}).$$

By virtue of $(1+y)^{-1}(1 + (1+a)y) \in K^m$ and $(1+y)^{-1}(1 + (1+a)y) \equiv 1 + (1+y)^{-1}((ay - ya) + (ya - ay)a) \pmod{K^n}$,

$$\rho_\theta(1 + (1+a)y) = \theta(1+y) \psi_\gamma((1+y)^{-1}(ay - ya)) \psi_\gamma((1+y)^{-1}(ya - ay)a).$$

Since

$$\begin{aligned}\psi_\gamma((1+y)^{-1}(ay - ya)) &= \psi(\text{Tr}(y\gamma(1+y)^{-1} - \gamma(1+y)^{-1}y)a) = 1, \\ \psi_\gamma((1+y)^{-1}(ya - ay)a) &= \psi_\gamma((ya - ay)a) \text{ and } |E^\times K^j / E^\times K^m| = q^{(l-1)(m-j)},\end{aligned}$$

we obtain

$$\chi_{\eta_\theta}(1+y) = \begin{cases} q^{m-(n-i)/2} \sum_{\sigma \in \text{Aut}_F E} \theta(1 + {}^\sigma y) & n-i \text{ even,} \\ q^{m-(n-i+1)/2} \sum_{\sigma \in \text{Aut}_F E} \theta(1 + {}^\sigma y) S(n-i, \sigma) & n-i \text{ odd,} \end{cases}$$

where

$$S(n-i, \sigma) = \sum_{a \in A^{(n-i+1)/2} + E \cap A^{(n-i-1)/2} \setminus A^{(n-i-1)/2}} \psi_\gamma(({}^\sigma ya - a^\sigma y)a).$$

Now we may assume $n-i$ odd and $\sigma = 1$. Put $y = \varpi_E^i y_0$, $a = \varpi_E^{(n-i-1)/2} a_0$ and $S = S(n-i, 1)$. Since

$$\begin{aligned}(ya - ay)a &= \varpi_E^{n-1} (y_0 \varpi_E^{-(n-i-1)/2} a_0 \varpi_E^{(n-i-1)/2} \\ &\quad - \varpi_E^{-(n+i-1)/2} a_0 \varpi_E^{(n+i-1)/2} y_0) a_0,\end{aligned}$$

we have by way of the map $R : A_0/A_1 \rightarrow k_F^l$ that

$$S = \sum_{(\alpha_j) \in k_F^l/\Delta} \psi \left(\sum_{j=0}^{l-1} \gamma \varpi_E^{n-1} y_0 (\alpha_{j-(n-i-1)/2} - \alpha_{j-(n+i-1)/2}) \alpha_j \right).$$

(The suffix is extended to \mathbb{Z} by $\alpha_j = \alpha_{j \bmod l}$.) At first replacing the suffix j by $j + (n + i - 1)/2$ and then replacing α_{ij} by α_j , we get our lemma. \square

Remark. In [6] and [7], it is proved that the Gauss sum $G(u, j)$ is a fourth root of unity when $l \neq 2$.

Next we calculate the character on $K^{n-1} - K^n$. We state the character formula including the case $x \notin E$. On $K^{n-1} - K^n$, the Kloosterman sum appears in the formula.

Definition 2.3. For $a \in k_F^\times$, we define the Kloosterman sum $\text{Kl}(a)$ by

$$(2.6) \quad \text{Kl}(a) = \sum_{\substack{(y_0, \dots, y_{l-1}) \in k_F^l \\ y_0 \cdots y_{l-1} = a}} \psi(y_0 + \cdots + y_{l-1}).$$

Lemma 2.4. Let $x = 1 + \varpi_E^{n-1} x_0$ for $x_0 = \text{diag}(k_0, \dots, k_{l-1})$, ($k_i \in \mathcal{O}_F^\times$). Then

$$\chi_{\eta_\theta}(x) = q^{(l-1)(m-1)} \text{Kl} \left((\gamma \varpi_E^{n-1})^l \prod_{j=0}^{l-1} k_j \right).$$

(Since $\gamma \varpi_E^{n-1} \in \mathcal{O}_E$ and $k_E = k_F$, we regard $\gamma \varpi_E^{n-1} \bmod P_E$ as an element of k_F .)

Proof. By the definition of η_θ , we have

$$\begin{aligned} & \chi_{\eta_\theta}(1 + \varpi_E^n \text{diag}(k_0, \dots, k_{l-1})) \\ &= q^{(l-1)(m-1)} \sum_{a \in E^\times K^1 \setminus B} \psi(\text{Tr } \gamma a \varpi_E^n \text{diag}(k_0, \dots, k_{l-1}) a^{-1}). \end{aligned}$$

It follows from (2.2) and (2.4) that the set $\{\text{diag}(1, y_1, \dots, y_{l-1}) \mid y_i \in k_F^\times\}$ makes a complete system of representatives of $E^\times K^1 \setminus B$. For convenience, put $y_0 = 1$. Since

$$\varpi_E \text{diag}(1, y_1, \dots, y_{l-1}) \varpi_E^{-1} = \text{diag}(y_1, \dots, y_{l-1}, 1),$$

we have

$$\begin{aligned} & \text{Tr } \gamma \text{diag}(1, y_1, \dots, y_{l-1}) \varpi_E^{n-1} \text{diag}(k_0, \dots, k_{l-1}) \text{diag}(1, y_1, \dots, y_{l-1})^{-1} \\ & \equiv \gamma \varpi_E^{n-1} \sum_{i=0}^{l-1} k_i y_{i-n+1} / y_i \bmod P_F, \end{aligned}$$

where $y_i = y_{i \bmod l}$. By replacing y_i by $k_i y_{i-n+1} / y_i$, we get our lemma. \square

On K^n , the character of $\pi = \pi_\theta$ becomes a constant function on elliptic regular conjugacy classes.

Lemma 2.5. Let x be an elliptic regular element in K^n . Then

$$\chi_\pi(x) = q^{(n-2)(l-1)/2} \frac{(q^l - 1)}{q - 1}.$$

Proof. We use the Deligne-Kazhdan correspondence (Theorem 1.5). Since the correspondence preserves the conductor exponents, there exists a generic character θ' such that $f(\theta') = n$ and $\chi_{\pi'_{\theta'}} = \chi_{\pi_{\theta}}$. Since $\pi'_{\theta'}$ is trivial on $1 + P_D^n$ and its degree is $q^{(n-2)(l-1)/2} \frac{(q^l-1)}{q-1}$, $\chi_{\pi'_{\theta'}}(x) = q^{(n-2)(l-1)/2} \frac{(q^l-1)}{q-1}$ for $x \in 1 + P_D^n$. Consequently we have $\chi_{\pi_{\theta}}(x) = q^{n-1}(q^2+q+1)$ if $x \in K^n$ is elliptic regular. \square

The character formula on elliptic regular conjugacy classes outside E^\times can be obtained easily.

Lemma 2.6. *Let x be an elliptic regular element of B . If x satisfies the condition that $F(x) \not\cong E$ and x is not conjugate to an element of $F^\times K^n$, then $\chi_\pi(x) = 0$.*

Proof. See Lemma 3.3 in [14]. \square

3. CALCULATION OF GAUSS SUMS

In this section, we determine the Gauss sum part $G(y, n-i)$ explicitly. Since $G(y, n-i)$ depends only on $n-i \pmod l$ and $y \pmod P_E$, we have only to treat the character of η_θ on U_1^* by replacing n big enough.

Lemma 3.1. *Assume $n = 2m$. Then for $x \in U_1^*$,*

$$(3.1) \quad \chi_{\eta_\theta}(x) = \sum_{\sigma \in \text{Aut}_F E} \sum_{a \in H \backslash E^\times K^{m-1}} \rho_\theta(a^\sigma x a^{-1}).$$

Proof. It follows from Lemma 2.1 that $axa^{-1} \in H$ implies $a \in E^\times K^{m-1}$. Hence our lemma. \square

In the calculation of the sum in the above lemma, we use the E^\times -module structure of various objects. When E/F is a Galois extension, it is easy to treat. Thus we first assume E/F is Galois, i.e., $q \equiv 1 \pmod l$. We recall ξ is the diagonal matrix $\text{diag}(1, \zeta^{l-1}, \zeta^{l-2}, \dots, \zeta)$ where ζ is an l -th root of unity in F and ξ satisfies $\xi^l = 1$ and $\xi x \xi^{-1} = {}^\sigma x$ for $x \in E$ where σ is the generator of $\text{Gal}(E/F)$ determined by ${}^\sigma \varpi_E = \varpi_E \zeta$. By the explicit matrix form of E and A_i , we obtain:

$$(3.2) \quad \begin{array}{rcl} M_l(F) & = & E \oplus E\xi \oplus \dots \oplus E\xi^{l-1} \\ A^0 & = & \mathcal{O}_E \oplus \mathcal{O}_E\xi \oplus \dots \oplus \mathcal{O}_E\xi^{l-1} \\ A^1 & = & P_E \oplus P_E\xi \oplus \dots \oplus P_E\xi^{l-1} \\ & & \dots \dots \dots \\ A^{l-1} & = & P_E^{l-1} \oplus P_E^{l-1}\xi \oplus \dots \oplus P_E^{l-1}\xi^{l-1}. \end{array}$$

Lemma 3.2. *A complete system of representatives of $H \backslash E^\times K^{m-1}$ is given by*

$$\{1 + \varpi_E^{m-1} \alpha_1 \xi + \dots + \varpi_E^{m-1} \alpha_{l-1} \xi^{l-1} \mid \alpha_i \in k_F\}.$$

Proof. It is obvious from (3.2). \square

For $a = 1 + \alpha_1 \xi + \dots + \alpha_{l-1} \xi^{l-1} \in A^{m-1}$, $\rho(axa^{-1})$ for $x \in U_1^*$ can be expressed explicitly in terms of $\alpha_1, \dots, \alpha_{l-1}$. At first, we determine the coefficients of a^{-1} with respect to the F -basis $\{1, \xi, \dots, \xi^{l-1}\}$.

Lemma 3.3. For $a = \sum_{j=0}^{l-1} \alpha_j \xi^j$ ($\alpha_j \in E$), put

$$\begin{aligned} \Lambda(a) &= (\sigma^j \alpha_{i-j \bmod l})_{0 \leq i, j \leq l-1} \\ &= \begin{pmatrix} \alpha_0 & \sigma \alpha_{l-1} & \cdots & \sigma^{l-1} \alpha_1 \\ \alpha_1 & \sigma \alpha_0 & \cdots & \vdots \\ \vdots & \cdots & \cdots & \sigma^{l-1} \alpha_{l-1} \\ \alpha_{l-1} & \cdots & \sigma^{l-2} \alpha_1 & \sigma^{l-1} \alpha_0 \end{pmatrix} \in M_l(E) \end{aligned}$$

and let $\Lambda_k(a)$ be the $(1, k+1)$ -cofactor of $\Lambda(a)$. Then

$$a^{-1} = \sum_{j=0}^{l-1} \frac{\Lambda_j(a)}{|\Lambda(a)|} \xi^j,$$

where $|\Lambda(a)|$ is the determinant of $\Lambda(a)$.

Proof. By the map $\Lambda : M_l(F) \rightarrow M_l(E)$, we can embed $M_l(F)$ into $M_l(E)$. Then our lemma follows from Cramer's formula. \square

Lemma 3.4. Assume $n = 2m$ and $3(m-1) \geq 2m$. Let $c \in F^\times$, $y \in P_E^{m-1}$ and $a = 1 + \sum_{j=1}^{l-1} \alpha_j \xi^j \in K^{m-1}$. Then

$$\rho_\theta(ac(1+y)a^{-1}) = \theta(c(1+y))\psi_E \left(\sum_{j=1}^{l-1} (\gamma \alpha_j^{\sigma^j} \alpha_{l-j} - \sigma^{-j} \gamma \alpha_{l-j} \sigma^{-j} \alpha_j) y \right).$$

Proof. It is obvious that we may assume $c = 1$. Since

$$\begin{aligned} g^{-1} a g a^{-1} &= 1 + (g^{-1}(a-1)g - (a-1))a^{-1} \\ &= 1 + \left(\sum_{j=1}^{l-1} (\sigma^j g g^{-1} - 1) \alpha_j \xi^j \right) a^{-1}, \end{aligned}$$

$\sum_{j=1}^{l-1} (\sigma^j g g^{-1} - 1) \alpha_j \xi^j \in A^m$ and $\text{Tr}(\gamma x \xi^j) = 0$ for all $x \in E$, we have:

$$\begin{aligned} \rho_\theta(g^{-1} a g a^{-1}) &= \psi_\gamma \left(\left(\sum_{j=1}^{l-1} (\sigma^j g g^{-1} - 1) \alpha_j \xi^j \right) a^{-1} \right) \\ &= \psi_\gamma \left(\sum_{j=1}^{l-1} (\sigma^j g g^{-1} - 1) \alpha_j^{\sigma^j} (f_{l-j}(a)) \right), \end{aligned}$$

where $f_j(a) \in E$ is defined by $a^{-1} = \sum_{j=0}^{l-1} f_j(a) \xi^j$. Put $g = 1 + y$. In the last equation, $\gamma \in P_E^{1-n}$, $f_{l-j} \in P_E^{m-1}$ and $\sigma^j g g^{-1} - 1 \equiv \sigma^j y - y \pmod{P_E^{2m-2}}$. Thus we get

$$\rho_\theta(g^{-1} a g a^{-1}) = \psi_E \left(\sum_{j=1}^{l-1} (\sigma^{-j} \gamma f_{l-j}(a) \sigma^{-j} \alpha_j - \gamma^{\sigma^j} (f_{l-j}(a)) \alpha_j) y \right)$$

by virtue of $\text{tr}_E u^{\sigma^j} v = \text{tr}_E \sigma^{-j} u v$ for any $u, v \in E$. By Lemma 3.3,

$$f_{l-j}(a) = \frac{\Lambda_{l-j}(a)}{|\Lambda(a)|} \equiv \alpha_{l-j} \pmod{P_E^{2m-2}}.$$

By the assumption $3m-3 \geq 2m$, we obtain the desired formula. \square

Proposition 3.5. *Assume $q \equiv 1 \pmod{l}$, $n = 2m$ and $m \geq 3$.*

1. For $x \in U_1^*$,

$$\chi_{\eta_\theta}(x) = q^{(l-1)/2} \sum_{j=0}^{l-1} \theta(\sigma^j x).$$

2. For an even integer n and $y \in \mathcal{O}_F$, $G(y, n - 1) = q^{(l-1)/2}$. In particular, $G(y, n - 1)$ depends neither on n nor on y .

Proof. By Lemmas 3.1, 3.2 and 3.4, we have for $c \in F^\times$ and $y \in 1 + P_E$

$$\chi_{\eta_\theta}(c(1 + y)) = \sum_{i=0}^{l-1} \theta(c(1 + \sigma^i y)) \sum_{(\alpha_1, \dots, \alpha_{l-1}) \in (P_E^{m-1})^{l-1}} f(\alpha_1, \dots, \alpha_{l-1}; \sigma^i y),$$

where

$$f(\alpha_1, \dots, \alpha_{l-1}; y) = \psi_E \left(\sum_{j=1}^{l-1} (\gamma \alpha_j \alpha_{l-j}(a)^{\sigma^j} - \gamma^{\sigma^{-j}} \alpha_{l-j}(a) \alpha_j^{\sigma^{-j}})^{\sigma^i y} \right).$$

Put $S_j = \{(\alpha_1, \dots, \alpha_{l-1}) \in (P_E^{m-1}/P_E^m)^{l-1} \mid \alpha_k = 0 \text{ for } k < j, \alpha_j \neq 0\}$ and $I_j(y) = \sum_{(\alpha_1, \dots, \alpha_{l-1}) \in S_j} f(\alpha_1, \dots, \alpha_{l-1}; y)$. Then

$$\chi_{\eta_\theta}(c(1 + y)) = \sum_{i=0}^{l-1} \theta(c(1 + \sigma^i y)) \sum_{j=1}^{l-1} I_j(\sigma^i y).$$

If $\alpha_1 = \dots = \alpha_{(l-1)/2} = 0$, $f(\alpha_1, \dots, \alpha_{l-1}; y) = 0$. Thus we have

$$\sum_{j=(l+1)/2}^{l-1} I_j(y) = q^{(l-1)/2}.$$

For $1 \leq j \leq (l - 1)/2$, $I_j(y)$ is proportional to

$$\sum_{\alpha_{l-j} \in P_E^m/P_E^{m+1}} \psi_E((\gamma \alpha_j^{\sigma^j} \alpha_{l-j} - \sigma^{-j} \gamma \alpha_{l-j} \sigma^{-j} \alpha_j) y).$$

Since $\alpha_j \neq 0$, the map

$$\alpha_{l-j} \mapsto \gamma \alpha_j^{\sigma^j} \alpha_{l-j} - \sigma^{-j} \gamma \alpha_{l-j} \sigma^{-j} \alpha_j$$

is a bijection from P_E^{m-1}/P_E^m to k_F . Therefore $I_j(y) = 0$. Consequently we get the first part of our lemma. $G(y, n - 1) = q^{(l-1)/2}$ follows from Proposition 2.2 and the first part. \square

Next we assume $q - 1 \not\equiv 0 \pmod{l}$. In this situation, it is rather difficult to describe E -module structure of various objects since F has no l -th primitive root of unity and E/F is not Galois. In order to apply the result of Galois case, we use the base change lift of simple characters by Bushnell-Henniart([3]). Let ζ be an l -th root of unity and $L = F(\zeta)$. Then L/F is an unramified extension of degree $l - 1$ and the generator τ of $\text{Gal}(L/F)$ is determined by $\tau \zeta = \zeta^k$ where k is a generator of $(\mathbb{Z}/l\mathbb{Z})^\times$. We add the subscript L to the base changed objects. Then $M_l(L) = M_l(F) \otimes_F L$ and $E_L = E \otimes_F L \simeq EL$. E_L is a ramified Galois extension over L of degree l , an unramified extension over E of degree $l - 1$ with $\text{Gal}(E_L/E) = \text{Gal}(L/F) = \langle \tau \rangle$ and a non-Abelian Galois extension over F of degree $l(l - 1)$. (We embed E into E_L by the map: $x \mapsto x \otimes 1$).

As in the previous section, we identify $M_l(L)$ with $\text{End}_L E_L$ and $G_L = \text{GL}_l(L)$ with $\text{Aut}_L E_L$ by the L -basis $\{\varpi_E^{l-1}, \dots, \varpi_E, 1\}$ of E_L , which is also an \mathcal{O}_L -basis of \mathcal{O}_{E_L} . By the lattice flag $\{P_{E_L}^i\}_{i \in \mathbb{Z}}$, we define

$$A_L^i = \{f \in M_l(L) \mid f(P_{E_L}^j) \subset P_{E_L}^{j+i} \text{ for all } j \in \mathbb{Z}\}.$$

Put $K_L = (A_L^0)^\times$, $B_L = E_L^\times K_L$ and $K_L^i = 1 + A_L^i$ for $i \geq 1$. For a subgroup $M_L \subset B_L$ (resp. $M \subset B$), we set $M_L^1 = M \cap L^\times K_L$ (resp. $M^1 = M \cap F^\times K$). By the result of Kutzko (Theorem 1.7), it suffices to calculate the character of $\kappa = \kappa_\theta$ instead of π_θ . In fact, we have only to get the character of $\eta_\theta|_{B^1}$. Therefore we have only to treat the base change of $\eta_\theta|_{B^1}$ to B_L^1 where $B_L^1 = L^\times K_L$.

Definition 3.6. Let θ be a generic character of E^\times with $f(\theta) = n$ and $\theta(1+x) = \psi(\text{tr}_E(\gamma x))$ for $x \in P_E^m$. We define a base change lift θ_L of θ to L^\times by $\theta_L = \theta \circ n_{E_L/E}$. Then $\theta_L(1+x) = \psi_L(\text{tr}_{E_L/L} \gamma x)$ for $x \in P_{E_L}^m$. (Recall $m = [(n+1)/2]$.) The base change lift ρ_L of $\rho|_{H^1}$ to $H_L^1 = L^\times(1 + P_{E_L})K_L^m$ is defined by

$$\rho_L(h \cdot g) = \theta_L(h)\psi_L(\text{Tr } \gamma(g-1)) \quad \text{for } h \in L^\times(1 + P_{E_L}), \quad g \in K_L^m.$$

We define the base change η_L of $\eta|_{B^1}$ to B_L^1 by

$$\eta_L = \text{Ind}_{H_L^1}^{B_L^1} \rho_L.$$

By virtue of $\theta_L \circ \tau = \theta_L$, we have $\rho_L \circ \tau = \rho_L$. Thus we can define an extension $\tilde{\rho}_L$ of ρ_L to $H_L^1 \rtimes \langle \tau \rangle$ by

$$\tilde{\rho}_L(x \rtimes \tau) = \rho_L(x) \quad \text{for } x \in H_L^1.$$

Now we apply the result of Bushnell-Henniart ([3]) to our case and get the character relation between η_L and $\tilde{\eta}_L$. Put $U_{E_L, i} = L^\times(1 + P_{E_L}^i)$ for $i > 0$ and $U_{E_L, i}^* = U_{E_L, i} - U_{E_L, i+1}$. By (12.19) Corollary in [3] and the fact $\langle \tau \rangle$ -fixed space $(L^\times K_L^i)^{\langle \tau \rangle}$ is equal to $F^\times K^i$, the following result follows.

Proposition 3.7. *Let $x \in U_{E_L, 1}$. Between the set*

$$\{g \in H^1 \setminus (E^\times K^{m-1})^1 \mid gn_{E_L/E}(x)g^{-1} \in H^1\}$$

and the set

$$\{h \in H_L^1 \setminus (E_L^\times K_L^{m-1})^1 \mid hx^\tau h^{-1} \in H_L^1\},$$

there is a bijection ψ with the property

$$\rho_L(\psi(g)x^\tau(\psi(g))^{-1}) = \rho(gn_{E_L/E}(x)g^{-1}).$$

Combining this with Lemma 3.1, we have:

Lemma 3.8.

$$(3.3) \quad \chi_{\eta_\theta}(n_{E_L/E}(x)) = \sum_{\substack{a \in H_L^1 \setminus (E_L^\times K_L^{m-1})^1 \\ ax^\tau a^{-1} \in H_L}} \rho_L(ax^\tau a^{-1}).$$

Since $n_{E_L/E}(L^\times(1 + P_{E_L}^i)) = F^\times(1 + P_E^i)$, it suffices to calculate the right hand side of (3.3) for $x \in U_{L, 1}^*$.

As in the Galois case, set $\xi = \text{diag}(1, \zeta^{l-1}, \zeta^{l-2}, \dots, \zeta) \in M_l(L)$. Then ξ satisfies $\xi^l = 1$, ${}^\tau \xi = \xi^k$ and

$$\xi x \xi^{-1} = {}^\sigma x \quad \text{for any } x \in E_L,$$

where σ is the generator of $\text{Gal}(E_L/L)$ determined by $\sigma\varpi_E = \varpi_E\zeta$. Moreover we have $\tau\sigma\tau^{-1} = \sigma^k$ and

$$(3.4) \quad \begin{aligned} M_l(L) &= E_L \oplus E_L\xi \oplus \cdots \oplus E_L\xi^{l-1} \\ A_L^0 &= \mathcal{O}_{E_L} \oplus \mathcal{O}_{E_L}\xi \oplus \cdots \oplus \mathcal{O}_{E_L}\xi^{l-1} \\ A_L^1 &= P_{E_L} \oplus P_{E_L}\xi \oplus \cdots \oplus P_{E_L}\xi^{l-1} \\ &\dots\dots\dots \\ A_L^{l-1} &= P_{E_L}^{l-1} \oplus P_{E_L}^{l-1}\xi \oplus \cdots \oplus P_{E_L}^{l-1}\xi^{l-1}. \end{aligned}$$

We note that any element of K_L^1 can be written in the form $(1 + \alpha_1\xi + \alpha_2\xi^2 + \cdots + \alpha_{l-1}\xi^{l-1})$ for $\alpha_i \in P_{E_L}$.

Lemma 3.9. *Let $i < m$ and $a = 1 + \alpha_1\xi + \alpha_2\xi^2 + \cdots + \alpha_{l-1}\xi^{l-1}$ for $\alpha_j \in \mathcal{O}_E$ and $x \in U_{E_L, i}^*$. Then $ax^\tau a^{-1} \in H_L$ is equivalent to $\alpha_j \in P_{m-i}$ and $\alpha_{kj} = \tau^{j-1}\alpha_k$ for all j . (The suffix of α_j is extended to \mathbb{Z} by $\alpha_j = \alpha_{j \bmod l}$.)*

Proof. It follows from Lemma 3.2 that if $a^{-1}x^\tau a \in H_L$, there exist $\gamma_0 \in \mathcal{O}_E^\times$ and $\gamma_j \in P_{E_L}^m$ for $1 \leq j \leq l-1$ such that

$$(1 + \alpha_1\xi + \cdots + \alpha_{l-1}\xi^{l-1})x = \gamma_0(1 + \gamma_1\xi + \gamma_2\xi^2 + \cdots + \gamma_{l-1}\xi^{l-1}) \\ (1 + \tau\alpha_1\xi^k + \tau\alpha_2\xi^{2k} + \cdots + \tau\alpha_{l-1}\xi^{(l-1)k}).$$

It implies

$$\begin{aligned} x &= \gamma_0(1 + \gamma_{l-k}\sigma^{l-k}\tau\alpha_1 + \gamma_{l-2k}\sigma^{l-2k}\tau\alpha_2 + \cdots + \gamma_k\sigma^k\tau\alpha_{l-1}) \\ \alpha_k\sigma^k &= \gamma_0(\gamma_k + \tau\alpha_1 + \gamma_{l-k}\sigma^{l-k}\tau\alpha_2 + \cdots + \gamma_{2k}\sigma^{2k}\tau\alpha_{l-1}) \\ &\dots\dots\dots \\ \alpha_{l-k}\sigma^{l-k}x &= \gamma_0(\gamma_{l-k} + \gamma_{l-2k}\sigma^{l-2k}\tau\alpha_1 + \cdots + \tau\alpha_{l-1}). \end{aligned}$$

Thus we have

$$\alpha_{jk}\sigma^{jk}x = x^\tau\alpha_j \bmod P_{E_L}^m \quad (j \in (\mathbb{Z}/l\mathbb{Z})^\times).$$

By eliminating $\alpha_{k^2}, \alpha_{k^3}, \dots, \alpha_{k^{l-1}}$, we get

$$\alpha_k = n_{E_L/E}(x)\sigma^k n_{E_L/E}(x)^{-1}\alpha_k \bmod P_{E_L}^m.$$

Since $n_{E_L/E}(x)\sigma^k n_{E_L/E}(x)^{-1} \in 1 + P_E^i - P_E^{i+1}$, $\alpha_k \in P_{E_L/E}^{m-i}$. By $x\sigma^{k^j}x^{-1} \in 1 + P_{E_L}^i$, we obtain $\alpha_{kj} \in P_{E_L}^{m-i}$ and $\alpha_{kj} = \tau^{(j-1)}\alpha_k \bmod P_{E_L}^m$ for $j = 1, \dots, l-1$. \square

Lemma 3.10. *Let $x \in 1 + P_{E_L} - P_{E_L}^2$ and $a = 1 + \sum_{j=1}^{l-1} \tau^{(j-1)}\alpha\xi^{kj}$ for $\alpha \in P_{E_L}^{m-1}$. Then*

$$\rho_L(ax^\tau a^{-1}x^{-1}) = \psi_E(\text{tr}_{E_L/E}(\alpha^{\sigma^k\tau^{(l-1)/2}})\alpha) \text{tr}_{E_L/E}(x-1).$$

Proof. By Lemma 3.9, $\tau a a^{-1} \in H_L$. Since $\rho_L(\tau a a^{-1}) = 1$, it implies $\rho_L(ax^\tau a^{-1}g^{-1}) = \rho_L(axa^{-1}g^{-1})$. Thus we can prove the lemma by the same way as Lemma 3.4. \square

It is time to get the character value of χ_η on U_1^* .

Proposition 3.11. *Let $x \in 1 + P_{E_L} - P_{E_L}^2$ and $n = 2m > 6$. Then*

$$\chi_\eta(n_{E_L/E}(x)) = -q^{(l-1)/2}\theta(n_{E_L/E}(x))$$

and

$$G(y, j) = -q^{(l-1)/2}$$

for all $y \in k_F$ and j odd.

Proof. By Proposition 3.7, Lemmas 3.8, 3.9, 3.10 and 3.11, we have:

$$\chi_\eta(n_{E_L/E}(x)) = \theta_L(x) \sum_{\alpha \in P_{E_L}^{m-1}/P_{E_L}^m} \psi_E(\text{tr}_{E_L/E}(\alpha^{\sigma\tau^{(l-1)/2}} \alpha) \text{tr}_{E_L/E}(x-1)).$$

Let E' be the $\langle \sigma\tau^{(l-1)/2} \rangle$ -fixed field. Then E_L/E' is a quadratic unramified extension. Since $\alpha^{\sigma\tau^{(l-1)/2}} \alpha = n_{E_L/E'}(\alpha)$ and $n_{E_L/E'}$ induces a surjection from $\varpi_E^{m-1} \mathcal{O}_{E_L}/1 + P_{E_L}$ to $\varpi_E^{2m-2} \mathcal{O}_{E'}/1 + P_{E'}$ and each fiber of the map has $q^{(l-1)/2} + 1$ elements. Thus we get

$$\chi_\eta(n_{E_L/E}(x)) = (1 - (q^{(l-1)/2} + 1))\theta_L(x).$$

and it follows from Proposition 2.2 that $G(y, j) = -q^{(l-1)/2}$ for all $y \in k_F$ and j odd. \square

Putting all these together, we can state the character formula.

Theorem 3.12. *Let E be a ramified extension of F with degree l , θ a generic quasi-character of E^\times with $f(\theta) = n$ and $\pi = \pi_\theta$ the irreducible supercuspidal representation of $\text{GL}_l(F)$ defined in section 1. Put $U_0 = F^\times \mathcal{O}_E^\times$, $U_j = F^\times(1 + P_E^j)$ and $U_j^* = U_j - U_{j+1}$ for $j \geq 1$. Let x be an elliptic regular element of $\text{GL}_l(F)$ and $\text{Aut}_F E$ the group of automorphism of E over F .*

1. *If $F(x)/F$ is unramified, then*

$$\chi_\pi(x) = \begin{cases} 0 & x \notin F^\times(1 + P_{F(x)}^n), \\ q^{(n-2)(l-1)/2} \frac{(q^l-1)}{q-1} \theta(c) & x = c(1+y) \\ & \text{for } c \in F^\times, y \in P_{F(x)}^n. \end{cases}$$

2. *If $F(x)/F$ is ramified and $F(x) \not\subseteq E$, then*

$$\chi_\pi(x) = \begin{cases} 0 & \text{if } x \notin F^\times(1 + P_{F(x)}^{n-1}), \\ q^{(n-2)(l-1)/2} \theta(c) \text{Kl}((\gamma\varpi_E^{n-1})^l \prod_{j=0}^{l-1} k_j) & \\ & \text{if } x = c(1 + \varpi_E^{n-1} \text{diag}(k_0, \dots, k_{l-1}) + z) \\ & \text{for } c \in F^\times, k_i \in k_F^\times, z \in P_{F(x)}^n, \\ q^{(n-2)(l-1)/2} \frac{(q^l-1)}{q-1} \theta(c) & \\ & \text{if } x = c(1+y) \text{ for } c \in F^\times, y \in P_{F(x)}^n. \end{cases}$$

3. *When $x \in E$, then*

$$\chi_\pi(x) = \begin{cases} \left(\frac{q}{l}\right)^n \sum_{\sigma \in \text{Aut}_F E} \theta(\sigma x) & \text{if } x \in E^\times - U_0, \\ (-1)^{n-j} q^{j(l-1)/2} \sum_{\sigma \in \text{Aut}_F E} \theta(\sigma x) & \\ & \text{if } x \in U_j^* \text{ for } 1 \leq j \leq n-1, \\ q^{(n-2)(l-1)/2} \theta(c) \text{Kl}((\gamma\varpi_E^{n-1} x_0)^l) & \\ & \text{if } x = c(1 + \varpi_E^{n-1} x_0) \text{ for } c \in F^\times, x_0 \in \mathcal{O}_E^\times, \\ q^{(n-2)(l-1)/2} \frac{(q^l-1)}{q-1} \theta(c) & \\ & \text{if } x = c(1+y) \text{ for } c \in F^\times, y \in P_E^n. \end{cases}$$

(See (2.6) for the definition of the Kloosterman sum $\text{Kl}(a)$.)

Proof. It follows from (1.8), Theorem 1.7, Lemmas 2.4, 2.5, 2.6, Propositions 2.2, 3.5 and 3.11 \square

Remark. By Theorem 1.6, the character formula of the representation π'_θ of D^\times is given by the same formula for π_θ .

REFERENCES

- [1] J. Arthur and L. Clozel, Simple algebras, base change, and the advanced theory of the trace formula, *Annals of Math. Studies* **120**, Princeton Univ. Press, 1989.
- [2] C. J. Bushnell and A. Fröhlich, Gauss Sums and p -adic Division Algebras, *Lecture Notes in Math.* **987**, Springer, Berlin, 1983.
- [3] C. J. Bushnell and G. Henniart, *Local Tame Lifting For $GL(N)$ I: Simple characters*, *Inst. Hautes Études Sci. Publ. Math. No. 83* (1996), 105–233.
- [4] H. Carayol, *Représentation cuspidales du groupe linéaire* *Ann. Scient. Éc. Norm. Sup.* **17** (1984), 191–226.
- [5] L. Corwin and R. Howe, *Computing characters of tamely ramified division algebras*, *Pac. J. Math.* **73** (1977), 461–477.
- [6] L. Corwin, A. Moy and P. J. Sally, Jr., *Supercuspidal character formulas for GL_l* , *Contemp. Math.* **191**, AMS, 1995, 1–11.
- [7] S. M. Debacker, *Supercuspidal characters of GL_l , l a prime*, Ph.D. thesis, Univ. of Chicago, 1997.
- [8] P. Deligne, D. Kazhdan, and M.-F. Vignéras, *Représentations des algèbres centrales simples p -adiques*, in: *Représentations des Groupes Réductifs sur un Corps Local*, Herman, Paris, 1984, pp. 33–117.
- [9] R. Godement and H. Jacquet, *Zeta functions of simple algebras*, *Lecture Notes in Math.* **260**, Springer, Berlin, 1972.
- [10] G. Henniart, *Correspondance de Jaquet-Langlands explicite I : le cas modéré de degré premier* in: *Séminaire de Théorie des Nombres, Paris 1990-91*, *Progress in Math.* **108**, Birkhäuser, Basel, 1993, pp. 85–114.
- [11] R. Howe, *Kirillov theory for compact p -adic groups*, *Pac. J. Math.* **73**, (1977), 365–381.
- [12] ———, *Tamely ramified supercuspidal representations of $GL_n(F)$* , *Pac. J. Math.* **73** (1977), 437–460.
- [13] H. Hijikata, H. Saito, and M. Yamauchi, *Representations of quaternion algebras over local fields and trace formula of Hecke operators*, *J. Number Theory* **43** (1993), 123–167.
- [14] P. Kutzko, *Character formulas for supercuspidal representations of GL_l , l a prime*, *Amer. J. Math.* **109** (1987), 201–222.
- [15] A. Moy, *Local constant and the tame Langlands correspondence*, *Amer. J. Math.* **108** (1986), 863–930.
- [16] F. Murnaghan, *Characters of supercuspidal representations of $SL(n)$* , *Pac. J. Math.* **170**, (1995), 217–235.
- [17] ———, *Characters of supercuspidal representations of classical groups*, *Ann. Sci. Éc. Norm. Sup.* (4) **29**, (1996), 49–105.
- [18] ———, *Local character expansions and Shalika germs for $GL(n)$* , *Math. Ann.* **304**, (1996), 423–455.
- [19] J. Rogawski, *Representations of $GL(n)$ and division algebras over p -adic field*, *Duke Math. J.* **50** (1983), 161–196.
- [20] T. Takahashi, *Characters of cuspidal unramified series for central simple algebras of prime degree*, *J. Math. Kyoto Univ.* **32-4** (1992), 873–888.
- [21] ———, *Character formula for representations of local quaternion algebras (Wildly ramified case)*, *J. Math. Kyoto Univ.* **36-1** (1996), 151–197.
- [22] ———, *Character formula for the representations of GL_3 over non-archimedean local field*, preprint.

DEPARTMENT OF MATHEMATICS AND INFORMATION SCIENCE, COLLEGE OF INTEGRATED ARTS AND SCIENCES, OSAKA PREFECTURE UNIVERSITY

E-mail address: takahasi@mi.cias.osakafu-u.ac.jp