

## The group-quark matrix ?

**Koichiro Harada/原田耕一郎**  
Ohio State University/オハイオ州立大学

### §1 The group-quark matrix ?

The main purpose of this article is to propose a problem. Let us consider the following  $3 \times 3$  matrix whose entries are finite groups.

$$\mathcal{A} = \begin{bmatrix} U_4(2).2 & S_6(2) & O_8^+(2) \\ U_6(2).2 & Conway_2 & Conway_3 \\ {}^2E_6(2).2 & Fischer_4 & Monster \end{bmatrix}$$

The orders of relevant simple groups are :

$$\begin{aligned} |U_4(2)| &= 25920 = 2^6 3^4 5 \\ |S_6(2)| &= 1451520 = 2^9 3^4 5 \cdot 7 \\ |O_8^+(2)| &= 174182400 = 2^{12} 3^5 5^2 7 \\ |U_6(2)| &= 2^{15} 3^6 5 \cdot 7 \cdot 11 \\ |Conway_2| &= 2^{18} 3^6 5^3 7 \cdot 11 \cdot 23 \\ |Conway_3| &= 2^{21} 3^9 5^4 7^2 11 \cdot 13 \cdot 23 \\ |{}^2E_6(2)| &= 2^{36} 3^9 5^2 7^2 11 \cdot 13 \cdot 17 \cdot 19 \\ |Fischer_4| &= 2^{41} 3^{13} 5^6 7^2 11 \cdot 13 \cdot 17 \cdot 19 \cdot 23 \cdot 31 \cdot 47 \\ |Monster| &= 2^{46} 3^{20} 5^9 7^6 11^2 13^3 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 41 \cdot 47 \cdot 59 \cdot 71 \end{aligned}$$

Note that Conway groups are numbered according to their orders. In particular,  $|Conway_1| = 2^{10} 3^7 5^3 7 \cdot 11 \cdot 23$ .  $U_4(2).2$  is the extension of  $U_4(2)$  by an outer automorphism of order 2, and  $U_6(2).2$  and  ${}^2E_6(2).2$  are analogously defined.

The columns of the matrix  $\mathcal{A}$  are indexed by the Dynkin diagrams of type  $E_6, E_7$  and  $E_8$ . Appearing in the first row of  $\mathcal{A}$  are the simple components of the Weyl groups of type  $E_6, E_7$  and  $E_8$ . The correct indexing of the rows of the matrix  $\mathcal{A}$  is left for the future research. We can, perhaps, index the rows of  $\mathcal{A}$  by three generations of quarks  $ud, cs, tb$  (up-down, charm-strange, top-bottom).

Let us next give the ‘transpose-inverse=tra-inv’ of the matrix  $\mathcal{A}$ .

$${}^t\mathcal{A}^{-1} = \begin{bmatrix} 2^{1+4} \cdot (S_3 \times S_3) & 2^{1+6^*} \cdot (S_3 \times S_3) & 2^{1+8} \cdot (S_3 \times S_3 \times S_3) \\ 2^{1+8} \cdot U_4(2) \cdot 2 & 2^{1+8} \cdot S_6(2) & 2^{1+8} O_8^+(2) \\ 2^{1+20} \cdot U_6(2) \cdot 2 & 2^{1+22} \text{Conway}_2 & 2^{1+24} \text{Conway}_3 \end{bmatrix}$$

If  $A_{ij}$  is the  $(i, j)$  entry of the matrix  $\mathcal{A}$ , then the corresponding entry of  ${}^t\mathcal{A}^{-1}$  is the centralizer of an involution in the center of a Sylow 2-subgroup of the group  $A_{ij}$ . Here  $2^{1+2n}$  denotes the extral-special group of order  $2^{1+2n}$ . An exception is  $2^{1+6^*}$ , which is almost extra-special but not exactly so. The main problem proposed here is : **Investigate the group-quark matrix  $\mathcal{A}$  algebro-geometrically.**

## §2 $\Gamma_{27}$ and $\Gamma_{28}$

Let  $S$  be the cubic surface defined in the projective space  $P^4(\mathbb{C})$  by the equations :

$$\begin{cases} x_0^3 + x_1^3 + x_2^3 + x_3^3 + x_4^3 = 0 \\ x_0 + x_1 + x_2 + x_3 + x_4 = 0. \end{cases}$$

The (projective) line defined by

$$\begin{cases} x_0 = 0 \\ x_1 + x_2 = 0 \\ x_3 + x_4 = 0 \end{cases}$$

lies completely on the surface  $S$ . Applying the permutations on the index set  $\{0, 1, 2, 3, 4\}$ , 15 lines on  $S$  can be obtained.

Next, let  $\alpha (= \frac{1 \pm \sqrt{5}}{2})$  be a zero of the quadratic equation :

$$X^2 - X - 1 = 0,$$

then the line defined by :

$$\begin{cases} x_0 + \alpha x_3 + x_4 = 0 \\ x_1 + x_3 + \alpha x_4 = 0 \\ x_2 - \alpha(x_3 + x_4) = 0 \end{cases}$$

is also completely on the surface  $S$ . Applying the permutations on  $\{0, 1, 2, 3, 4\}$  again, 12 lines can be obtained. Therefore there are

altogether 27 lines on  $S$ . That this is the exact number of lines on  $S$  comes from the theory of algebraic geometry, although our special case itself was known already in the middle of the 19th century.

**Theorem.** A general (complex) cubic surface contains exactly 27 lines.

Let  $\Gamma_{27}$  be the graph of 27 lines with their configuration on a general cubic surface. Then  $\Gamma_{27}$  satisfies the following properties :

(1). Any line  $A$  of  $\Gamma_{27}$  meets exactly ten other lines of  $\Gamma_{27}$ . Those ten lines split into five pairs  $(B_1, C_1), \dots, (B_5, C_5)$ , and if  $i = 1, 2, 3, 4, 5$ , then  $B_i$  and  $C_i$  meet and the triangle  $AB_iC_i$  is formed. There are  $5 \cdot 27/3 = 45$  triangles so formed. (Note. If  $i \neq j$ , then  $B_i$  and  $C_j$  do not meet. In particular, there are no three lines that meet at a point. This applies to a general cubic surface. A specialization of it may contain three lines that meet at a point.)

(2). Let  $ABC, A'B'C'$  be any two triangles having no side in common. Then they determine uniquely a third triangle  $A''B''C''$  such that each of three triples of lines  $\{A, A', A''\}, \{B, B', B''\}, \{C, C', C''\}$  intersect and form three new triangles  $AA'A'', BB'B'', CC'C''$ .

Those two properties (1), (2) uniquely determines the configuration of 45 triangles formed by the elements of  $\Gamma_{27}$ .

**Theorem(C.Jordan).**  $\text{Aut}(\Gamma_{27}) \cong U_4(2).2 \cong \text{Aut}(U_4(2))$

This is the (1, 1) entry of the matrix  $\mathcal{A}$ . The isomorphisms of simple groups

$$U_4(2) \cong S_4(3) \cong O_5(3) \cong O_6^-(2)$$

is significant in the history of group theory.

Let us next discuss the (1, 2) entry of the matrix  $\mathcal{A}$ . The graph of the quartic curve

$$x^4 + y^4 + x^2y^2 - 8(x^2 + y^2) + 16.25 = 0$$

is drawn at the end of this article.

It is easy to see that  $28 = 4 + (12 \cdot 4/2)$  double tangents to the curve can be drawn. If the constant 16.25 is replaced by a number smaller than about 15.5 then four regions merge into a single region and if it is replaced by a number larger than about 17.5, then we get four convex regions and only 24 double tangents can actually be visible.

In general, it is known :

**Theorem.** A nonsingular (complex) plane curve of degree 4 possesses exactly 28 double tangents.

The number of double tangents to a nonsingular plane curve of degree  $m$  is given by the formula of Plücker :

$$\text{Number of double tangents} = \frac{1}{2}m(m-2)(m^2-9).$$

Let  $\Gamma_{28}$  be the set of 28 double tangents. The configuration satisfied by the 28 double tangents was investigated by Steiner, Aronhold and many others.

(1). (Steiner) Let  $x_1, y_1$  be two distinct elements of  $\Gamma_{28}$ . Then there exist five pairs  $(x_2, y_2), (x_3, y_3), \dots, (x_6, y_6)$  of elements in  $\Gamma_{28}$  and if we put

$$\mathfrak{S} = \{(x_i, y_i) | i = 1, 2, 3, \dots, 6\}$$

then, the eight tangent points of any pair of double tangents  $(x_i, y_i), (x_j, y_j) \in \mathfrak{S}$  lie on a same conic (an irreducible plane curve of degree 2).  $\mathfrak{S}$  is called a *Steiner complex*.  $\Gamma_{28}$  possesses 63 Steiner complexes in total.

Let  $P_1, \dots, P_7$  be seven points given in the complex plane. The cubic curves passing through these seven points form a vector space  $\mathfrak{T}$ . Every pair of curves  $\{C_1, C_2\}$  of  $\mathfrak{T}$  intersect two more points by Bézout's theorem. If these two points coincide then the pair  $\{C_1, C_2\}$  possesses a common tangent. The totality of common tangents so obtained forms a plane curve  $D'$  of class 4, or equivalently the dual curve of a plane curve of degree 4.

The dual of the statement above will read as follows.

(2)(Aronhold). Let  $L_1, \dots, L_7$  be seven lines on the plane. The totality of all curves of class 3 containing these seven lines forms a vector space  $\mathfrak{X}'$ . Every pair of curves  $\{C'_1, C'_2\}$  in  $\mathfrak{X}'$  contains two more lines  $\{L_8, L_9\}$  in common. If  $L_8 = L_9$ , then the pair  $\{C'_1, C'_2\}$  possesses a tangent point  $z$  and  $z$  is on a curve  $D$  of degree 4 uniquely determined by  $L_1, \dots, L_7$ . Moreover,  $L_1, \dots, L_7$  are double tangents of this curve  $D$ .

Let  $D$  be the curve of degree 4 uniquely determined by the seven lines  $\{L_1, \dots, L_7\}$ . Then  $D$  possesses 28 double tangents  $\Gamma_{28} = \{L_1, L_2, \dots, L_{28}\}$ . Moreover, the following properties hold.

- (i).  $L_1, \dots, L_7$  is a maximal aszygetic set (defined below) of  $\Gamma_{28}$ .
- (ii). The remaining 21 double tangents are rationally constructible by  $L_1, \dots, L_7$  (their coefficients are rational functions of the coefficients of  $L_1, \dots, L_7$ ).
- (iii). Every curve of degree 4 without double points can be obtained by this construction.
- (iv). Every aszygetic set of seven double tangents of  $\Gamma_{28}$  defines  $D$ .

Let  $L_1, L_2, L_3$  be three distinct lines in  $\Gamma_{28}$ . Those three lines determine six tangent points. If those six tangent points are on a same conic, then the triple  $\{L_1, L_2, L_3\}$  is called *syzygetic*. In the contrary case, the triple is called *asyzygetic*. A subset  $S$  of  $\Gamma_{28}$  is called aszygetic if every triple of  $S$  is aszygetic.

Let us call a maximal aszygetic seven-line set mentioned in (i) an *Aronhold set*. Therefore, an Aronhold set is a maximal aszygetic subset of  $\Gamma_{28}$  consisting of seven elements. It is known that  $\Gamma_{28}$  contains exactly 288 Aronhold sets.

**Theorem(Jordan).**  $Aut(\Gamma_{28}) \cong S_6(2)$ .

Note that  $|S_6(2)| = 288 \times 7!$ . In fact,  $S_6(2)$  transitively permutes all Aronhold sets and the fixing subgroup of an Aronhold set  $A$  acts as the symmetric group of degree 7 on  $A$ .

$\Gamma_{28}$  can not be determined only by vertices and edges since  $Aut(\Gamma_{28})$  acts doubly transitively on the 28 points. Therefore,  $\Gamma_{28}$  is not a

graph in an usual sense.

Let  $L_1, L_2$  be a pair of elements in  $\Gamma_{28}$ , then there are 10 elements  $X$  in  $\Gamma_{28}$  such that  $\{L_1, L_2, X\}$  is a syzygetic triple. In fact, all such  $X$  are in the Steiner complex determined by the pair  $\{L_1, L_2\}$ . Therefore,  $\Gamma_{28}$  possesses  $28 \cdot 27 \cdot 10 / 6 = 1260$  syzygetic triples. If all syzygetic triples are given in  $\Gamma_{28}$ , then the configuration of  $\Gamma_{28}$  is completely determined. The author is not aware if any combinatorial characterization of  $\Gamma_{28}$  is known. (Note. A combinatorial characterization of  $\Gamma_{27}$  is known as mentioned in this article before.)

Let  $L$  be an element of  $\Gamma_{28}$ . Consider  $\Gamma'_{27} = \Gamma_{28} \setminus \{L\}$ . For a pair of elements  $X, Y$  in  $\Gamma'_{27}$ , if  $L, X, Y$  is syzygetic, connect  $X$  and  $Y$  by an edge. Then a graph of 27 vertices and 135 edges is obtained. The  $\Gamma'_{27}$  is isomorphic with  $\Gamma_{27}$  discussed before (Geiger, 1869).

We have thus obtained the (1,2) entry of the matrix  $\mathcal{A}$ .

**Problem.** Define the (1,3) entry of the group-quark matrix  $\mathcal{A}$  algebro-geometrically.

Since

$$[O_8^+(2) : S_6(2)] = 120,$$

the algebro-geometric model on which  $O_8^+(2)$  acts should contain 120 elements in it. Let us denote the object by  $\Gamma_{120}$ . The fixing subgroup of a point  $\alpha$  of  $\Gamma_{120}$  should be  $S_6(2)$ .

Therefore,  $\Gamma_{120}$  is, as an  $O_8^+(2)$ -set, equivalent to the quotient space  $O_8^+(2)/S_6(2)$ . The action of  $O_8^+(2)$  on  $O_8^+(2)/S_6(2)$  is well known and it induces a rank 3-permutation representation. Equivalently one point stabilizer  $S_6(2)$  has exactly two orbits on the remaining 119 points  $\Gamma_{120} \setminus \{\alpha\}$ . The suborbit lengths are 56 and 63, and the stabilizer of a point in  $S_6(2)$  is  $U_4(2)$  or  $E_{32} \cdot S_5$  respectively. Let us write

$$\Gamma_{120} = \{\alpha\} + \Delta + \Omega$$

where,  $|\Delta| = 56, |\Omega| = 63$ .

We are assuming that the configuration graph of  $\Gamma_{120}$  contains  $\Gamma_{28}$  as a subgraph. Therefore, we should be able to identify  $\Delta$  and  $\Omega$  in terms of  $\Gamma_{28}$ .  $\Omega$  is of length 63 and so it is natural to assume that  $\Omega$  is the totality of all Steiner complexes.

There are 28 double tangents and so obviously there are 56 tangent points. Therefore, it is natural again to choose  $\Delta$  to be the set of all (double) tangent points of the plane curve of degree 4 that we initially began with.

In Heinrich Weber's *Lehrbuch der Algebra*, Vol II(1899), there is a 50 page chapter entirely devoted to the structure of  $\Gamma_{28}$ . In it, it is proved also that  $S_6(2)$  is the automorphism group of the configuration.

There are other 120 mathematical objects.

- (1). A nonsingular plane curve of degree 5 possesses 120 double tangents (easy by Plücker's formula).
- (2). There is a curve ( called del Pizzo surface) of degree 6 and of genus 4 possessing 120 tritangents planes.
- (3). The root system of type  $E_8$  possesses 240 roots. If the sign of each root is ignored then a set  $\Gamma$  of 120 objects and its graph are obtained.

It must be an interesting problem to investigate the configuration  $\Gamma_{120}$  purely group theoretically also.

### §3 The second and third rows of $\mathcal{A}$ .

The second and third rows of the group-quark matrix are up in the air at this moment. McKay [Finite Groups, Proceedings of Symposia in Pure Mathematics, Vol. 37, Amer. Math. Soc. 1980] observed that if  $s$  and  $t$  are involutions of the Monster both of which are conjugate to the involutions of  $2A$  type, then its product  $st$  belongs to the conjugacy classes of the Monster of type

$$1A, 2A, 3A, 4A, 5A, 6A, 3C, 4B, 2B.$$


Recall that if  $-\alpha_0$  is the highest root of the Lie algebra of type  $E_8$ , then

$$1\alpha_0 + 2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 5\alpha_4 + 6\alpha_5 + 3\alpha_6 + 4\alpha_7 + 2\alpha_8 = 0.$$

The numbers  $\{1, 2, 3, 4, 5, 6, 3, 4, 2\}$  are called the weights of  $E_8$ . McKay lists *Fischer*<sub>3</sub> and *Fischer*<sub>4</sub> as groups having similar property with respect to  $E_6$  and  $E_7$ , respectively. *Fischer*<sub>3</sub> is replaced by  ${}^2E_6(2)$  in this article, since it fits better if we consider the (2,1) entry of the tra-inv  ${}^t\mathcal{A}^{-1}$  of the group-quark matrix.

Similar coincidences between weights of Dynkin diagrams and orders of groups elements have been observed by Glauberman and Norton [to appear in the Proceedings of Monster Workshop at Montreal, 1999] . At Kyoto symposium, the (2,1) and (3,1) entries of the matrix  $\mathcal{A}$  were the sporadic simple groups *Suzuki* and *Fischer*<sub>3</sub>, respectively. The new entries  $U_6(2)$  and  ${}^2E_6(2)$ , however, appear to fit its tra-inv matrix  ${}^t\mathcal{A}^{-1}$  better, although leaving the main realm of the 3-transposition groups may be a problem.

I thank M.Kitazume, J.Matsuzawa, T.Asai, and J.McKay for valuable discussions.



$$x^4 + y^4 + x^2y^2 - 8(x^2 + y^2) + 16.25 < 0$$