

LEXICOGRAPHIC GRÖBNER BASES OF TORIC IDEALS
ARISING FROM ROOT SYSTEMS

大杉英史
HIDEFUMI OHSUGI

立教大学理学部数学科
Department of Mathematics, Rikkyo University

ABSTRACT. The present paper is a brief draft based on a joint work with Takayuki Hibi. Gröbner bases of toric ideals arising from root systems are studied.

INTRODUCTION

Let $\mathcal{A} \subset \mathbb{Z}^n$ be a finite set and let $K[t, t^{-1}, s] = K[t_1, t_1^{-1}, \dots, t_n, t_n^{-1}, s]$ denote the Laurent polynomial ring over a field K . We associate each $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}^n$ with the monomial $t^\alpha s = t_1^{\alpha_1} \dots t_n^{\alpha_n} s \in K[t, t^{-1}, s]$ and write $\mathcal{R}_K[\mathcal{A}]$ for the subalgebra of $K[t, t^{-1}, s]$ generated by all monomials $t^\alpha s$ with $\alpha \in \mathcal{A}$. Let $K[\mathbf{x}] = K[\{x_\alpha; \alpha \in \mathcal{A}\}]$ denote the polynomial ring in $\#\mathcal{A}$ variables over K and $I_{\mathcal{A}} \subset K[\mathbf{x}]$ the kernel of the surjective homomorphism $\pi : K[\mathbf{x}] \rightarrow \mathcal{R}_K[\mathcal{A}]$ defined by setting $\pi(x_\alpha) = t^\alpha s$ for all $\alpha \in \mathcal{A}$. The ideal $I_{\mathcal{A}}$ is called the *toric ideal* of the configuration \mathcal{A} . It is known [9] that if $I_{\mathcal{A}}$ possesses a squarefree initial ideal, then the convex hull of \mathcal{A} possesses a unimodular triangulation.

Fix $n \geq 2$. Let \mathbf{e}_i denote the i -th unit coordinate vector of \mathbb{R}^n . We write \mathbf{A}_{n-1}^+ , \mathbf{B}_n^+ , \mathbf{C}_n^+ , \mathbf{D}_n^+ and \mathbf{BC}_n^+ for the set of positive roots of root systems \mathbf{A}_{n-1} , \mathbf{B}_n , \mathbf{C}_n , \mathbf{D}_n and \mathbf{BC}_n , respectively ([3, pp. 64 – 65]):

$$\begin{aligned} \mathbf{A}_{n-1}^+ &= \{\mathbf{e}_i - \mathbf{e}_j; 1 \leq i < j \leq n\}; \\ \mathbf{B}_n^+ &= \{\mathbf{e}_i; 1 \leq i \leq n\} \cup \{\mathbf{e}_i + \mathbf{e}_j; 1 \leq i < j \leq n\} \cup \{\mathbf{e}_i - \mathbf{e}_j; 1 \leq i < j \leq n\}; \\ \mathbf{C}_n^+ &= \{2\mathbf{e}_i; 1 \leq i \leq n\} \cup \{\mathbf{e}_i + \mathbf{e}_j; 1 \leq i < j \leq n\} \cup \{\mathbf{e}_i - \mathbf{e}_j; 1 \leq i < j \leq n\}; \\ \mathbf{D}_n^+ &= \{\mathbf{e}_i + \mathbf{e}_j; 1 \leq i < j \leq n\} \cup \{\mathbf{e}_i - \mathbf{e}_j; 1 \leq i < j \leq n\}; \\ \mathbf{BC}_n^+ &= \mathbf{B}_n^+ \cup \mathbf{C}_n^+. \end{aligned}$$

Let, in addition, $\tilde{\Phi}^+ = \Phi^+ \cup \{(0, 0, \dots, 0)\}$, where $\Phi = \mathbf{A}_{n-1}, \mathbf{B}_n, \mathbf{C}_n, \mathbf{D}_n$ or \mathbf{BC}_n and where $(0, 0, \dots, 0)$ is the origin of \mathbb{R}^n .

In their combinatorial study of hypergeometric functions associated with root systems, Gelfand, Graev and Postnikov [2, Theorem 6.3] discovered a squarefree quadratic initial ideal of the toric ideal $I_{\tilde{\mathbf{A}}_{n-1}^+}$ of $\tilde{\mathbf{A}}_{n-1}^+$. Moreover, for any subconfiguration \mathcal{A} of \mathbf{A}_{n-1}^+ , the configuration $\tilde{\mathcal{A}} = \mathcal{A} \cup (0, 0, \dots, 0)$ possesses a regular unimodular triangulation ([7, Example 2.4 (a)]). Stanley [8, Exercise 6.31 (b), p. 234] computed the Ehrhart polynomial of the convex polytope $\text{conv}(\tilde{\mathbf{A}}_{n-1}^+)$. Fong [1] constructed certain triangulations of the configurations $\tilde{\mathbf{B}}_n^+ (= \text{conv}(\tilde{\mathbf{D}}_n^+) \cap \mathbb{Z}^n)$

and $\text{conv}(\widetilde{C}_n^+) \cap \mathbb{Z}^n (= \widetilde{BC}_n^+)$, and computes the Ehrhart polynomials of $\text{conv}(\widetilde{B}_n^+)$ and $\text{conv}(\widetilde{C}_n^+)$. The triangulations studied in [1] are, however, non-unimodular. Motivated by their results, Ohsugi–Hibi [6] showed that

Proposition 0.1. *Let $\Phi \subset \mathbb{Z}^n$ be one of the root systems A_{n-1} , B_n , C_n , D_n and BC_n . Then, there exists a reverse lexicographic order such that the initial ideal of $I_{\widetilde{\Phi}^+}$ is generated by squarefree quadratic monomials.*

Moreover, Ohsugi–Hibi [5] discussed subconfigurations $\widetilde{\mathcal{A}} = \mathcal{A} \cup \{(0, 0, \dots, 0)\}$ of $\widetilde{B}_n^+ \cup \widetilde{C}_n^+$ which possesses a (regular) unimodular triangulation (i.e., $I_{\widetilde{\mathcal{A}}}$ which possesses a squarefree initial ideal).

Hence, it is natural to study the same problem as above for I_{Φ^+} where $\Phi \subset \mathbb{Z}^n$ is one of the root systems A_{n-1} , B_n , C_n , D_n and BC_n . (Then, I_{Φ^+} is not generated by quadratic binomials if $n \geq 6$.)

1. SQUAREFREE LEXICOGRAPHIC INITIAL IDEALS

Let $\Phi^+ \subset \mathbb{Z}^n$ denote one of the configurations A_{n-1}^+ , B_n^+ , C_n^+ , D_n^+ and BC_n^+ . Let $K[A_{n-1}^+]$, $K[B_n^+]$, $K[C_n^+]$, $K[D_n^+]$ and $K[BC_n^+]$ denote the polynomial rings

$$\begin{aligned} K[A_{n-1}^+] &= K[\{f_{i,j}\}_{1 \leq i < j \leq n}], \\ K[B_n^+] &= K[\{y_i\}_{1 \leq i \leq n} \cup \{e_{i,j}\}_{1 \leq i < j \leq n} \cup \{f_{i,j}\}_{1 \leq i < j \leq n}], \\ K[C_n^+] &= K[\{a_i\}_{1 \leq i \leq n} \cup \{e_{i,j}\}_{1 \leq i < j \leq n} \cup \{f_{i,j}\}_{1 \leq i < j \leq n}], \\ K[D_n^+] &= K[\{e_{i,j}\}_{1 \leq i < j \leq n} \cup \{f_{i,j}\}_{1 \leq i < j \leq n}], \\ K[BC_n^+] &= K[\{a_i\}_{1 \leq i \leq n} \cup \{y_i\}_{1 \leq i \leq n} \cup \{e_{i,j}\}_{1 \leq i < j \leq n} \cup \{f_{i,j}\}_{1 \leq i < j \leq n}] \end{aligned}$$

over K . Write $\pi : K[\Phi^+] \rightarrow K[t, t^{-1}, s]$ for the homomorphism defined by setting

$$\pi(a_i) = t_i^2 s, \quad \pi(y_i) = t_i s, \quad \pi(e_{i,j}) = t_i t_j s, \quad \pi(f_{i,j}) = t_i t_j^{-1} s.$$

Thus the kernel of π is the toric ideal I_{Φ^+} .

First, an explicit initial ideals of $I_{A_{n-1}^+}$ generated by squarefree monomials of degree ≤ 3 will be constructed. Let $<_{lex}$ be the lexicographic order induced by the ordering of variables

$$f_{n-1,n} > f_{n-2,n-1} > f_{n-2,n} > \cdots > f_{1,2} > f_{1,3} > \cdots > f_{1,n},$$

and let $<_{rev}$ be the reverse lexicographic order induced by the ordering of variables

$$f_{n-1,n} > f_{n-2,n} > f_{n-2,n-1} > \cdots > f_{2,3} > f_{1,n} > \cdots > f_{1,3} > f_{1,2}.$$

Then, the reduced Gröbner basis with respect to $<_{lex}$ (and $<_{rev}$) is as follows.

Theorem 1.1 ([4]). *The set of the binomials*

$$\begin{aligned} f_{i,\ell} f_{j,k} - f_{i,k} f_{j,\ell}, & \quad i < j < k < \ell, \\ f_{i,j} f_{j,k} - f_{i,i+1} f_{i+1,k}, & \quad i+1 < j < k, \\ f_{i,j} f_{k,k+1} f_{k+1,\ell} - f_{i,i+1} f_{i+1,j} f_{k,\ell}, & \quad i+1 < j < k < \ell-1, \end{aligned}$$

is the reduced Gröbner basis of the toric ideal $I_{A_{n-1}^+}$ with respect to both $<_{lex}$ and $<_{rev}$, where the initial monomial of each binomial is the first monomial.

Then, we can associate the initial ideal of $I_{\mathbf{A}_{n-1}^+}$ with respect to \langle_{lex} with the regular unimodular triangulation $\Delta_{\langle_{lex}}$. A graph-theoretical characterization of the maximal faces of the triangulation $\Delta_{\langle_{lex}}$ is given in [4].

Second, we discuss the existence of squarefree initial ideals of the toric ideal I_{Φ^+} where $\Phi \subset \mathbb{Z}^n$ is one of the root systems \mathbf{B}_n , \mathbf{C}_n , \mathbf{D}_n and \mathbf{BC}_n . The similar argument as in [5] plays an important role in the proof of Theorems 1.2 and 1.4.

Let \langle_{lex}^c be the lexicographic order induced by the ordering of variables

$$\begin{aligned} a_1 &> a_2 > \cdots > a_n \\ &> f_{n-1,n} > f_{n-2,n-1} > f_{n-2,n} > \cdots > f_{1,2} > f_{1,3} > \cdots > f_{1,n} \\ &> e_{n-1,n} > e_{n-2,n-1} > e_{n-2,n} > \cdots > e_{1,2} > e_{1,3} > \cdots > e_{1,n}. \end{aligned}$$

Theorem 1.2. *The initial ideal of the toric ideal $I_{\mathbf{C}_n^+}$ with respect to \langle_{lex}^c is generated by squarefree monomials.*

Let \langle_{lex}^d denote the lexicographic order obtained by restricting \langle_{lex}^c to $K[\mathbf{D}_n^+]$. By the elimination property of the lexicographic order \langle_{lex}^c , we have the following corollary from Theorem 1.2.

Corollary 1.3. *The initial ideal of the toric ideal $I_{\mathbf{D}_n^+}$ with respect to \langle_{lex}^d is generated by squarefree monomials.*

We now consider the root systems \mathbf{B}_n and \mathbf{BC}_n . Let \langle_{lex}^{bc} be the lexicographic order induced by the ordering of variables

$$\begin{aligned} a_1 &> a_2 > \cdots > a_n \\ &> e_{n-1,n} > e_{n-2,n-1} > e_{n-2,n} > \cdots > e_{1,2} > e_{1,3} > \cdots > e_{1,n} \\ &> y_1 > y_2 > \cdots > y_n \\ &> f_{n-1,n} > f_{n-2,n-1} > f_{n-2,n} > \cdots > f_{1,2} > f_{1,3} > \cdots > f_{1,n}. \end{aligned}$$

Theorem 1.4. *The initial ideal of the toric ideal $I_{\mathbf{BC}_n^+}$ with respect to \langle_{lex}^{bc} is generated by squarefree monomials.*

Let \langle_{lex}^b denote the lexicographic order obtained by restricting \langle_{lex}^{bc} to $K[\mathbf{B}_n^+]$. By the elimination property of the lexicographic order \langle_{lex}^{bc} , we have the following corollary from Theorem 1.4.

Corollary 1.5. *The initial ideal of the toric ideal $I_{\mathbf{B}_n^+}$ with respect to \langle_{lex}^b is generated by squarefree monomials.*

Remark 1.6. Let $n \geq 6$ and let Φ^+ denote one of the configurations \mathbf{A}_{n-1}^+ , \mathbf{B}_n^+ , \mathbf{C}_n^+ , \mathbf{D}_n^+ and \mathbf{BC}_n^+ . Then I_{Φ^+} is not generated by quadratic binomials. Hence, in particular, I_{Φ^+} does not possess a quadratic Gröbner basis.

REFERENCES

- [1] W. Fong, *Triangulations and Combinatorial Properties of Convex Polytopes*, Dissertation, M. I. T., June, 2000.
- [2] I. M. Gelfand, M. I. Graev and A. Postnikov, *Combinatorics of hypergeometric functions associated with positive roots*, in “Arnold–Gelfand Mathematics Seminars, Geometry and Singularity Theory” (V. I. Arnold, I. M. Gelfand, M. Smirnov and V. S. Retakh, Eds.), Birkhäuser, Boston, 1997, pp. 205 – 221.
- [3] J. E. Humphreys, “*Introduction to Lie Algebras and Representation Theory*,” Second Printing, Revised, Springer–Verlag, Berlin, Heidelberg, New York, 1972.
- [4] T. Kitamura, H. Ohsugi and T. Hibi, *Gröbner bases associated with positive roots and Catalan numbers*, preprint.
- [5] H. Ohsugi and T. Hibi, *Unimodular triangulations and coverings of configurations arising from root systems*, *J. Algebraic Combinatorics*, **14** (2001), 199 – 219.
- [6] H. Ohsugi and T. Hibi, *Quadratic initial ideals of root systems*, *Proc. Amer. Math. Soc.*, **130** (2002), 1913 – 1922.
- [7] R. P. Stanley, *Decompositions of rational convex polytopes*, *Ann. Discrete Math.* **6** (1980), 333 – 342.
- [8] R. P. Stanley, “*Enumerative Combinatorics, Volume II*,” Cambridge University Press, Cambridge, New York, Sydney, 1999.
- [9] B. Sturmfels, “*Gröbner Bases and Convex Polytopes*,” Amer. Math. Soc., Providence, RI, 1995.

Department of Mathematics,
Rikkyo University,
Nishi-Ikebukuro, Tokyo 171–8501, Japan
E-mail: ohsugi@rkmath.rikkyo.ac.jp