

## ON THE LIFTING OF HERMITIAN MODULAR FORMS

TAMOTSU IKEDA

### Notation

Let  $K$  be an imaginary quadratic field with discriminant  $-D = -D_K$ . We denote by  $\mathcal{O} = \mathcal{O}_K$  the ring of integers of  $K$ . The non-trivial automorphism of  $K$  is denoted by  $x \mapsto \bar{x}$ . The primitive Dirichlet character corresponding to  $K/\mathbb{Q}$  is denoted by  $\chi = \chi_D$ . We denote by  $\mathcal{O}^\# = (\sqrt{-D})^{-1}\mathcal{O}$  the inverse different ideal of  $K/\mathbb{Q}$ .

The special unitary group  $G = \mathrm{SU}(m, m)$  is an algebraic group defined over  $\mathbb{Q}$  such that

$$G(R) = \left\{ g \in \mathrm{SL}_{2m}(R \otimes K) \mid g \begin{pmatrix} \mathbf{0}_m & -\mathbf{1}_m \\ \mathbf{1}_m & \mathbf{0}_m \end{pmatrix} {}^t \bar{g} = \begin{pmatrix} \mathbf{0}_m & -\mathbf{1}_m \\ \mathbf{1}_m & \mathbf{0}_m \end{pmatrix} \right\}$$

for any  $\mathbb{Q}$ -algebra  $R$ . We put  $\Gamma_K^{(m)} = G(\mathbb{Q}) \cap \mathrm{GL}_{2m}(\mathcal{O})$ .

The hermitian upper half space  $\mathcal{H}_m$  is defined by

$$\mathcal{H}_m = \{ Z \in M_m(\mathbb{C}) \mid \frac{1}{2\sqrt{-1}}(Z - {}^t \bar{Z}) > 0 \}.$$

Then  $G(\mathbb{R})$  acts on  $\mathcal{H}_m$  by

$$g \langle Z \rangle = (AZ + B)(CZ + D)^{-1}, \quad Z \in \mathcal{H}_m, g = \begin{pmatrix} A & B \\ C & D \end{pmatrix}.$$

We put

$$\begin{aligned} \Lambda_m(\mathcal{O}) &= \{ h = (h_{ij}) \in M_m(K) \mid h_{ii} \in \mathbb{Z}, h_{ij} = \bar{h}_{ji} \in \mathcal{O}^\#, i \neq j \}, \\ \Lambda_m(\mathcal{O})^+ &= \{ h \in \Lambda_m(\mathcal{O}) \mid h > 0 \}. \end{aligned}$$

We set  $e(T) = \exp(2\pi\sqrt{-1}\mathrm{tr}(T))$  if  $T$  is a square matrix with entries in  $\mathbb{C}$ . For each prime  $p$ , the unique additive character of  $\mathbb{Q}_p$  such that  $e_p(x) = \exp(-2\pi\sqrt{-1}x)$  for  $x \in \mathbb{Z}[p^{-1}]$  is denoted  $e_p$ . Note that  $e_p$  is of order 0. We put  $e(x) = e(x_\infty) \prod_{p < \infty} e_p(x_p)$  for an adèle  $x = (x_v)_v \in \mathbb{A}$ .

Let  $\underline{\chi} = \otimes_v \underline{\chi}_v$  be the Hecke character of  $\mathbb{A}^\times / \mathbb{Q}^\times$  determined by  $\chi$ . Then  $\underline{\chi}_v$  is the character corresponding to  $\mathbb{Q}_v(\sqrt{-D})/\mathbb{Q}$  and given by

$$\underline{\chi}_v(t) = \left( \frac{-D, t}{\mathbb{Q}_v} \right).$$

Let  $Q_D$  be the set of all primes which divides  $D$ . For each prime  $q \in Q_D$ , we put  $D_q = q^{\text{ord}_q D}$ . We define a primitive Dirichlet character  $\chi_q$  by

$$\chi_q(n) = \begin{cases} \chi(n') & \text{if } (n, q) = 1 \\ 0 & \text{if } q|n, \end{cases}$$

where  $n'$  is an integer such that

$$n' \equiv \begin{cases} n & \text{mod } D_q, \\ 1 & \text{mod } D_q^{-1}D \end{cases}$$

Then we have  $\chi = \prod_{q|D} \chi_q$ . Note that

$$\chi_q(n) = \left( \frac{\chi_q(-1)D_q, n}{\mathbb{Q}_q} \right) = \prod_{p|n} \left( \frac{\chi_q(-1)D_q, n}{\mathbb{Q}_p} \right)$$

for  $q \nmid n$ ,  $n > 0$ . One should not confuse  $\chi_q$  with  $\underline{\chi}_q$ .

### 1. Fourier coefficients of Eisenstein series on $\mathcal{H}_m$

In this section, we consider Siegel series associated to non-degenerate hermitian matrices. Fix a prime  $p$ . Put  $\xi_p = \chi(p)$ , i.e.,

$$\xi_p = \begin{cases} 1 & \text{if } -D \in (\mathbb{Q}_p^\times)^2 \\ -1 & \text{if } \mathbb{Q}_p(\sqrt{-D})/\mathbb{Q}_p \text{ is unramified quadratic extension} \\ 0 & \text{if } \mathbb{Q}_p(\sqrt{-D})/\mathbb{Q}_p \text{ is ramified quadratic extension.} \end{cases}$$

For  $H \in \Lambda_m(\mathcal{O})$ ,  $\det H \neq 0$ , we put

$$\begin{aligned} \gamma(H) &= (-D)^{\lfloor m/2 \rfloor} \det(H) \\ \zeta_p(H) &= \underline{\chi}_p(\gamma(H))^{m-1}. \end{aligned}$$

The Siegel series for  $H$  is defined by

$$b_p(H, s) = \sum_{R \in \text{Her}_m(K_p)/\text{Her}_m(\mathcal{O}_p)} e_p(\text{tr}(BR)) p^{-\text{ord}_p(\nu(R))s}, \quad \text{Re}(s) \gg 0.$$

Here,  $\text{Her}_m(K_p)$  (resp.  $\text{Her}_m(\mathcal{O}_p)$ ) is the additive group of all hermitian matrices with entries in  $K_p$  (resp.  $\mathcal{O}_p$ ). The ideal  $\nu(R) \subset \mathbb{Z}_p$  is defined

as follows: Choose a coprime pair  $\{C, D\}$ ,  $C, D \in M_{2n}(\mathcal{O}_p)$  such that  $C^t \bar{D} = D^t \bar{C}$ , and  $D^{-1}C = R$ . Then  $\nu(R) = \det(D)\mathcal{O}_p \cap \mathbb{Z}_p$ .

We define a polynomial  $t_p(K/\mathbb{Q}; X) \in \mathbb{Z}[X]$  by

$$t_p(K/\mathbb{Q}; X) = \prod_{i=1}^{\lfloor (m+1)/2 \rfloor} (1 - p^{2i} X) \prod_{i=1}^{\lfloor m/2 \rfloor} (1 - p^{2i-1} \xi_p X).$$

There exists a polynomial  $F_p(H; X) \in \mathbb{Z}[X]$  such that

$$F_p(H; p^{-s}) = b_p(H, s) t_p(K/\mathbb{Q}; p^{-s})^{-1}.$$

This is proved in [9].

Moreover,  $F_p(H; X)$  satisfies the following functional equation:

$$F_p(H; p^{-2m} X^{-1}) = \zeta_p(H) (p^m X)^{-\text{ord}_p \gamma(H)} F_p(H; X).$$

This functional equation is a consequence of [7], Proposition 3.1. We will discuss it in the next section.

The functional equation implies that  $\deg F_p(H; X) = \text{ord}_p \gamma(H)$ . In particular, if  $p \nmid \gamma(H)$ , then  $F_p(H; X) = 1$ . Put

$$\tilde{F}_p(H; X) = X^{-\text{ord}_p \gamma(H)} F_p(H; p^{-m} X^2).$$

Then following lemma is a immediate consequence of the functional equation of  $F(H; X)$ .

**Lemma 1.** *We have*

$$\tilde{F}_p(H; X^{-1}) = \tilde{F}_p(H; X), \quad \text{if } m \text{ is odd.}$$

$$\tilde{F}_p(H; \xi_p X^{-1}) = \tilde{F}_p(H; X), \quad \text{if } m \text{ is even and } \xi_p \neq 0.$$

Let  $k$  be a sufficiently large integer. Put  $n = \lfloor m/2 \rfloor$ . The Eisenstein series  $E_{2k+2n}^{(m)}(Z)$  of weight  $2k + 2n$  on  $\mathcal{H}_m$  is defined by

$$E_{2k+2n}^{(m)}(Z) = \sum_{\{C, D\}/\sim} \det(CZ + D)^{-2k-2n},$$

where  $\{C, D\}/\sim$  extends over coprime pairs  $\{C, D\}$ ,  $C, D \in M_{2n}(\mathcal{O})$  such that  $C^t \bar{D} = D^t \bar{C}$  modulo the action of  $GL_m(\mathcal{O})$ . We put

$$\mathcal{E}_{2k+2n}^{(m)}(Z) = A_m^{-1} \prod_{i=1}^m L(1 + i - 2k - 2n, \chi^{i-1}) E_{2k+2n}^{(m)}(Z).$$

Here

$$A_m = \begin{cases} 2^{-4n^2-4n} \mathbf{D}^{2n^2+n} & \text{if } m = 2n + 1, \\ (-1)^n 2^{-4n^2+4n} \mathbf{D}^{2n^2-n} & \text{if } m = 2n. \end{cases}$$

TAMOTSU IKEDA

Then the  $H$ -th Fourier coefficient of  $\mathcal{E}_{2k+2n}^{(2n+1)}(Z)$  is equal to

$$\begin{aligned} |\gamma(H)|^{2k-1} \prod_{p|\gamma(H)} F_p(H; p^{-2k-2n}) &= |\gamma(H)|^{k-(1/2)} \prod_{p|\gamma(H)} \tilde{F}_p(H; p^{-k+(1/2)}) \\ &= |\gamma(H)|^{k-(1/2)} \prod_{p|\gamma(H)} \tilde{F}_p(H; p^{k-(1/2)}) \end{aligned}$$

for any  $H \in \Lambda_{2n+1}(\mathcal{O})^+$  and any sufficiently large integer  $k$ .

The  $H$ -th Fourier coefficient of  $\mathcal{E}_{2k+2n}^{(2n)}(Z)$  is equal to

$$|\gamma(H)|^{2k} \prod_{p|\gamma(H)} F_p(H; p^{-2k-2n}) = |\gamma(H)|^k \prod_{p|\gamma(H)} \tilde{F}_p(H; p^{-k})$$

for any  $H \in \Lambda_{2n}(\mathcal{O})^1$  and any sufficiently large integer  $k$ .

## 2. Main theorems

We first consider the case when  $m = 2n$  is even.

Let  $f(\tau) = \sum_{N=1}^{\infty} a(N)q^N \in S_{2k+1}(\Gamma_0(\mathbf{D}), \chi)$  be a primitive form, whose  $L$ -function is given by

$$\begin{aligned} L(f, s) &= \sum_{N=1}^{\infty} a(N)N^{-s} \\ &= \prod_{p \mid \mathbf{D}} (1 - a(p)p^{-s} + \chi(p)p^{2k-2s})^{-1} \prod_{q \nmid \mathbf{D}} (1 - a(q)q^{-s})^{-1}. \end{aligned}$$

For each prime  $p \nmid \mathbf{D}$ , we define the Satake parameter  $\{\alpha_p, \beta_p\} = \{\alpha_p, \chi(p)\alpha_p^{-1}\}$  by

$$(1 - a(p)X + \chi(p)p^{2k}X^2) = (1 - p^k\alpha_p X)(1 - p^k\beta_p X).$$

For  $q \mid \mathbf{D}$ , we put  $\alpha_q = q^{-k}a(q)$ .

Put

$$A(H) = |\gamma(H)|^k \prod_{p|\gamma(H)} \tilde{F}_p(H, \alpha_p), \quad H \in \Lambda_{2n}(\mathcal{O})^+$$

$$F(Z) = \sum_{H \in \Lambda_{2n}(\mathcal{O})^+} A(H)\mathbf{e}(HZ), \quad Z \in \mathcal{H}_{2n}.$$

Then our first main theorem is as follows:

**Theorem 1.** *Assume that  $m = 2n$  is even. Let  $f(\tau)$ ,  $A(H)$  and  $F(Z)$  be as above. Then we have  $F \in S_{2k+2n}(\Gamma_K^{(2n)})$ . Moreover,  $F$  is a Hecke eigenform.  $F = 0$  if and only if  $f(\tau)$  comes from a Hecke character of  $K$  and  $n$  is odd.*

Now we consider the case when  $m = 2n + 1$  is odd.

Let  $f(\tau) = \sum_{N=1}^{\infty} a(N)q^N \in S_{2k}(\mathrm{SL}_2(\mathbb{Z}))$  be a normalized Hecke eigenform, whose  $L$ -function is given by

$$\begin{aligned} L(f, s) &= \sum_{N=1}^{\infty} a(N)N^{-s} \\ &= \prod_p (1 - a(p)p^{-s} + p^{2k-1-2s})^{-1} \end{aligned}$$

For each prime  $p$ , we define the Satake parameter  $\{\alpha_p, \alpha_p^{-1}\}$  by

$$(1 - a(p)X + p^{2k-1}X^2) = (1 - p^{k-(1/2)}\alpha_p X)(1 - p^{k-(1/2)}\alpha_p^{-1}X).$$

Put

$$A(H) = |\gamma(H)|^{k-(1/2)} \prod_{p|\gamma(H)} \tilde{F}_p(H, \alpha_p), \quad H \in \Lambda_{2n+1}(\mathcal{O})^+$$

$$F(Z) = \sum_{H \in \Lambda_{2n+1}(\mathcal{O})^+} A(H)e(HZ), \quad Z \in \mathcal{H}_{2n+1}.$$

**Theorem 2.** *Assume that  $m = 2n + 1$  is odd. Let  $f(\tau)$ ,  $A(H)$  and  $F(Z)$  be as above. Then we have  $F \in S_{2k+2n}(\Gamma_K^{(2n+1)})$ . Moreover,  $F$  is a non-zero Hecke eigenform.*

## REFERENCES

- [1] J. Arthur, *Unipotent automorphic representations: conjectures*, *Astérisque* **171-172** (1989), 13–71.
- [2] S. Böcherer, *Über die Fourier-Jacobi-Entwicklung Siegelscher Eisensteinreihen. I*, *Math. Z.* **183** (1983), 21–46.
- [3] S. Breulmann and M. Kuss, *On a conjecture of Duke-Imamoglu*, *Proc. Amer. Math. Soc.* **107** (2000),
- [4] M. Eichler and D. Zagier, *The theory of Jacobi forms*, *Progress in Mathematics* **55** Birkhäuser Boston, Inc., Boston, Mass. 1985.
- [5] T. Ikeda, *On the theory of Jacobi forms and the Fourier-Jacobi coefficients of Eisenstein series*, *J. Math. Kyoto Univ.* **34** (1994), 615–636.
- [6] ———, *On the lifting of elliptic cusp forms to Siegel cusp forms of degree  $2n$* , *Ann. of Math.* **154** (2001), 641–681.
- [7] S. Kudla and W. J. Sweet Jr. *Degenerate principal series representations for  $U(n, n)$* . *Israel J. Math.* **98** (1997), 253–306.
- [8] G. Shimura *Introduction to the arithmetic theory of automorphic functions*, *Publ. Math. Soc. Japan* **11** Iwanami Shoten and Princeton University Press, 1971.
- [9] ———, *Euler products and Eisenstein series*, *CBMS Regional Conference Series in Mathematics* **93** the American Mathematical Society, Providence, RI, 1997.