

Bivariate Chebyshev maps of \mathbf{C}^2 and their dynamics

東海大学・理学部 内村桂輔 (Keisuke Uchimura)
Department of Mathematics
Tokai University

Abstract

We study the properties of bivariate (two-dimensional) Chebyshev maps $T_n(x, y)$ from \mathbf{C}^2 to \mathbf{C}^2 and study the properties and dynamics of the maps.

(A) The properties of T_n .

- (1) Solutions of $T_n(x, y) = (0, 0)$ are obtained.
- (2) A critical set $\det(DT_n) = 0$ is written in a simple formula.

These properties are similar to those of Chebyshev maps of \mathbf{C} .

(B) The dynamics of T_n .

- (1) T_n is strictly critically finite.
- (2) Any periodic point of T_n is repelling.
- (3) The exact form of the invariant probability measure μ of maximal entropy associated with T_n is obtained.
- (4) External rays for $J_2(T_n)$ and foliations of $J_1(T_n)$ are studied.

These properties are also similar to those of Chebyshev maps of \mathbf{C} .

1 Bivariate Chebyshev maps

The Chebyshev map is a typical chaotic map. Generalized Chebyshev maps are studied by several researchers, Koornwinder [1974], Lidle [1975], Veselov [1987] and Hoffman & Withers [1988].

In this paper, we study bivariate Chebyshev maps T_n from \mathbf{C}^2 to \mathbf{C}^2 , $n \in \mathbf{Z}$.

$$T_n(x, y) = (g^{(n)}(x, y), g^{(n)}(y, x)).$$

This definition is due to [V]. Here $g^{(n)}(x, y)$ is a generalized Chebyshev polynomial defined by Lidle [L].

Let

$$x = t_1 + t_2 + t_3, \quad y = t_1t_2 + t_1t_3 + t_2t_3, \quad 1 = t_1t_2t_3.$$

Then

$$g^{(n)}(x, y) := t_1^n + t_2^n + t_3^n.$$

So

$$g^{(n)}(y, x) = (1/t_1)^n + (1/t_2)^n + (1/t_3)^n = g^{(-n)}(x, y).$$

For instance,

$$T_2(x, y) = (x^2 - 2y, y^2 - 2x),$$

$$T_3(x, y) = (x^3 - 3xy + 3, y^3 - 3xy + 3),$$

$$T_4(x, y) = (x^4 - 4x^2y + 2y^2 + 4x, y^4 - 4xy^2 + 2x^2 + 4y).$$

$\{g^n(x, y)\}$ satisfy the following recurrence equation:

$$g^{(n)}(x, y) = xg^{(n-1)}(x, y) - yg^{(n-2)}(x, y) + g^{(n-3)}(x, y).$$

First, we show a branch covering over \mathbb{C}^2 .

The following diagram is commutative.

$$\begin{array}{ccc} (\mathbb{C} - \{0\})^2 & \xrightarrow{g^n} & (\mathbb{C} - \{0\})^2 \\ \downarrow \Psi & & \downarrow \Psi \\ \mathbb{C}^2 & \xrightarrow{T_n} & \mathbb{C}^2 \end{array}$$

$$\text{where} \quad g_n(u, v) = (u^n, v^n),$$

and

$$(x, y) = \Psi(u, v) = \left(u + v + \frac{1}{uv}, \frac{1}{u} + \frac{1}{v} + uv\right).$$

The covering map

$$\Psi : \mathbb{C}^2 - \Psi^{-1}(D) \rightarrow \mathbb{C}^2 - D$$

is a 6-sheeted covering map. Branch locus D of Ψ is written as

$$x^2y^2 - 4x^3 - 4y^3 + 18xy - 27 = 0.$$

In the case $n = 2$, Ueda[*Ue*] showed this diagram.

$T_n(x, y)$ restricted on $\{x = \bar{y}\}$ is a Chebyshev polynomial defined by Koornwinder [K]

$$P_{n,0}^{-\frac{1}{2}}(z, \bar{z}) = e^{in\sigma} + e^{-in\tau} + e^{i(n\tau - n\sigma)}.$$

Set

$$z(\sigma, \tau) := e^{i\sigma} + e^{-i\tau} + e^{i(\tau - \sigma)} = u + iv.$$

The mapping

$$z : (\sigma, \tau) \rightarrow (u, v)$$

is a diffeomorphism from R onto S . See Koornwinder [K].

Proposition 1. *There are n^2 solutions of $T_n(x, y) = (0, 0)$. All solutions lie in the closed domain S in $\{x = \bar{y}\}$. They are written in the (σ, τ) coordinate.*

$$(1) \quad (\sigma, \tau) = \left(\frac{2(1+j+h)\pi}{3n}, \frac{2(1+2j+h)\pi}{3n} \right)$$

$$j = 0, 1, \dots, n-1, \text{ and } h = 0, 1, \dots, j.$$

$$(2) \quad (\sigma, \tau) = \left(\frac{2(2+j+h)\pi}{3n}, \frac{2(2+2j-h)\pi}{3n} \right)$$

$$j = 0, 1, \dots, n-2, \text{ and } h = 0, 1, \dots, j.$$

Proof. By definition,

$$T_n(x, y) = (g^{(n)}(x, y), g^{(n)}(y, x)).$$

$g^{(n)}(x, y)$ and $g^{(n)}(y, x)$ are polynomials of degree n with no common components.

We can find n^2 zeros on S . See Uchimura [Uc1]. \square

We see that the zeros of T_n and T_{n+1} "mutually separate each other".

Next we consider critical set of $T_n(x, y)$.

$$C_n := \{(x, y) \in \mathbb{C}^2 : \det(DT_n) = 0\}.$$

Proposition 2. *Let $n \in \mathbb{Z}$. Assume that*

$$x = t_1 + t_2 + t_3, \quad y = t_1 t_2 + t_1 t_3 + t_2 t_3, \quad t_1 t_2 t_3 = 1.$$

Then

$$\text{Det}(DT_n) = n^2 \frac{t_1^n - t_2^n}{t_1 - t_2} \cdot \frac{t_1^n - t_3^n}{t_1 - t_3} \cdot \frac{t_2^n - t_3^n}{t_2 - t_3}.$$

Proof.

$$\text{Det}(DT_n) = \text{Det}(D(T_n \circ \Psi)) / \text{Det}(D\Psi). \quad \square$$

The similar result is holds for generalized Chebyshev maps from \mathbb{C}^n to \mathbb{C}^n .

Corollary 1. *Any irreducible component of C_n is a rational curve of degree 2 or 4.*

Proof. From Proposition 2, we have

$$x = t + \epsilon^k t + \frac{1}{\epsilon^k t^2}, \quad \epsilon = e^{\frac{2\pi i}{n}}$$

$$y = \frac{1}{t} + \frac{1}{\epsilon^k t} + \epsilon^k t^2.$$

When $\epsilon^k = -1$, the degree of the rational curve is 2. □

We see that C_n and C_{n+1} "mutually separate each other", and

$$C_n \cap S \neq \emptyset \quad (S = J_2(T_n)).$$

Note that $\{T_m : m \in \mathbf{Z}\}$ is a semigroup satisfying

$$T_m \circ T_n = T_{mn}.$$

2 Dynamics of Bivariate Maps

We study the dynamics of $T_n(x, y)$. Let

$$K(T_n) := \{(x, y) : \{T_n^m(x, y)\} \text{ is bounded for any } m\}.$$

In our setting we have the following proposition.

Proposition 3.

$$K(T_n) = \{|t_1| = |t_2| = 1\} = S \subset \{x = \bar{y}\}.$$

Proof

$$\begin{array}{ccc} (t_1, t_2) & \xrightarrow{g_n} & (t_1^n, t_2^n) \\ \downarrow \Psi & & \downarrow \Psi \\ (x, y) & \xrightarrow{T_n} & (g^{(n)}, g^{(-n)}) \end{array}$$

□

f is called *critically finite* if each irreducible component of the critical set of f is periodic or preperiodic. Dihn and Sibony [DS] show that generalized Chebyshev maps are critically finite. Here using proposition 2, we give a direct proof.

Proposition 4. T_n is strictly critically finite.

Proof.

$$\begin{array}{ccccc} C_n & \xrightarrow{T_n} & T_n(C_n) & \xrightarrow{T_n} & T_n(C_n) \\ (t, \epsilon t) & & (t^n, t^n) & & (t^{n^2}, t^{n^2}) \end{array} \quad \square$$

Next we study the second Julia set J_2 of $T_n(x, y)$.

Proposition 5. All periodic points of T_n lie on S and are equidistributed in S .

Proof. From [FS], we know that number of periodic points with period k equals n^{2k} . For the distribution of periodic points, see [Uc2]. \square

Proposition 6. *Any periodic point of T_n is repelling.*

To prove this proposition we consider the following function.

$$S_n := T_n | \{x = \bar{y}\} : \mathbf{R}^2 \rightarrow \mathbf{R}^2$$

$$\text{e.g. } S_2(z) = z^2 - 2\bar{z} : (u, v) \mapsto (u^2 - 2u - v^2, 2uv + 2v).$$

Lemma 1. *Let p be a periodic point of S_n . Let α and β be eigen values of $DS_n(p)$. Then*

$$|\alpha|, |\beta| > 1.$$

Proposition 7. *Let*

$$f(x, y) \in \mathbf{R}[x, y].$$

$$T(x, y) := (f(x, y), f(y, x)) : \mathbf{C}^2 \rightarrow \mathbf{C}^2.$$

$$t(z) := T | \{x = \bar{y}\} : \mathbf{R}^2 \rightarrow \mathbf{R}^2.$$

Then

$$U^{-1}DT(z, \bar{z})U = Dt(z),$$

where

$$U = \frac{1}{2} \begin{pmatrix} 1+i & -1+i \\ 1+i & 1-i \end{pmatrix}.$$

From Lemma 1 and Proposition 7, Proposition 6 follows. \square

Next we study the invariant measure μ of maximal entropy for T_n .

Proposition 8. *Under the above notation,*

$$\text{supp } \mu = S.$$

$$\mu = \left(\frac{2}{\pi}\right)^2 \frac{dx_1 dx_2}{\sqrt{-x^2 \bar{x}^2 + 4x^3 + 4\bar{x}^3 - 18x\bar{x} + 27}}.$$

$(x = x_1 + ix_2)$

This is an extension of invariant measure

$$\mu = \frac{1}{\pi} \frac{dx}{\sqrt{(x+1)(3-x)}}$$

for Chebyshev maps in one variable on $[-1, 3]$.

Proof. We prove this proposition in the following three steps.

(1) Briend and Duval [BD] shows that

$$\text{let } \mu_n := \frac{1}{d^{nk}} \sum_{f^n(y)=y, y \text{ repelling}} \delta_y,$$

then

$$\mu_n \rightarrow \mu \quad (\text{weak convergence}).$$

(2) From Proposition 5, we see that the periodic points are repelling and equidistributed in the triangle on the (s,t) plane (see [Uc2]).

(3) Pullback of Lebesgue measure under ϕ . □

Next we consider the properties of external rays of $T_n(x, y)$. We use the definitions of external rays by Bedford and Jonsson [BJ]. We extend the map

$$T_n(x, y) : \mathbb{C}^2 \rightarrow \mathbb{C}^2$$

$$\text{to } \hat{T}_n(x : y : z) : \mathbb{P}^2 \rightarrow \mathbb{P}^2.$$

Let $\Pi := \mathbb{P}^2 - \mathbb{C}^2$ be the line at infinity.

Then

$$\hat{T}_n | \Pi : (x : y : 0) \rightarrow (x^n : y^n : 0).$$

Therefore

$$J_\Pi = \{(x : y : 0) : |x| = |y|\} \simeq S^1.$$

The stable set of J_Π for T_n is defined by

$$W^s(J_\Pi, T_n) := \{x \in \mathbb{P}^2 : d(T_n^j x, J_\Pi) \rightarrow 0, \quad j \rightarrow \infty\}.$$

Bedford and Jonsson [BJ] state that there exists a Böttcher coordinate Ψ such that

$$\Psi : W^s(J_\Pi, f_n) \rightarrow W^s(J_\Pi, T_n)$$

conjugating f_n to T_n , where

$$f_n(x, y) = (x^n, y^n).$$

They also show that $W^s(J_\Pi, T_n)$ is foliated by stable disks W_a . They define a local stable manifold $W_{loc}^s(a)$, ($a \in J_\Pi$) and then a stable disk $W_a \supset W_{loc}^s(a)$ and an external ray $R(a, \theta)$. They show that $J_0(T_n) = J_1(T_n)$ is laminated by stable disks W_a .

Nakane [N] shows the following results on $T_2(x, y)$:

(1) The map Ψ defined by Ueda is essentially the inverse of Böttcher coordinate ϕ .

$$\Psi(u, v) = \Psi(t, at), |t| > 1.$$

(2) The stable disk W_a is the set of points $R(r, \phi, \theta)$

$$x = re^{-2\pi i\theta} + \frac{1}{r}e^{2\pi i(\theta-\phi)} + e^{2\pi i\phi},$$

$$y = re^{2\pi i(\phi-\theta)} + \frac{1}{r}e^{2\pi i\theta} + e^{-2\pi i\phi}, \quad a = e^{2\pi i\phi}, \quad (r > 1).$$

An external ray is written as

$$R(\phi, \theta) := \{R(r, \phi, \theta) : r > 1\}.$$

From this,

$$J_2 = S \subset \{x = \bar{y}\}.$$

(3) Each point $z \in S$ is the landing point of exactly 1, 3, or 6 external rays if z is a cusp point on ∂S , z is non-cusp point on ∂S or $z \in \text{int}(S)$ respectively.

We can show that Nakane's results are also true for any $T_n(x, y)$, $n \neq 0$.

Next we study the structure of foliations W_a of

$$J_1(T_n) = W^s(J_\Pi, T_n).$$

Proposition 9. *For any point $z \in \text{int}(S)$, there exist three stable disks W_a such that boundaries of these three disks intersect at z . At the point, two external rays on each W_a land from opposite directions.*

Metaphorically speaking, three mouths (stable disks) eat a sandwich (the second Julia set S).

Two external rays $R(\phi, \theta)$ and $R(\phi, \phi - \theta)$ lie on the stable disk

$$W_a \quad (a = e^{2\pi i\phi}).$$

Two points $R(r, \phi, \theta)$ and $R(r, \phi, \phi - \theta)$ are "symmetrical" about $\{x = \bar{y}\}$ in the following sense.

- (1) The midpoint of the segment $\overline{R(r, \phi, \theta)R(r, \phi, \phi - \theta)}$ lies on the plane $\{x = \bar{y}\}$,
- (2) The segment connecting two points is perpendicular to $\{x = \bar{y}\}$.

We compare the external rays of $T_n(x, y)$ with those of Chebyshev map $T_n(z)$ in one variable. The external rays $T_n(z)$ is written as

$$R(r, \phi) : u = re^{2\pi i\phi} + \frac{1}{r}e^{2\pi i(-\phi)}, \quad (r > 1).$$

Clearly,

$$R(r, -\phi) : v = re^{2\pi i(-\phi)} + \frac{1}{r}e^{2\pi i\phi},$$

$$v = \bar{u}.$$

It is well-known that $R(r, \phi)$ and $R(r, -\phi)$ are "symmetrical" about the real axis. Note that symmetric group S_2 acts on external rays of $T_n(z)$. On the other hand, S_3 acts on external rays of $T_n(x, y)$.

Using the notations in Sect. 1, we can write

$$W^s(J_{\Pi}, T_n) = \{\Psi(t_1, t_2) : |t_1| = \frac{1}{|t_2|} > 1\}.$$

Then

$$C_n \cap W^s(J_{\Pi}, T_n) = \phi.$$

Lastly we consider periodic rays $R(\phi, \theta)$ of $T_n(x, y)$.

Proposition 10. *If one periodic ray lands at the point $z_0 \in S$, all rays which land at z_0 are all periodic with the same period.*

References

- [BJ] E. Bedford and M. Jonsson, *Dynamics of regular polynomial endomorphisms of \mathbf{C}^k* , Amer. J. Math. 122 (2000), 153-212.
- [BD] J. -Y. Briend and J. Duval, *Exposants de Liapounoff et distribution des points periodiques d'un endomorphisme de \mathbf{CP}^k* , Acta Math. 182 (1999), 143-157.
- [DS] T. Dihn and N. Sibony, *Sur les endomorphismes holomorphes permutables*

- de \mathbf{P}^k , Math. Ann. 324 (2002), 33-70.
- [FS] J. Fornæss and N. Sibony, *Complex dynamics in higher dimension I*, Asterisque 222 (1994), 201-231.
- [HW] M. E. Hoffman and W. D. Withers, *Generalized Chebyshev polynomials associated with affine Weyle groups*, Trans. Amer. Math. 308, 1 (1988), 91-104.
- [K] T. H. Koornwinder, *Orthogonal polynomials in two variables which are eigenfunctions of two algebraically independent partial differential Operator, III, IV*, Indag. Math. 38 (1974), 357-381.
- [L] R. Lidl, *Tschebysheff polynome in mehreren variablen*, J. Reine Angew. Math. 273 (1975), 178-198.
- [N] S. Nakane, *External rays for a regular polynomial endomorphism of \mathbf{C}^2 associated with Chebyshev mappings*, preprint (2004).
- [Uc1] K. Uchimura, *Zeros of Chebyshev polynomials in two variables z and \bar{z}* , Rend. Matematica VII, 8 (1988), 187-209.
- [Uc2] K. Uchimura, *The dynamical systems associatted with Chebyshev polynomials in two variables*, Int. J. Bifurcation and Chaos 12B, (6) (1996), 2611-2618.
- [Uc3] K. Uchimura, *The sets of points with bounded orbits for generalized Chebyshev mappings*, Int. J. Bifurcation and Chaos 11, (1) (2001), 91-107.
- [Uc4] K. Uchimura, *Dynamics of Symmetric polynomial endomorphism of \mathbf{C}^2* , preprint (2005).
- [Ue] T. Ueda, *On critically finite maps on the complex projective space*, RIMS Kokyuroku 1087 (1999), 132-138.
- [V] A. P. Veselov, *Integrable mappings and Lie Algebras*, Dokl. Akad. Nauk. SSSR 292 (1987), 1289-1291.