

# Topological Radon Transforms and Projective Duality

Yutaka Matsui (松井 優)

Graduate School of Mathematical Sciences, the University of Tokyo  
(東京大学大学院数理科学研究科)<sup>1</sup>

Kiyoshi Takeuchi (竹内 潔)

Institute of Mathematics, University of Tsukuba  
(筑波大学数学系)<sup>2</sup>

30, November, 2005

## Abstract

We study various topological properties of projective duality in algebraic geometry by using the microlocal theory of sheaves developed by Kashiwara-Schapira [21]. In particular, in the real algebraic case we obtain some results similar to Ernström's ones [9] obtained in the complex case. For this purpose, we use constructible functions and their topological Radon transforms. We also generalize a class formula (i.e. a formula which expresses the degrees of dual varieties) in [10] to the case of associated varieties studied by Gelfand-Kapranov-Zelevinsky [12] etc. For the detail, see [26] and [27].

## 1 Introduction

We denote the projective space of dimension  $n$  over  $\mathbb{K}$  ( $= \mathbb{R}$  or  $\mathbb{C}$ ) by  $\mathbb{P}_n$  and its dual space by  $\mathbb{P}_n^*$ . These spaces are naturally identified with the following sets.

$$\mathbb{P}_n = \{l \mid l \text{ is a line in } \mathbb{K}^{n+1} \text{ through the origin}\}, \quad (1.1)$$

$$\mathbb{P}_n^* = \{H' \mid H' \text{ is a hyperplane in } \mathbb{K}^{n+1} \text{ through the origin}\}. \quad (1.2)$$

Note that if we projectivize a hyperplane  $H'$  in  $\mathbb{K}^{n+1}$  we obtain a hyperplane  $H$  in  $\mathbb{P}_n$ . Therefore in what follows we identify the dual projective space  $\mathbb{P}_n^*$  with the set

$$\{H \mid H \text{ is a hyperplane in } \mathbb{P}_n\}. \quad (1.3)$$

---

<sup>1</sup>3-8-1, Komaba, Meguro-ku, Tokyo, 153-8914, Japan, E-mail: you317@ms.u-tokyo.ac.jp

<sup>2</sup>1-1-1, Tennodai, Tsukuba, Ibaraki, 305-8571, Japan, E-mail: takemicro@nifty.com

**Definition 1.1** Let  $V$  be a projective variety in  $\mathbb{P}_n$ . We define the dual variety  $V^*$  of  $V$  by

$$V^* := \overline{\{H \in \mathbb{P}_n^* \mid \exists x \in V_{\text{reg}} \cap H \text{ s.t. } T_x V \subset T_x H\}} \subset \mathbb{P}_n^*. \quad (1.4)$$

When  $V$  is smooth,  $V^*$  is the set of hyperplanes tangent to  $V$ . As we see in the example below, even if  $V$  is smooth,  $V^*$  may be very singular in general.

**Example 1.2** (i) Let  $\iota_n : \mathbb{P}_1 \hookrightarrow \mathbb{P}_n$  be the Veronese embedding given by  $[x : y] \mapsto [x^n : x^{n-1}y : \dots : xy^{n-1} : y^n]$  and set  $V = \iota_n(\mathbb{P}_1) \subset \mathbb{P}_n$ . Then the dual  $V^* \subset \mathbb{P}_n^*$  is a hypersurface defined by the classical discriminant for polynomials of degree  $n$ .

(ii) For  $n \geq m$ , consider the Segre embedding  $\iota_{n,m} : \mathbb{P}_n \times \mathbb{P}_m \hookrightarrow \mathbb{P}_{(n+1)(m+1)-1}$  given by  $([x_0 : \dots : x_n], [y_0 : \dots : y_m]) \mapsto [\dots : x_i y_j : \dots]$ . Set  $W = \iota_{n,m}(\mathbb{P}_n \times \mathbb{P}_m) \subset \mathbb{P}_{(n+1)(m+1)-1}$ . Then the dual variety  $W^* \subset \mathbb{P}_{(n+1)(m+1)-1}^*$  has very complicated singularities and the dual defect  $\delta^*(W)$  of  $W$  (see (2.2) below) is  $n - m$ . Indeed, let  $M_{(n+1),(m+1)}$  be the space of  $(n+1) \times (m+1)$  matrices and identify the dual projective space  $\mathbb{P}_{(n+1)(m+1)-1}^*$  with its projectivization  $\mathbb{P}(M_{(n+1),(m+1)})$ . Then the dual  $W^* \subset \mathbb{P}_{(n+1)(m+1)-1}^*$  is explicitly written by

$$W^* = \mathbb{P}(\{A \in M_{(n+1),(m+1)} \mid \text{rank} A \leq m\}). \quad (1.5)$$

Therefore the dual  $W^*$  admits a stratification defined by the ranks of matrices.

Many mathematicians were interested in the mysterious relations between projective varieties and their duals. Above all, they observed that the tangency of a hyperplane  $H \in V^*$  with  $V$  is related to the singularity of the dual  $V^*$  at  $H$ . For example, consider the case of a plane curve  $C \subset \mathbb{P}_2$  over  $\mathbb{C}$ . Then a tangent line  $l$  at an inflection point of  $C$  corresponds to a cusp of the dual curve  $C^*$ , and a bitangent (double tangent) line  $l$  of  $C$  corresponds to an ordinary double point of  $C^*$ . The most general results for complex plane curves were found in the 19th century by Klein, Plücker and Clebsch etc. (see for example, [34, Theorem 1.6] and [38, Chapter 7] etc.).

In the last two decades, this beautiful correspondence was extended to higher-dimensional complex projective varieties from the viewpoint of the geometry of hyperplane sections. In particular, after some important contributions by Viro [37] and Dimca [8] etc., Ernström proved the following remarkable result in 1994.

**Theorem 1.3** [9, Corollary 3.9] *Let  $V \subset \mathbb{P}_n$  be a smooth projective variety over  $\mathbb{C}$ . Take a generic hyperplane  $H$  in  $\mathbb{P}_n$  such that  $H \notin V^*$ . Then for any hyperplane  $L \in V^*$ , we have*

$$\chi(V \cap L) - \chi(V \cap H) = (-1)^{n-1+\dim V - \dim V^*} \text{Eu}_{V^*}(L), \quad (1.6)$$

where  $\chi$  stands for the topological Euler characteristic and  $\text{Eu}_{V^*}: V^* \rightarrow \mathbb{Z}$  is the Euler obstruction of  $V^*$  (introduced by Kashiwara [17] and MacPherson [24] independently).

Recall that the Euler obstruction  $\text{Eu}_{V^*}$  of  $V^*$  is a  $\mathbb{Z}$ -valued function on  $V^*$  which measures the singularity of  $V^*$  at each point of  $V^*$ . For example,  $\text{Eu}_{V^*}$  takes the value 1 on the regular part of  $V^*$ . Moreover, if we take a Whitney stratification  $\bigsqcup_{\alpha \in A} V_\alpha^*$  of  $V^*$  consisting of connected strata, then  $\text{Eu}_{V^*}$  is constant on each stratum  $V_\alpha^*$ . The values of  $\text{Eu}_{V^*}$  on a stratum  $V_\alpha^*$  is determined by those on  $V_\beta^*$ 's satisfying the condition  $V_\alpha^* \subset \overline{V_\beta^*}$  (for more detail, see e.g. [18]).

Hence Ernström's result says that the jumping number of the topological Euler characteristics of hyperplane sections of  $V$  at  $L$  is expressed by  $\text{Eu}_{V^*}(L)$ , that is, the singularity of the dual variety  $V^*$  at  $L$ .

The aim of this article is to introduce our results in the real algebraic case similar to this Ernström's one and to survey its theoretical background.

## 2 Main results

Consider a real projective space  $X = \mathbb{R}\mathbb{P}_n$  of dimension  $n$  and its dual  $Y = \mathbb{R}\mathbb{P}_n^*$ . Let  $M \subset X$  be a smooth real projective variety and  $M^* \subset Y$  its dual variety.

We fix a  $\mu$ -stratification  $Y = \bigsqcup_{\alpha \in A} Y_\alpha$  of  $Y = \mathbb{R}\mathbb{P}_n^*$  consisting of connected strata and adapted to  $M^*$ . Note that Trotman [35] proved that this  $\mu$ -condition is equivalent to famous Verdier's w-regularity condition.

**Definition 2.1** We define a  $\mathbb{Z}$ -valued function  $\varphi_M: Y \rightarrow \mathbb{Z}$  on  $Y = \mathbb{R}\mathbb{P}_n^*$  by

$$\varphi_M(H) = \chi(M \cap H) \quad (H \in Y). \quad (2.1)$$

Since the function  $\varphi_M$  is defined by the topological Euler characteristics of hyperplane sections  $M \cap H$  of  $M$ , to obtain results similar to Ernström's formula (1.6) it suffices to describe the function  $\varphi_M$  in terms of the singularities of  $M^*$ . We will show that the whole function  $\varphi_M$  can be reconstructed

from one value  $\varphi_M(y)$  at a point  $y \in Y \setminus M^*$  and the singularities of  $M^*$ . First of all, for the above  $\mu$ -stratification  $Y = \bigsqcup_{\alpha \in A} Y_\alpha$  of  $Y = \mathbb{RP}_n^*$  we can prove the following basic result.

**Proposition 2.2** *The function  $\varphi_M$  is constant on each stratum  $Y_\alpha$ .*

We denote the value of  $\varphi_M$  on  $Y_\alpha$  by  $\varphi_\alpha$ . Our main results are reconstruction theorems of  $\varphi_M$ . Namely, we can determine all the values  $\varphi_\alpha$ 's of  $\varphi_M$  from only one value  $\varphi_M(y)$  at a point  $y \in Y \setminus M^*$  and the topology of  $M^*$ .

To state the first theorem, we introduce two notations concerning dual varieties. Recall that the dual variety  $M^*$  is usually a hypersurface in  $Y = \mathbb{RP}_n^*$ .

**Definition 2.3** (i) We denote the dual defect of  $M$  by

$$\delta^*(M) = (n - 1) - \dim M^*. \quad (2.2)$$

(ii) For a conormal vector  $\vec{p} \in T_{M_{\text{reg}}^*}^* Y$  at  $y \in M_{\text{reg}}^*$ , consider the second fundamental form

$$h_{M^*, \vec{p}}: T_y M^* \times T_y M^* \longrightarrow \mathbb{R}, \quad (2.3)$$

with respect to the canonical (Fubini-Study) metric of  $Y = \mathbb{RP}_n^*$  and set

$$J_{\vec{p}} := \#\{\text{positive eigenvalues of } h_{M^*, \vec{p}}\} + \delta^*(M). \quad (2.4)$$

Now, let us state our first main theorem which describes the values of  $\varphi_M$  on  $Y \setminus M_{\text{sing}}^*$ .

**Theorem 2.4** ([26])

(i) *Assume that  $\delta^*(M) > 0$ . Then on  $Y \setminus M^*$  the function  $\varphi_M$  is constant. Moreover for any  $y \in M_{\text{reg}}^*$  there exists a neighborhood  $U$  of  $y$  such that we have*

$$\varphi_M = d \cdot 1_Y + (-1)^{J_{\vec{p}}} 1_{M_{\text{reg}}^*}. \quad (2.5)$$

on  $U$ , where  $d$  is the value of  $\varphi_M$  on  $Y \setminus M^*$  and  $\vec{p} \in T_{M_{\text{reg}}^*}^* Y$  is a conormal vector at  $y \in M_{\text{reg}}^*$ .

(ii) Assume that  $\delta^*(M) = 0$ , that is  $M^*$  is a hypersurface in  $Y = \mathbb{RP}_n^*$ , and consider the following local situation. Let  $Y_{\alpha_1}$  and  $Y_{\alpha_2}$  be two strata in  $Y \setminus M^*$ ,  $Y_\beta$  an open stratum in  $M_{\text{reg}}^*$  such that  $Y_\beta \subset \overline{Y_{\alpha_i}}$  for  $i = 1, 2$  and  $\vec{p} \in T_{M_{\text{reg}}^*} Y$  a conormal vector at a point  $y \in Y_\beta$  pointing from  $Y_{\alpha_1}$  to  $Y_{\alpha_2}$  (see Figure 2.4.1 below). Then we have

$$\varphi_{\alpha_2} - \varphi_{\alpha_1} = (-1)^{J_{\vec{p}}} - (-1)^{J_{-\vec{p}}} (= 0, \pm 2), \tag{2.6}$$

$$\varphi_\beta = \begin{cases} \frac{1}{2}(\varphi_{\alpha_1} + \varphi_{\alpha_2}) & \text{if } \varphi_{\alpha_1} \neq \varphi_{\alpha_2}, \\ \varphi_{\alpha_1} + (-1)^{J_{\vec{p}}} & \text{if } \varphi_{\alpha_1} = \varphi_{\alpha_2}. \end{cases} \tag{2.7}$$




Figure 2.4.1

**Remark 2.5** (i) If we rewrite (2.5) by Euler characteristics, we obtain an equality analogous to Ernström's formula (1.6). Namely, we have

$$\chi(M \cap L) - \chi(M \cap H) = (-1)^{J_{\vec{p}}} \tag{2.8}$$

for any  $L \in M_{\text{reg}}^*$ , where  $H \in Y \setminus M^*$  is a generic hyperplane in  $Y = \mathbb{RP}_n^*$ .

(ii) In the case where  $M^*$  is a hypersurface, the complement of  $M^*$  is divided into several connected components in general. So when we cross the hypersurface  $M^*$ , the value of the function  $\varphi_M$  may jump. Our formula (2.6) describes this jumping number in terms of the principal curvature  $J_{\vec{p}}$  of  $M_{\text{reg}}^*$ .

Next, we state our second main theorem which reconstructs the values of  $\varphi_M$  on  $M_{\text{sing}}^*$ .

**Theorem 2.6** ([26]) *Let  $k \geq \text{codim}_Y M^*$ . Suppose that the values  $\varphi_\alpha$ 's on  $Y_\alpha$ 's satisfying  $\text{codim}_Y Y_\alpha \leq k$  are already determined. Then the value  $\varphi_\beta$  on  $Y_\beta$  satisfying  $\text{codim}_Y Y_\beta = k + 1$  is given by*

$$\varphi_\beta = \sum_{\alpha: Y_\alpha \cap B \neq \emptyset} \varphi_\alpha \cdot \{\chi(\overline{Y_\alpha} \cap B) - \chi(\partial Y_\alpha \cap B)\}. \tag{2.9}$$

Here we set  $B = B(y, \varepsilon) \cap \{\psi < 0\}$  by taking a small enough open ball  $B(y, \varepsilon)$  centered at a point  $y \in Y_\beta$  and a real-valued real analytic function  $\psi$  defined in a neighborhood of  $y$  satisfying  $\psi^{-1}(0) \supset Y_\beta$  and

$$(y; \text{grad}\psi(y)) \in T_{Y_\beta}^* Y \setminus \bigcup_{\alpha \neq \beta} \overline{T_{Y_\alpha}^* Y} \quad (2.10)$$

(see Figure 2.5.1 below).




Figure 2.5.1

By Theorem 2.6, we can recursively determine the values  $\varphi_\alpha$ 's of  $\varphi_M$  by induction on the codimensions of strata  $Y_\alpha$ 's. Note that this representation of the function  $\varphi_M$  is completely analogous to that of the Euler obstructions in [18].

**Example 2.7** Consider a smooth projective curve  $M$  defined by the homogeneous equation  $x^4 + x^3z + z^4 - y^3z = 0$  in  $\mathbb{RP}_2$  (see Figure 2.6.1 below).





Figure 2.6.1

Then the dual curve  $M^* \subset \mathbb{RP}_2^*$  has a shape as in Figure 2.6.2 below. More precisely, as a  $\mu$ -stratification of  $Y = \mathbb{RP}_2^*$  adapted to  $M^*$ , we can take  $Y = \bigsqcup_{i=0}^{11} Y_i$  in Figure 2.6.2. Since the last strata  $Y_{11}$  is contained in the line at infinity ( $\simeq \mathbb{RP}_1$ ) of  $\mathbb{RP}_2^*$  it does not appear in the figure.



- $\text{codim}_Y Y_0 = 0$
- $\text{codim}_Y Y_1 = 0$
- $\text{codim}_Y Y_2 = 0$
- $\text{codim}_Y Y_3 = 1$
- $\text{codim}_Y Y_4 = 1$
- $\text{codim}_Y Y_5 = 1$
- $\text{codim}_Y Y_6 = 1$
- $\text{codim}_Y Y_7 = 1$
- $\text{codim}_Y Y_8 = 2$
- $\text{codim}_Y Y_9 = 2$
- $\text{codim}_Y Y_{10} = 2$

Figure 2.6.2

Now let us apply our two main theorems to this case. Denote by  $\varphi_i$  the value of the function  $\varphi_M$  on  $Y_i$ . Then we can easily see that  $\varphi_0$  is 0. Starting from this value  $\varphi_0 = 0$ , we can recursively determine all the values  $\varphi_i$ 's of  $\varphi_M$  as follows.




Figure 2.6.3

For example, by Theorem 2.4 the values  $\varphi_1$  and  $\varphi_3$  on  $Y_1$  and  $Y_3$  respectively can be calculated in the following way.

$$\varphi_1 = \varphi_0 + (-1)^2 - (-1)^1 = 2, \tag{2.11}$$

$$\varphi_3 = \frac{1}{2}(\varphi_0 + \varphi_1) = 1. \tag{2.12}$$

Moreover, by Theorem 2.6 the value  $\varphi_{10}$  on  $Y_{10}$  is determined by  $\varphi_1, \varphi_2, \varphi_5$  and  $\varphi_6$  as follows.

$$\varphi_{10} = \varphi_1 \cdot 0 + \varphi_5 \cdot 1 + \varphi_2 \cdot (-1) + \varphi_6 \cdot 1 + \varphi_1 \cdot 0 = 2. \tag{2.13}$$

In this case, we can easily check these results simply by counting the intersection numbers of  $M$  and lines in  $\mathbb{R}P_2$ . Namely, our results are the generalization of this very simple example to higher dimensional cases.

### 3 Theoretical background

Since the function  $\varphi_M$  in our main theorems is constant on each stratum of  $Y$ , we consider a class of such functions to study  $\varphi_M$ , which are called constructible functions.

**Definition 3.1** Let  $X$  be a real analytic manifold. We say that a function  $\varphi: X \rightarrow \mathbb{Z}$  is constructible if there exists a locally finite family  $\{X_i\}$  of compact subanalytic subsets  $X_i$  of  $X$  such that  $\varphi$  is expressed by

$$\varphi = \sum_i c_i \mathbf{1}_{X_i} \quad (c_i \in \mathbb{Z}). \quad (3.1)$$

We denote the abelian group of constructible functions on  $X$  by  $CF(X)$ .

We define the operations of constructible functions in the following way.

**Definition 3.2** ([21] and [37]) Let  $f: Y \rightarrow X$  be a morphism of real analytic manifolds.

- (i) (The inverse image) For  $\varphi \in CF(X)$ , we define a function  $f^*\varphi \in CF(Y)$  by

$$f^*\varphi(y) := \varphi(f(y)). \quad (3.2)$$

- (ii) (The integral) Let  $\varphi = \sum_i c_i \mathbf{1}_{X_i} \in CF(X)$  such that  $\text{supp}(\varphi)$  is compact. Then we define a topological (Euler) integral  $\int_X \varphi \in \mathbb{Z}$  of  $\varphi$  by

$$\int_X \varphi := \sum_i c_i \cdot \chi(X_i). \quad (3.3)$$

- (iii) (The direct image) Let  $\psi \in CF(Y)$  such that  $f|_{\text{supp}(\psi)}: \text{supp}(\psi) \rightarrow X$  is proper. Then we define a function  $\int_f \psi \in CF(X)$  by

$$\left( \int_f \psi \right) (x) := \int_Y (\psi \cdot \mathbf{1}_{f^{-1}(x)}). \quad (3.4)$$



From now on, we shall use various notions concerning derived categories of constructible sheaves. For the detail of these notions, see [21] etc. We denote by  $\mathbf{D}^b(X)$  the derived category of bounded complexes sheaves of  $\mathbf{C}_X$ -modules on  $X$ . Its full subcategory consisting of complexes whose cohomology sheaves are  $\mathbb{R}$ -constructible is denoted by  $\mathbf{D}_{\mathbb{R}-c}^b(X)$ .

Recall also that the Grothendieck group  $\mathbf{K}_{\mathbb{R}-c}(X)$  of  $\mathbf{D}_{\mathbb{R}-c}^b(X)$  is a quotient group of the free abelian group generated by objects of  $\mathbf{D}_{\mathbb{R}-c}^b(X)$  by the subgroup generated by

$$[F] - [F'] - [F''] \quad (F' \longrightarrow F \longrightarrow F'' \xrightarrow{+1} \text{ is a distinguished triangle}). \quad (3.5)$$

Then the natural morphism

$$\chi: \mathbf{K}_{\mathbb{R}-c}(X) \longrightarrow CF(X) \quad (3.6)$$

defined by  $\chi([F])(x) = \sum_{j \in \mathbf{Z}} (-1)^j \dim H^j(F)_x$  ( $x \in X$ ) is an isomorphism.

Moreover, by the isomorphism  $\chi: \mathbf{K}_{\mathbb{R}-c}(X) \xrightarrow{\sim} CF(X)$  the operations of constructible functions that we introduced in Definition 3.2 correspond to those for  $\mathbb{R}$ -constructible sheaves. For example, let  $f: Y \longrightarrow X$  be a morphism of real analytic manifolds and for  $\psi \in CF(Y)$  take an object  $G \in \mathbf{D}_{\mathbb{R}-c}^b(Y)$  such that  $\psi = \chi(G)$ . Then we have  $\chi(Rf_*G) = \int_f \psi$  in  $CF(X)$ . In the same way, we can slightly generalize the notion of topological integrals of constructible functions as follows.

**Definition 3.3** ([26]) Let  $U$  be a relatively compact subanalytic open subset of  $X$  and  $\varphi \in CF(X)$ . Take an object  $F \in \mathbf{D}_{\mathbb{R}-c}^b(X)$  such that  $\varphi = \chi(F)$  and set

$$\int_U \varphi := \chi(R\Gamma(U; F)). \quad (3.7)$$

We can easily check that the definition above does not depend on the choice of  $F$  such that  $\varphi = \chi(F)$ . Note that we do not have to assume that the support of  $\varphi$  is compact in  $U$  as in the usual definition (Definition 3.2 (ii)). Using this slight modification of the notion of topological integrals, we can express the R.H.S of (2.9) simply by  $\int_B \varphi_M$ . In fact, we used this fact in the proof of Theorem 2.6.

Now, let  $X$  be a real analytic manifold and denote by  $\mathcal{L}_X$  the sheaf of conic ( $\mathbb{R}_{>0}$ -invariant) subanalytic Lagrangian cycles in the cotangent bundle  $T^*X$  of  $X$ . Its global section  $H^0(T^*X; \mathcal{L}_X)$  is the abelian group of conic

subanalytic Lagrangian cycles in  $T^*X$ . In 1985, Kashiwara [19] constructed a group homomorphism  $CC: \mathbf{K}_{\mathbb{R}-c}(X) \rightarrow H^0(T^*X; \mathcal{L}_X)$  and associated with each  $[F] \in \mathbf{K}_{\mathbb{R}-c}(X)$  a Lagrangian cycle  $CC([F])$  in  $T^*X$ . This Lagrangian cycle  $CC([F])$  is called the characteristic cycle of  $[F] \in \mathbf{K}_{\mathbb{R}-c}(X)$ . The following very important theorem was proved also in [19] (see [21] for the detail).

**Theorem 3.4** [21, Theorem 9.7.11] *There exists a commutative diagram*

$$\begin{array}{ccc}
 & H^0(T^*X; \mathcal{L}_X) & \\
 \nearrow \sim_{CC} & & \searrow \sim \\
 \mathbf{K}_{\mathbb{R}-c}(X) & \xrightarrow{\sim_{\chi}} & CF(X)
 \end{array} \tag{3.8}$$

*in which all arrows are isomorphisms.*

By this theorem, we can reduce the problem of constructible functions (sheaves) to that of Lagrangian cycles.

### 4 Outline of the proof of main theorems

In this section, we give an outline of the proof of our main theorems. Let  $X = \mathbb{R}P_n$  and  $Y = \mathbb{R}P_n^*$  as before. Consider the incidence submanifold  $S = \{(x, H) \in X \times Y \mid x \in H\}$  of  $X \times Y$  and the diagram

$$\begin{array}{ccc}
 & X \times Y & \\
 p_1 \swarrow & \uparrow S & \searrow p_2 \\
 X & & Y, \\
 f \swarrow & & \searrow g
 \end{array} \tag{4.1}$$

where  $p_1$  and  $p_2$  are natural projections and  $f$  and  $g$  are restrictions of  $p_1$  and  $p_2$  to  $X$  and  $Y$  respectively.

**Definition 4.1** Let  $\varphi \in CF(X)$ . We define the topological Radon transform  $\mathcal{R}_S(\varphi) \in CF(Y)$  of  $\varphi$  by

$$\mathcal{R}_S(\varphi) := \int_g f^* \varphi. \tag{4.2}$$

In particular, for a real analytic submanifold  $M$  of  $X = \mathbb{R}P_n$  and a hyperplane  $H$  in  $X = \mathbb{R}P_n$  ( $\iff H \in Y = \mathbb{R}P_n^*$ ) we have

$$\mathcal{R}_S(\mathbf{1}_M)(H) = \chi(M \cap H) \quad \left( = \varphi_M(H) \right). \tag{4.3}$$

Therefore for the study of the function  $\varphi_M \in CF(Y)$  it suffices to study the topological Radon transform  $\mathcal{R}_S(\mathbf{1}_M)$ . Using the isomorphisms in Theorem 3.4, instead of the topological Radon transform  $\mathcal{R}_S: CF(X) \rightarrow CF(Y)$  itself, we studied the corresponding operation for Lagrangian cycles (characteristic cycles). Then we found an isomorphism

$$\Psi: H^0(\dot{T}^*X; \mathcal{L}_X) \xrightarrow{\sim} H^0(\dot{T}^*Y; \mathcal{L}_Y), \quad (4.4)$$

where we set  $\dot{T}^*X = T^*X \setminus T_X^*X$  and  $\dot{T}^*Y = T^*Y \setminus T_Y^*Y$  (the zero-sections are removed). Moreover this operation  $\Psi$  is (up to some sign  $\varepsilon = \pm 1$ ) the isomorphism of Lagrangian cycles induced by the canonical diffeomorphism  $\Phi: \dot{T}^*X \xrightarrow{\sim} \dot{T}^*Y$  which coincides with the classical Legendre transform in the standard affine charts of  $X = \mathbb{R}\mathbb{P}_n$  and  $Y = \mathbb{R}\mathbb{P}_n^*$ . Since the characteristic cycle  $CC(\mathbf{1}_M)$  of  $\mathbf{1}_M \in CF(X)$  is the conormal cycle  $[\dot{T}_M^*X]$  in  $\dot{T}^*X$ , the characteristic cycle  $CC(\mathcal{R}_S(\mathbf{1}_M))$  of the topological Radon transform  $\mathcal{R}_S(\mathbf{1}_M) = \varphi_M$  is  $\varepsilon[\Phi(\dot{T}_M^*X)]$ . Set  $\dot{\pi}_Y: \dot{T}^*Y \rightarrow Y$  and  $N = (\dot{\pi}_Y \circ \Phi)(\dot{T}_M^*X) \subset Y$ . Then we can easily prove that  $N$  coincides with the dual variety  $M^*$  of  $M$ , which is a closed subanalytic subset of  $Y = \mathbb{R}\mathbb{P}_n^*$  (in classical terminology we call it a caustic or Legendre singularity). Moreover it turns out that the closure  $\overline{\dot{T}_{N_{\text{reg}}}^*Y}$  of the conormal bundle  $\dot{T}_{N_{\text{reg}}}^*Y$  in  $\dot{T}^*Y$  is nothing but  $\Phi(\dot{T}_M^*X)$  (see [16] for a similar argument). Then by using this very nice property of the characteristic cycle  $CC(\varphi_M)$  we can reconstruct the function  $\varphi_M$  from the geometry of the dual variety  $M^* = N$ . Theorem 2.6 was proved in this way. To prove Theorem 2.4, we have to determine the sign  $\varepsilon = \pm 1$ , which is the most difficult part of our study. We could determine it by employing the theory of pure sheaves in [21]. More precisely, we expressed the Maslov indices of the Lagrangian submanifolds  $\dot{T}_M^*X$  and  $\dot{T}_{N_{\text{reg}}}^*Y$  by the principal curvatures of  $M$  and  $N_{\text{reg}}$  respectively with the help of results in [11].

**Remark 4.2** By the same argument as above, we can give a more transparent proof to the main results of Ernström [9] in the complex case.

## 5 Grassmann cases and class formulas

### 5.1 $k$ -dual varieties

We shall generalize the situation considered in the previous sections to Grassmann cases and obtain similar results. Let  $0 \leq k \leq n-1$  be an integer.

Recall that the Grassmann manifold consisting of  $k$ -dimensional planes in  $\mathbb{P}_n$  is defined by

$$\mathbf{G}_{n,k} = \{L' \mid L' \text{ is a } (k+1)\text{-dimensional linear subspace in } \mathbb{K}^{n+1}\} \quad (5.1)$$

$$= \{L \mid L \text{ is a } k\text{-dimensional linear subspace in } \mathbb{P}_n\}. \quad (5.2)$$

Note that  $\mathbb{G}_{n,0} = \mathbb{P}_n$  and  $\mathbb{G}_{n,n-1} = \mathbb{P}_n^*$ . Then the notion of dual varieties is generalized to Grassmann cases as follows.

**Definition 5.1** Let  $V \subset \mathbb{P}_n$  be a projective variety. We define the  $k$ -dual variety  $V^{(k)}$  of  $V$  by

$$V^{(k)} := \overline{\{L \in \mathbb{G}_{n,k} \mid \exists x \in V_{\text{reg}} \cap L \text{ s.t. } V \not\subset L \text{ at } x\}} \subset \mathbb{G}_{n,k}. \quad (5.3)$$

If  $k = n - 1$  the  $k$ -dual  $V^{(k)} \subset \mathbb{G}_{n,k} \simeq \mathbb{P}_n^*$  is nothing but the classical dual variety of  $V$ . In [12], Gelfand-Kapranov-Zelevinsky called  $V^{(k)}$  the associated variety of  $V$  and showed that  $V^{(n-\dim V-1)}$  is a hypersurface.

## 5.2 Topological class formulas

From now on, we always assume that the ground field  $\mathbb{K}$  is  $\mathbb{C}$ . Let  $V \subset \mathbb{P}_n$  be a projective variety over  $\mathbb{C}$  and  $0 \leq k \leq n - 1$  an integer. Assume that  $V^{(k)}$  is a hypersurface in  $\mathbb{G}_{n,k}$ .

**Definition 5.2** [12, Proposition 2.1 of Chapter 3] Consider the Plücker embedding:

$$V^{(k)} \subset \mathbb{G}_{n,k} \subset \mathbb{P}_{\binom{n+1}{k+1}-1}. \quad (5.4)$$

We call the degree of the defining polynomial of  $V^{(k)}$  in  $\mathbb{P}_{\binom{n+1}{k+1}-1}$  the degree of  $V^{(k)}$  and denote it by  $\deg V^{(k)}$ .

In [27], we proved the following topological class formula (i.e. a formula which expresses the degrees of dual varieties) for  $k$ -dual varieties by using Ernström's result [9] and some elementary formulas on constructible functions.

**Theorem 5.3** ([27]) *In the situation as above, for generic linear subspaces  $L_1 \simeq \mathbb{P}_{k-1}$ ,  $L_2 \simeq \mathbb{P}_k$  and  $L_3 \simeq \mathbb{P}_{k+1}$  of  $\mathbb{P}_n$  we have*

$$\deg V^{(k)} = (-1)^{(n-k)+\dim V+1} \left\{ \int_{L_1} \text{Eu}_V - 2 \int_{L_2} \text{Eu}_V + \int_{L_3} \text{Eu}_V \right\}. \quad (5.5)$$

**Corollary 5.4** *Let  $L \simeq \mathbb{P}_{k+1}$  be a generic  $(k+1)$ -dimensional linear subspace of  $\mathbb{P}_n$  and consider the usual dual variety  $(V \cap L)^* \subset \mathbb{P}_{k+1}^*$  of  $V \cap L \subset L \simeq \mathbb{P}_{k+1}$ . Then we have*

$$\deg V^{(k)} = \deg(V \cap L)^*. \quad (5.6)$$

The formula in Theorem 5.3 expresses the algebraic invariant  $\deg V^{(k)}$  of  $V^{(k)}$  by the topological data of  $V$ . In the case where  $k = n - 1$ , we thus reobtain the topological class formulas obtained by Ernström [10], Parusinski and Kleiman [22] etc. See [34, Section 10.1] for an excellent review on this subject. In a forthcoming paper [28], from these topological class formulas we derive various more computable class formulas which extend the previous results obtained by Teissier and Kleiman [23] etc.

## References

- [1] V. I. Arnold, Singularities of caustics and wave fronts, *Mathematics and its Applications* 62, Kluwer, 1987
- [2] V. I. Arnold, S. M. Gusein-Zade and A. N. Varchenko, *Singularities of differential maps I*, Birkhäuser, 1985
- [3] R. Benedetti and J. J. Risler, *Real algebraic and semi-algebraic sets*, Hermann, Paris, 1990
- [4] J. Boman and E. T. Quinto, Support theorems for real-analytic Radon transforms, *Duke Math. J.* 55 (1987), no. 4, 943-948
- [5] J.L. Brylinski, Transformations canoniques, dualité projective, théorie de Lefschetz, transformations de Fourier et sommes trigonométriques. *Géométrie et analyse microlocales*, Astérisque 140-141 (1986), 3-134
- [6] G. Comte, Equisingularité réelle: nombres de Lelong et images polaires, *Ann. Sci. École Norm. Sup.* 33 (2000), 757-788
- [7] A. D'Agnolo and P. Schapira, Radon-Penrose transform for  $\mathcal{D}$ -modules, *J. Funct. Anal.* 139 (1996), no. 2, 349-382
- [8] A. Dimca, Milnor numbers and multiplicities of dual varieties, *Rev. Roumaine Math. Pures Appl.* 31 (1986), 535-538
- [9] L. Ernström, Topological Radon transforms and the local Euler obstruction, *Duke Math. J.* 76 (1994), 1-21
- [10] L. Ernström, A Plücker formula for singular projective varieties, *Communications in algebra*, 25 (1997), 2897-2901
- [11] G. Fischer and J. Piontowski, *Ruled varieties*, Vieweg, 2001
- [12] I. M. Gelfand, M. M. Kapranov and A. V. Zelevinsky, *Discriminants, resultants and multidimensional determinants*, Birkhäuser, 1994

- [13] P. Griffiths and J. Harris, Algebraic geometry and local differential geometry, *Ann. Sci. École Norm. Sup.* (4) 12 (1979), 355-452
- [14] V. Guillemin and S. Stenberg, Geometric asymptotics, Amer. Math. Soc. Providence, Math. Surveys 14, 1977
- [15] S. Helgason, The Radon transform, Progress in Mathematics 5, Birkhäuser, Boston, 1980
- [16] G. Ishikawa and T. Morimoto, Solution surfaces of Monge-Ampère equations, *Differential Geom. Appl.* 14 (2001), no. 2, 113-124
- [17] M. Kashiwara, Index theorem for maximally overdetermined systems of linear differential equations, *Proc. Japan Acad.* 49 (1973), 803-804
- [18] M. Kashiwara, Systems of microdifferential equations, Progress in Mathematics 34, Birkhäuser, Boston, 1983
- [19] M. Kashiwara, Index theorem for constructible sheaves, *Astérisque* 130 (1985), 193-209
- [20] M. Kashiwara and P. Schapira, Microlocal study of sheaves, *Astérisque* 128 (1985)
- [21] M. Kashiwara and P. Schapira, Sheaves on manifolds, Grundlehren Math. Wiss. 292, Springer-Verlag, Berlin-Heidelberg-New York, 1990
- [22] S. L. Kleiman, The enumerative theory of singularities, Real and complex singularities, Sijthoff and Nordhoff international Publishers, Alphen an den Rijn (1977), 297-396
- [23] S. L. Kleiman, A generalized Teissier-Plücker formula, *Contemp. Math.* 162 (1994), 249-260
- [24] R. MacPherson, Chern classes for singular varieties, *Ann. of Math.* 100 (1974), 423-432
- [25] Y. Matsui, Radon transforms of constructible functions on Grassmann manifolds, *Publ. Res. Inst. Math. Sci.*, to appear
- [26] Y. Matsui and K. Takeuchi, Microlocal study of topological Radon transforms and real projective duality, submitting
- [27] Y. Matsui and K. Takeuchi, Topological Radon transforms and degree formulas for dual varieties, submitting
- [28] Y. Matsui and K. Takeuchi, Generalized Plücker-Teissier-Kleiman formulas for varieties with arbitrary dual defect, submitting

- [29] A. Parusinski, Multiplicity of the dual variety, *Bull. London Math. Soc.* 23 (1991), 429-436
- [30] P. Schapira, Tomography of constructible functions, *Lecture Notes Computer Science* 948, Springer Berlin (1995), 427-435
- [31] P. Schapira, Operations on constructible functions, *J. Pure Appl. Algebra* 72, (1991), 83-93
- [32] J. Schürmann, A general intersection formula for Lagrangian cycles, *Compositio Math.* , 140 (2004), 1037-1052
- [33] K. Takeuchi, Microlocal boundary value problem in higher codimensions, *Bull. Soc. Math. France* 124 (1996), 243-276
- [34] E. Tevelev, Projective duality and homogeneous spaces, *Encyclopaedia of mathematical sciences* 133, Springer, 2005
- [35] D. Trotman, Une version microlocale de la condition (w) de Verdier, *Ann. Inst. Fourier Grenoble* 39, (1989), 825-829
- [36] T. Urabe, Duality of the second fundamental form, *Pitman Res. Notes Math. Ser. 381, Real analytic and algebraic singularities (Nagoya/Sapporo/Hachioji, 1996)*, Longman, Harlow, (1998), pp.145-148
- [37] O.Y. Viro, Some integral calculus based on Euler characteristics, *Lecture Notes in Math.* 1346, Springer-Verlag, Berlin (1988), 127-138
- [38] C. T. C. Wall, *Singular points of plane curves*, London Math. Soc. Student Texts 63, Cambridge Univ. Press, 2004