

On dense subsets of boundaries of Coxeter groups

宇都宮大学教育学部

保坂 哲也 (Tetsuya Hosaka)

The purpose of this note is to introduce some results of recent papers [7], [8] and [9] about dense subsets of boundaries of Coxeter groups.

A *Coxeter group* is a group W having a presentation

$$\langle S \mid (st)^{m(s,t)} = 1 \text{ for } s, t \in S \rangle,$$

where S is a finite set and $m : S \times S \rightarrow \mathbb{N} \cup \{\infty\}$ is a function satisfying the following conditions:

- (i) $m(s, t) = m(t, s)$ for any $s, t \in S$,
- (ii) $m(s, s) = 1$ for any $s \in S$, and
- (iii) $m(s, t) \geq 2$ for any $s, t \in S$ such that $s \neq t$.

The pair (W, S) is called a *Coxeter system*. Let (W, S) be a Coxeter system. For a subset $T \subset S$, W_T is defined as the subgroup of W generated by T , and called a *parabolic subgroup*. A subset $T \subset S$ is called a *spherical subset* of S , if the parabolic subgroup W_T is finite. For each $w \in W$, we define $S(w) = \{s \in S \mid \ell(ws) < \ell(w)\}$, where $\ell(w)$ is the minimum length of word in S which represents w . For a subset $T \subset S$, we also define $W^T = \{w \in W \mid S(w) = T\}$.

Let (W, S) be a Coxeter system and let \mathcal{S}^f be the family of spherical subsets of S . We denote $W\mathcal{S}^f$ as the set of all cosets of the form wW_T , with $w \in W$ and $T \in \mathcal{S}^f$. The sets \mathcal{S}^f and $W\mathcal{S}^f$ are partially ordered by inclusion. Contractible simplicial complexes $K(W, S)$ and $\Sigma(W, S)$ are defined as the geometric realizations of the partially ordered sets \mathcal{S}^f and $W\mathcal{S}^f$, respectively ([4]). The natural embedding $\mathcal{S}^f \rightarrow W\mathcal{S}^f$ defined by $T \mapsto W_T$ induces an embedding $K(W, S) \rightarrow \Sigma(W, S)$ which we regard as an inclusion. The group W acts on $\Sigma(W, S)$ via simplicial automorphism. Then $\Sigma(W, S) = WK(W, S)$

and $\Sigma(W, S)/W \cong K(W, S)$ ([4]). For each $w \in W$, $wK(W, S)$ is called a *chamber* of $\Sigma(W, S)$. If W is infinite, then $\Sigma(W, S)$ is noncompact. In [10], G. Moussong proved that a natural metric on $\Sigma(W, S)$ satisfies the CAT(0) condition. Hence, if W is infinite, $\Sigma(W, S)$ can be compactified by adding its ideal boundary $\partial\Sigma(W, S)$ ([4], [3]). This boundary $\partial\Sigma(W, S)$ is called the *boundary of (W, S)* . We note that the natural action of W on $\Sigma(W, S)$ is properly discontinuous and cocompact ([4]), and this action induces an action of W on $\partial\Sigma(W, S)$.

A subset A of a space X is said to be *dense* in X , if $\bar{A} = X$. A subset A of a metric space X is said to be *quasi-dense*, if there exists $N > 0$ such that each point of X is N -close to some point of A .

Let (W, S) be a Coxeter system. Then W has the *word metric* d_ℓ defined by $d_\ell(w, w') = \ell(w^{-1}w')$ for each $w, w' \in W$.

In [7], the following theorems were proved.

Theorem 1. *Let (W, S) be a Coxeter system. Suppose that $W^{\{s_0\}}$ is quasi-dense in W with respect to the word metric and $o(s_0t_0) = \infty$ for some $s_0, t_0 \in S$, where $o(s_0t_0)$ is the order of s_0t_0 in W . Then there exists $\alpha \in \partial\Sigma(W, S)$ such that the orbit $W\alpha$ is dense in $\partial\Sigma(W, S)$.*

Suppose that a group Γ acts properly and cocompactly by isometries on a CAT(0) space X . Every element $\gamma \in \Gamma$ such that $o(\gamma) = \infty$ is a hyperbolic transformation of X , i.e., there exists a geodesic axis $c : \mathbb{R} \rightarrow X$ and a real number $a > 0$ such that $\gamma \cdot c(t) = c(t + a)$ for each $t \in \mathbb{R}$ ([3]). Then, for all $x \in X$, the sequence $\{\gamma^i x\}$ converges to $c(\infty)$ in $X \cup \partial X$. We denote $\gamma^\infty = c(\infty)$.

Theorem 2. *Let (W, S) be a Coxeter system. If the set*

$$\bigcup \{W^{\{s\}} \mid s \in S \text{ such that } o(st) = \infty \text{ for some } t \in S\}$$

is quasi-dense in W , then $\{w^\infty \mid w \in W \text{ such that } o(w) = \infty\}$ is dense in $\partial\Sigma(W, S)$.

Remark. For a negatively curved group G and the boundary ∂G of G ,

- (1) we can show that $G\alpha$ is dense in ∂G for each $\alpha \in \partial G$ by an easy argument, and

(2) it is known that $\{g^\infty \mid g \in G \text{ such that } o(g) = \infty\}$ is dense in ∂G ([2]).

As an application of Theorems 1 and 2, we obtained the following theorem in [7].

Theorem 3. *Let (W, S) be a Coxeter system. Suppose that there exist a maximal spherical subset T of S and an element $s_0 \in S$ such that $o(s_0 t) \geq 3$ for each $t \in T$ and $o(s_0 t_0) = \infty$ for some $t_0 \in T$. Then*

- (1) $W\alpha$ is dense in $\partial\Sigma(W, S)$ for some $\alpha \in \partial\Sigma(W, S)$, and
- (2) $\{w^\infty \mid w \in W \text{ such that } o(w) = \infty\}$ is dense in $\partial\Sigma(W, S)$.

Example. The Coxeter system defined by the diagram in Figure 1 is not hyperbolic in Gromov sense, since it contains a copy of \mathbb{Z}^2 , and it satisfies the condition of Theorem 3.




FIGURE 1

Suppose that a group G acts on a compact metric space X by homeomorphisms. Then X is said to be *minimal*, if every orbit Gx is dense in X .

For a negatively curved group G and the boundary ∂G of G , $G\alpha$ is dense in ∂G for each $\alpha \in \partial G$, that is, ∂G is minimal.

We note that Coxeter groups are non-positive curved groups and not negatively curved groups in general. There exist examples of Coxeter systems whose boundaries are not minimal as follows.

Example. Let $S = \{s, t, u\}$ and let

$$W = \langle S \mid s^2 = t^2 = u^2 = (st)^3 = (tu)^3 = (us)^3 = 1 \rangle.$$

Then (W, S) is a Coxeter system and $\Sigma(W, S)$ is the flat Euclidean plane. For any $\alpha \in \partial\Sigma(W, S)$, $W\alpha$ is a finite-points set and not dense in $\partial\Sigma(W, S)$ which is a circle. This example implies that we can not omit the assumption " $m(s_0, t_0) = \infty$ " in Theorem 3.

Example. Let $S = \{s_1, s_2, s_3, s_4\}$ and let

$$W = \langle S \mid s_1^2 = s_2^2 = s_3^2 = s_4^2 = (s_1s_2)^2 = (s_2s_3)^2 = (s_3s_4)^2 = (s_4s_1)^2 = 1 \rangle.$$

Then (W, S) is a Coxeter system and $\Sigma(W, S)$ is the Euclidean plane. For any $\alpha \in \partial\Sigma(W, S)$, $W\alpha$ is a finite-points set and not dense in $\partial\Sigma(W, S)$ which is a circle. Here we note that $\{s_1, s_2\}$ is a maximal spherical subset of S , $m(s_1, s_3) = \infty$ and $m(s_2, s_3) = 2$. This example implies that we can not omit the assumption “ $m(s_0, t) \geq 3$ ” in Theorem 3.

As an extension of Theorem 3, we have obtained the following theorem in [9].

Theorem 4. *Let (W, S) be a Coxeter system which satisfies the condition in Theorem 3. Then every orbit $W\alpha$ is dense in $\partial\Sigma(W, S)$, that is, $\partial\Sigma(W, S)$ is minimal.*

Here the following problems are open.

Problem. Does there exist a Coxeter system (W, S) such that some orbit $W\alpha$ is dense in $\partial\Sigma(W, S)$ and $\partial\Sigma(W, S)$ is not minimal?

Problem. Suppose that a group G acts geometrically on two CAT(0) spaces X and X' . Is it the case that ∂X is minimal if and only if $\partial X'$ is minimal?

Problem (Ruane). Suppose that a group G acts geometrically on a CAT(0) space X . Is it always the case that the set $\{g^\infty \mid g \in G, o(g) = \infty\}$ is dense in ∂X ?

In [8], we also have obtained the following theorem.

Theorem 5. *Let (W, S) be a Coxeter system and let T be a subset of S such that W_T is infinite. If the set*

$$\bigcup \{W^{ts} \mid s \in S \text{ such that } o(ss_0) = \infty \text{ and } s_0t \neq ts_0 \\ \text{for some } s_0 \in S \setminus T \text{ and } t \in \tilde{T}\}$$

is quasi-dense in W with respect to the word metric, then $W\partial\Sigma(W_T, T)$ is dense in $\partial\Sigma(W, S)$, where $W_{\tilde{T}}$ is the essential parabolic subgroup of (W_T, T) .

If W is a hyperbolic Coxeter group, then $W\partial\Sigma(W_T, T)$ is dense in $\partial\Sigma(W, S)$ for any $T \subset S$ such that W_T is infinite.

As an application of Theorem 5, we have obtained the following corollary in [8].

Corollary 6. *Let (W, S) be a Coxeter system and let T be a subset of S such that W_T is infinite. Suppose that there exist a maximal spherical subset U of S and an element $s \in S$ such that $o(su) \geq 3$ for every $u \in U$ and $o(su_0) = \infty$ for some $u_0 \in U$. If*

- (1) $s \notin T$ and $u_0 \in \tilde{T}$, or
- (2) $u_0 \notin T$ and $s \in \tilde{T}$,

then $W\partial\Sigma(W_T, T)$ is dense in $\partial\Sigma(W, S)$.

Concerning W -invariantness of $\partial\Sigma(W_T, T)$, the following theorem is known.

Theorem 7 ([6]).

- (1) *Let (W, S) be a Coxeter system and $T \subset S$. Then $\partial\Sigma(W_T, T)$ is W -invariant if and only if $W = W_{\tilde{T}} \times W_{S \setminus \tilde{T}}$.*
- (2) *Let (W, S) be an irreducible Coxeter system and let T be a proper subset of S such that W_T is infinite. Then $\partial\Sigma(W_T, T)$ is not W -invariant.*

Here the following problem is open.

Problem. Let (W, S) be a Coxeter system and let T be a subset of S such that W_T is infinite. Is it the case that if $\partial\Sigma(W_T, T)$ is not W -invariant then $W\partial\Sigma(W_T, T)$ is dense in $\partial\Sigma(W, S)$? Particulary, is it the case that if (W, S) is an irreducible Coxeter system then $W\partial\Sigma(W_T, T)$ is dense in $\partial\Sigma(W, S)$ for any subset T of S such that W_T is infinite?

REFERENCES

- [1] N. Bourbaki, *Groupes et Algèbres de Lie*, Chapters IV-VI, Masson, Paris, 1981.
- [2] P. L. Bowers and K. Ruane, *Fixed points in boundaries of negatively curved groups*, Proc. Amer. Math. Soc. **124** (no.4) (1996), 1311–1313.
- [3] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Springer-Verlag, Berlin, 1999.
- [4] M. W. Davis, *Nonpositive curvature and reflection groups*, in Handbook of geometric topology (Edited by R. J. Daverman and R. B. Sher), pp. 373–422, North-Holland, Amsterdam, 2002.

- [5] M. Gromov, *Hyperbolic groups*, in Essays in group theory (Edited by S. M. Gersten), pp. 75–263, M.S.R.I. Publ. 8, 1987.
- [6] T. Hosaka, *Parabolic subgroups of finite index in Coxeter groups*, J. Pure Appl. Algebra **169** (2002), 215–227.
- [7] T. Hosaka, *Dense subsets of the boundary of a Coxeter system*, Proc. Amer. Math. Soc. **132** (2004), 3441–3448.
- [8] T. Hosaka, *Addendum to “Dense subsets of the boundary of a Coxeter system”*, Proc. Amer. Math. Soc. **133** (2005), 3745–3747.
- [9] T. Hosaka, *On dense orbits in the boundary of a Coxeter system*, J. Kyoto Univ., to appear.
- [10] G. Moussong, *Hyperbolic Coxeter groups*, Ph.D. thesis, The Ohio State University, 1988.

DEPARTMENT OF MATHEMATICS, UTSUNOMIYA UNIVERSITY, UTSUNOMIYA,
321-8505, JAPAN

E-mail address: hosaka@cc.utsunomiya-u.ac.jp