

# Algebraic independence of the values of certain functions at distinct algebraic points

慶應義塾大学理工学部 田中 孝明 (Taka-aki Tanaka)  
Faculty of Science and Technology, Keio Univ.

## 1 Introduction and results

Loxton and van der Poorten [2] obtained the following result: Let

$$F(z) = \sum_{k=0}^{\infty} z^{d^k},$$

where  $d$  is an integer greater than 1, and let  $\alpha_1, \dots, \alpha_r$  be algebraic numbers with  $0 < |\alpha_i| < 1$  ( $1 \leq i \leq r$ ). Then the following three properties are equivalent:

- (i)  $F(\alpha_1), \dots, F(\alpha_r)$  are algebraically dependent.
- (ii)  $1, F(\alpha_1), \dots, F(\alpha_r)$  are linearly dependent over the field  $\overline{\mathbb{Q}}$  of algebraic numbers.
- (iii) There exist a non-empty subset  $\{\alpha_{i_1}, \dots, \alpha_{i_t}\}$  of  $\{\alpha_1, \dots, \alpha_r\}$ , nonnegative integers  $k_1, \dots, k_t$ , roots of unity  $\zeta_1, \dots, \zeta_t$ , an algebraic number  $\gamma$  with  $\alpha_{i_l}^{d^{k_l}} = \zeta_l \gamma$  ( $1 \leq l \leq t$ ), and algebraic numbers  $\xi_1, \dots, \xi_t$ , not all zero, such that

$$\sum_{l=1}^t \xi_l \zeta_l^{d^k} = 0 \quad (k = 0, 1, 2, \dots).$$

In contrast with this result we consider the power series

$$f(z) = \sum_{k=0}^{\infty} z^{a_k},$$

where  $\{a_k\}_{k \geq 0}$  is a linear recurrence of positive integers which is not a geometric progression and which satisfies

$$a_{k+n} = c_1 a_{k+n-1} + \dots + c_n a_k \quad (k = 0, 1, 2, \dots), \quad (1)$$

where  $c_1, \dots, c_n$  are nonnegative integers with  $c_n \neq 0$ . We assume that the polynomial  $\Phi(X) = X^n - c_1 X^{n-1} - \dots - c_n$  associated with (1) satisfies  $\Phi(\pm 1) \neq 0$  and the ratio of

any pair of distinct roots of  $\Phi(X)$  is not a root of unity. For this power series  $f(z)$ , the author obtained the necessary and sufficient condition for the numbers  $f(\alpha_1), \dots, f(\alpha_r)$  to be algebraically dependent.

**DEFINITION 1.** We say that the algebraic numbers  $\alpha_1, \dots, \alpha_r$  with  $0 < |\alpha_i| < 1$  ( $1 \leq i \leq r$ ) are  $\{a_k\}_{k \geq 0}$ -dependent if there exist a non-empty subset  $\{\alpha_{i_1}, \dots, \alpha_{i_t}\}$  of  $\{\alpha_1, \dots, \alpha_r\}$ , roots of unity  $\zeta_1, \dots, \zeta_t$ , an algebraic number  $\gamma$  with  $\alpha_{i_l} = \zeta_l \gamma$  ( $1 \leq l \leq t$ ), and algebraic numbers  $\xi_1, \dots, \xi_t$ , not all zero, such that

$$\sum_{l=1}^t \xi_l \zeta_l^{a_k} = 0$$

for all sufficiently large  $k$ .

**Theorem 1** (A special case of Theorem 2 in [6]). *Let  $\{a_k\}_{k \geq 0}$  be a linear recurrence satisfying (1). Let  $\alpha_1, \dots, \alpha_r$  be algebraic numbers with  $0 < |\alpha_i| < 1$  ( $1 \leq i \leq r$ ). Then the following three properties are equivalent:*

- (i)  $f(\alpha_1), \dots, f(\alpha_r)$  are algebraically dependent.
- (ii)  $1, f(\alpha_1), \dots, f(\alpha_r)$  are linearly dependent over  $\overline{\mathbb{Q}}$ .
- (iii)  $\alpha_1, \dots, \alpha_r$  are  $\{a_k\}_{k \geq 0}$ -dependent.

**REMARK 1.** In Theorem 1 it is obvious that the property (iii) implies (ii), since  $\sum_{l=1}^t \xi_l f(\alpha_{i_l}) \in \overline{\mathbb{Q}}$  if  $\alpha_1, \dots, \alpha_r$  are  $\{a_k\}_{k \geq 0}$ -dependent.

**REMARK 2.** As a special case of the result of Nishioka [4], the three properties (i)–(iii) in Theorem 1 are equivalent also for a gap series  $\sum_{k=0}^{\infty} z^{a_k}$  with  $\{a_k\}_{k \geq 0}$  an increasing sequence of positive integers such that  $\lim_{k \rightarrow \infty} a_{k+1}/a_k = \infty$ . In the case of our linear recurrence  $\{a_k\}_{k \geq 0}$  satisfying (1), we have  $\lim_{k \rightarrow \infty} a_{k+1}/a_k = \rho$  with  $1 < \rho < \infty$ .

In what follows, let

$$f(z) = \sum_{k=0}^{\infty} z^{a_k}, \quad g(z) = \sum_{k=0}^{\infty} \frac{z^{a_k}}{1 - z^{a_k}}, \quad h(z) = \prod_{k=0}^{\infty} (1 - z^{a_k}).$$

The author proved the following:

**Theorem 2** ([7, Theorem 5]). *Let  $\{a_k\}_{k \geq 0}$  be a linear recurrence satisfying (1). Let  $\alpha_1, \dots, \alpha_r$  be algebraic numbers with  $0 < |\alpha_i| < 1$  ( $1 \leq i \leq r$ ) such that none of  $\alpha_i/\alpha_j$  ( $1 \leq i < j \leq r$ ) is a root of unity. Then the  $3r$  numbers  $f(\alpha_i), g(\alpha_i), h(\alpha_i)$  ( $1 \leq i \leq r$ ) are algebraically independent.*

**REMARK 3.** If  $\{a_k\}_{k \geq 0}$  is a geometric progression, namely  $a_k = ad^k$  ( $k \geq 0$ ) for some integers  $a \geq 1$  and  $d \geq 2$ , each of the  $3r$  numbers in Theorem 2 is transcendental by the

theorem of Mahler [3] ; however Theorem 2 is not valid in this case, since there exist the following relations over  $\overline{\mathbb{Q}}$ : Let

$$F(z) = \sum_{k=0}^{\infty} z^{ad^k}, \quad G(z) = \sum_{k=0}^{\infty} \frac{z^{ad^k}}{1 - z^{ad^k}}, \quad H(z) = \prod_{k=0}^{\infty} (1 - z^{ad^k}),$$

and let  $\alpha$  be an algebraic number with  $0 < |\alpha| < 1$ . Then

$$F(\alpha) - F(\alpha^d) = \alpha^a, \quad G(\alpha) - G(\alpha^d) = \frac{\alpha^a}{1 - \alpha^a}, \quad \frac{H(\alpha)}{H(\alpha^d)} = 1 - \alpha^a,$$

whereas  $\alpha/\alpha^d$  is not a root of unity.

**REMARK 4.** The assumption in Theorem 2 that none of  $\alpha_i/\alpha_j$  ( $1 \leq i < j \leq r$ ) is a root of unity cannot be removed. For example, suppose that the initial values  $a_0, \dots, a_{n-1}$  are divided by an integer  $d > 1$ . Then by the linear recurrence relation (1),  $a_k$  is divided by  $d$  for any  $k \geq 0$ . If  $\alpha_i/\alpha_j$  is a  $d$ -th root of unity for some distinct  $i$  and  $j$ , then  $\alpha_i^{a_k} = \alpha_j^{a_k}$  ( $k \geq 0$ ) and so the numbers considered in Theorem 2 are algebraically dependent. Even in some cases where  $a_0, \dots, a_{n-1}$  have no common factor, the assumption is also inevitable as shown in the following example:

Let  $\{a_k\}_{k \geq 0}$  be a linear recurrence defined by

$$a_0 = 2, \quad a_1 = 3, \quad a_{k+2} = 6a_{k+1} + a_k \quad (k = 0, 1, 2, \dots).$$

We put

$$f(z) = \sum_{k=0}^{\infty} z^{a_k}, \quad g(z) = \sum_{k=0}^{\infty} \frac{z^{a_k}}{1 - z^{a_k}}, \quad h(z) = \prod_{k=0}^{\infty} (1 - z^{a_k}).$$

Let  $\alpha$  be an algebraic number with  $0 < |\alpha| < 1$  and let  $\zeta = e^{\pi\sqrt{-1}/3} = (1 + \sqrt{-3})/2$ . Then

$$\begin{aligned} 2f(\alpha) + f(\zeta\alpha) - f(\zeta^2\alpha) - 2f(\zeta^3\alpha) - f(\zeta^4\alpha) + f(\zeta^5\alpha) &= 0, \\ 2g(\alpha) + g(\zeta\alpha) - g(\zeta^2\alpha) - 2g(\zeta^3\alpha) - g(\zeta^4\alpha) + g(\zeta^5\alpha) &= 0, \end{aligned}$$

and

$$h(\alpha)^2 h(\zeta\alpha) h(\zeta^2\alpha)^{-1} h(\zeta^3\alpha)^{-2} h(\zeta^4\alpha)^{-1} h(\zeta^5\alpha) = 1,$$

since  $a_{2k} \equiv 2 \pmod{6}$  and  $a_{2k+1} \equiv 3 \pmod{6}$  for any  $k \geq 0$ .

The author obtained the necessary and sufficient condition for the  $3r$  numbers  $f(\alpha_i), g(\alpha_i), h(\alpha_i)$  ( $1 \leq i \leq r$ ) in Theorem 2 to be algebraically dependent:

**Theorem 3** ([8]). *Let  $\{a_k\}_{k \geq 0}$  be a linear recurrence satisfying (1). Let  $\alpha_1, \dots, \alpha_r$  be algebraic numbers with  $0 < |\alpha_i| < 1$  ( $1 \leq i \leq r$ ). Then the numbers  $f(\alpha_i), g(\alpha_i), h(\alpha_i)$  ( $1 \leq i \leq r$ ) are algebraically dependent if and only if the algebraic numbers  $\alpha_1, \dots, \alpha_r$  are  $\{a_k\}_{k \geq 0}$ -dependent.*

Combining Theorems 1 and 3, we immediately have the following:

**Theorem 4** ([8]). *Let  $\alpha_1, \dots, \alpha_r$  be algebraic numbers with  $0 < |\alpha_i| < 1$  ( $1 \leq i \leq r$ ). If the numbers  $f(\alpha_1), \dots, f(\alpha_r)$  are algebraically independent, then so are the numbers  $f(\alpha_i), g(\alpha_i), h(\alpha_i)$  ( $1 \leq i \leq r$ ).*

Theorem 4 implies the following:

**Theorem 5** ([8]). *Let  $\alpha_1, \dots, \alpha_r$  be algebraic numbers with  $0 < |\alpha_i| < 1$  ( $1 \leq i \leq r$ ). Then*

$$\begin{aligned} & \text{trans. deg}_{\mathbb{Q}} \mathbb{Q}(f(\alpha_1), \dots, f(\alpha_r), g(\alpha_1), \dots, g(\alpha_r), h(\alpha_1), \dots, h(\alpha_r)) \\ & \geq 3 \text{trans. deg}_{\mathbb{Q}} \mathbb{Q}(f(\alpha_1), \dots, f(\alpha_r)). \end{aligned} \quad (2)$$

The following is an example in which the equality of (2) holds:

**EXAMPLE 1.** Let  $\{a_k\}_{k \geq 0}$  be a linear recurrence defined by

$$a_0 = 1, \quad a_1 = 2, \quad a_{k+2} = 3a_{k+1} + a_k \quad (k = 0, 1, 2, \dots).$$

We put

$$f(z) = \sum_{k=0}^{\infty} z^{a_k}, \quad g(z) = \sum_{k=0}^{\infty} \frac{z^{a_k}}{1 - z^{a_k}}, \quad h(z) = \prod_{k=0}^{\infty} (1 - z^{a_k}).$$

Let  $\alpha$  be an algebraic number with  $0 < |\alpha| < 1$  and let  $\omega = e^{2\pi\sqrt{-1}/3} = (-1 + \sqrt{-3})/2$ . Since  $a_{2k} \equiv 1 \pmod{3}$  and  $a_{2k+1} \equiv 2 \pmod{3}$  for any  $k \geq 0$ , the numbers  $\alpha, \omega\alpha$ , and  $\alpha^3$  are not  $\{a_k\}_{k \geq 0}$ -dependent. Therefore the numbers  $f(\alpha), f(\omega\alpha), f(\alpha^3), g(\alpha), g(\omega\alpha), g(\alpha^3), h(\alpha), h(\omega\alpha), h(\alpha^3)$  are algebraically independent by Theorem 3. Noting that  $f(\alpha) + f(\omega\alpha) + f(\omega^2\alpha) = 0$ ,  $g(\alpha) + g(\omega\alpha) + g(\omega^2\alpha) = 3g(\alpha^3)$ , and  $h(\alpha)h(\omega\alpha)h(\omega^2\alpha) = h(\alpha^3)$ , we see that

$$\text{trans. deg}_{\mathbb{Q}} \mathbb{Q}(f(\alpha), f(\omega\alpha), f(\omega^2\alpha), f(\alpha^3)) = 3,$$

$$\text{trans. deg}_{\mathbb{Q}} \mathbb{Q}(g(\alpha), g(\omega\alpha), g(\omega^2\alpha), g(\alpha^3)) = 3,$$

$$\text{trans. deg}_{\mathbb{Q}} \mathbb{Q}(h(\alpha), h(\omega\alpha), h(\omega^2\alpha), h(\alpha^3)) = 3,$$

and

$$\begin{aligned} & \text{trans. deg}_{\mathbb{Q}} \mathbb{Q}(f(\alpha), f(\omega\alpha), f(\omega^2\alpha), f(\alpha^3), \\ & g(\alpha), g(\omega\alpha), g(\omega^2\alpha), g(\alpha^3), h(\alpha), h(\omega\alpha), h(\omega^2\alpha), h(\alpha^3)) = 9. \end{aligned}$$

As shown in Remark 4 or in the example above, it seems complicated to state the necessary and sufficient condition for the values of the Lambert series  $g(z)$  and the infinite product  $h(z)$  at  $\{a_k\}_{k \geq 0}$ -dependent algebraic numbers  $\alpha_1, \dots, \alpha_r$  to be algebraically

independent. In Theorem 6 below we establish an easily confirmable condition under which such values are algebraically independent.

**DEFINITION 2.** We say that the algebraic numbers  $\alpha_1, \dots, \alpha_r$  with  $0 < |\alpha_i| < 1$  ( $1 \leq i \leq r$ ) are *strongly*  $\{a_k\}_{k \geq 0}$ -*dependent* if there exist a non-empty subset  $\{\alpha_{i_1}, \dots, \alpha_{i_t}\}$  of  $\{\alpha_1, \dots, \alpha_r\}$ ,  $N$ -th roots of unity  $\zeta_1, \dots, \zeta_t$ , an algebraic number  $\gamma$  with  $\alpha_{i_l} = \zeta_l \gamma$  ( $1 \leq l \leq t$ ), and algebraic numbers  $\xi_1, \dots, \xi_t$ , not all zero, such that

$$\sum_{l=1}^t \xi_l \zeta_l^{ma_k} = 0, \quad m = 1, \dots, N-1, \quad \text{g.c.d.}(m, N) = 1,$$

for all sufficiently large  $k$ .

It is clear that, if the algebraic numbers  $\alpha_1, \dots, \alpha_r$  with  $0 < |\alpha_i| < 1$  ( $1 \leq i \leq r$ ) are strongly  $\{a_k\}_{k \geq 0}$ -dependent, then they are  $\{a_k\}_{k \geq 0}$ -dependent.

The following theorem is more precise than Theorem 4 above.

**Theorem 6** ([8]). *Let  $\{a_k\}_{k \geq 0}$  be a linear recurrence satisfying (1). Let  $\alpha_1, \dots, \alpha_r$  be algebraic numbers with  $0 < |\alpha_i| < 1$  ( $1 \leq i \leq r$ ). Suppose that the algebraic numbers  $\alpha_1, \dots, \alpha_r$  are not strongly  $\{a_k\}_{k \geq 0}$ -dependent. Assume further that  $\alpha_1, \dots, \alpha_\rho$  ( $\rho \leq r$ ) are not  $\{a_k\}_{k \geq 0}$ -dependent or equivalently that the numbers  $f(\alpha_1), \dots, f(\alpha_\rho)$  are algebraically independent. Then the numbers  $f(\alpha_1), \dots, f(\alpha_\rho), g(\alpha_1), \dots, g(\alpha_r), h(\alpha_1), \dots, h(\alpha_r)$  are algebraically independent.*

Using Theorem 6, we have an example in which the strict inequality of (2) holds:

**EXAMPLE 2.** Let  $\{a_k\}_{k \geq 0}$  be a linear recurrence defined by

$$a_0 = 1, \quad a_1 = 3, \quad a_{k+2} = 3a_{k+1} + a_k \quad (k = 0, 1, 2, \dots).$$

We put

$$f(z) = \sum_{k=0}^{\infty} z^{a_k}, \quad g(z) = \sum_{k=0}^{\infty} \frac{z^{a_k}}{1 - z^{a_k}}, \quad h(z) = \prod_{k=0}^{\infty} (1 - z^{a_k}).$$

Let  $\alpha$  be an algebraic number with  $0 < |\alpha| < 1$  and let  $\omega = e^{2\pi\sqrt{-1}/3} = (-1 + \sqrt{-3})/2$ . Since  $a_{2k} \equiv 1 \pmod{3}$  and  $a_{2k+1} \equiv 0 \pmod{3}$  for any  $k \geq 0$ , the numbers  $\alpha$ ,  $\omega\alpha$ ,  $\omega^2\alpha$ , and  $\alpha^3$  are not strongly  $\{a_k\}_{k \geq 0}$ -dependent and the numbers  $\alpha$ ,  $\omega\alpha$ , and  $\alpha^3$  are not  $\{a_k\}_{k \geq 0}$ -dependent. Therefore the numbers  $f(\alpha), f(\omega\alpha), f(\alpha^3), g(\alpha), g(\omega\alpha), g(\omega^2\alpha), g(\alpha^3), h(\alpha), h(\omega\alpha), h(\omega^2\alpha), h(\alpha^3)$  are algebraically independent by Theorem 6 with  $\rho = 3$  and  $r = 4$ . Noting that  $\omega f(\alpha) - (\omega + 1)f(\omega\alpha) + f(\omega^2\alpha) = 0$ , we see that

$$\text{trans. deg}_{\mathbb{Q}} \mathbb{Q}(f(\alpha), f(\omega\alpha), f(\omega^2\alpha), f(\alpha^3)) = 3,$$

$$\begin{aligned} & \text{trans. deg}_{\mathbb{Q}} \mathbb{Q}(f(\alpha), f(\omega\alpha), f(\omega^2\alpha), f(\alpha^3), \\ & g(\alpha), g(\omega\alpha), g(\omega^2\alpha), g(\alpha^3), h(\alpha), h(\omega\alpha), h(\omega^2\alpha), h(\alpha^3)) = 11, \end{aligned}$$

and so

$$\begin{aligned} & \text{trans. deg}_{\mathbb{Q}} \mathbb{Q}(f(\alpha), f(\omega\alpha), f(\omega^2\alpha), f(\alpha^3), \\ & g(\alpha), g(\omega\alpha), g(\omega^2\alpha), g(\alpha^3), h(\alpha), h(\omega\alpha), h(\omega^2\alpha), h(\alpha^3)) \\ & > 3 \text{ trans. deg}_{\mathbb{Q}} \mathbb{Q}(f(\alpha), f(\omega\alpha), f(\omega^2\alpha), f(\alpha^3)). \end{aligned}$$

## 2 Proof of Theorems 3 and 6

*Proof of Theorem 3.* If the algebraic numbers  $\alpha_1, \dots, \alpha_r$  are  $\{a_k\}_{k \geq 0}$ -dependent, then the numbers  $f(\alpha_i), g(\alpha_i), h(\alpha_i)$  ( $1 \leq i \leq r$ ) are algebraically dependent, since so are the numbers  $f(\alpha_1), \dots, f(\alpha_r)$  by Theorem 1 with Remark 1. Conversely, if the algebraic numbers  $\alpha_1, \dots, \alpha_r$  are not  $\{a_k\}_{k \geq 0}$ -dependent, then by Theorem 6 with  $\rho = r$  the numbers  $f(\alpha_i), g(\alpha_i), h(\alpha_i)$  ( $1 \leq i \leq r$ ) are algebraically independent. This completes the proof of the theorem.

*Sketch of the proof of Theorem 6.* Suppose on the contrary that the numbers  $f(\alpha_1), \dots, f(\alpha_r), g(\alpha_1), \dots, g(\alpha_r), h(\alpha_1), \dots, h(\alpha_r)$  are algebraically dependent. There exist multiplicatively independent algebraic numbers  $\beta_1, \dots, \beta_s$  with  $0 < |\beta_j| < 1$  ( $1 \leq j \leq s$ ) such that

$$\alpha_i = \zeta_i \prod_{j=1}^s \beta_j^{e_{ij}} \quad (1 \leq i \leq r), \quad (3)$$

where  $\zeta_1, \dots, \zeta_r$  are roots of unity and  $e_{ij}$  ( $1 \leq i \leq r, 1 \leq j \leq s$ ) are nonnegative integers (cf. Nishioka [5, Lemma 3.4.9]). Take a positive integer  $N$  such that  $\zeta_i^N = 1$  for any  $i$  ( $1 \leq i \leq r$ ). We can choose a positive integer  $p$  and a nonnegative integer  $q$  such that  $a_{k+p} \equiv a_k \pmod{N}$  for any  $k \geq q$ . Let  $y_{jl}$  ( $1 \leq j \leq s, 1 \leq l \leq n$ ) be variables and let  $\mathbf{y} = (y_{11}, \dots, y_{1n}, \dots, y_{s1}, \dots, y_{sn})$ . Define the auxiliary functions

$$\begin{aligned} f_i(\mathbf{y}) &= \sum_{k=q}^{\infty} \zeta_i^{a_k} \prod_{j=1}^s (y_{j1}^{a_{k+n-1}} \dots y_{jn}^{a_k})^{e_{ij}} \quad (1 \leq i \leq \rho), \\ g_i(\mathbf{y}) &= \sum_{k=q}^{\infty} \frac{\zeta_i^{a_k} \prod_{j=1}^s (y_{j1}^{a_{k+n-1}} \dots y_{jn}^{a_k})^{e_{ij}}}{1 - \zeta_i^{a_k} \prod_{j=1}^s (y_{j1}^{a_{k+n-1}} \dots y_{jn}^{a_k})^{e_{ij}}} \quad (1 \leq i \leq r), \end{aligned}$$

and

$$h_i(\mathbf{y}) = \prod_{k=q}^{\infty} \left( 1 - \zeta_i^{a_k} \prod_{j=1}^s (y_{j1}^{a_{k+n-1}} \dots y_{jn}^{a_k})^{e_{ij}} \right) \quad (1 \leq i \leq r).$$

Letting

$$\beta = (\underbrace{1, \dots, 1}_{n-1}, \beta_1, \dots, \dots, \underbrace{1, \dots, 1}_{n-1}, \beta_s),$$

we see by (3) that

$$f_i(\beta) = \sum_{k=q}^{\infty} \alpha_i^{a_k}, \quad g_i(\beta) = \sum_{k=q}^{\infty} \frac{\alpha_i^{a_k}}{1 - \alpha_i^{a_k}}, \quad h_i(\beta) = \prod_{k=q}^{\infty} (1 - \alpha_i^{a_k}).$$

Hence the values  $f_1(\beta), \dots, f_\rho(\beta), g_1(\beta), \dots, g_r(\beta), h_1(\beta), \dots, h_r(\beta)$  are algebraically dependent. Let  $\Omega$  be a multiplicative transformation for the variables  $y_{11}, \dots, y_{1n}, \dots, y_{s1}, \dots, y_{sn}$  sending  $y_{j1}^{a_{k+n-1}} \dots y_{jn}^{a_k}$  to  $y_{j1}^{a_{k+p+n-1}} \dots y_{jn}^{a_{k+p}}$  for  $j = 1, \dots, s$ . Then  $f_1(\mathbf{y}), \dots, f_\rho(\mathbf{y}), g_1(\mathbf{y}), \dots, g_r(\mathbf{y}), h_1(\mathbf{y}), \dots, h_r(\mathbf{y})$  satisfy the functional equations

$$\begin{aligned} f_i(\mathbf{y}) &= f_i(\Omega\mathbf{y}) + \sum_{k=q}^{p+q-1} \zeta_i^{a_k} \prod_{j=1}^s (y_{j1}^{a_{k+n-1}} \dots y_{jn}^{a_k})^{e_{ij}}, \\ g_i(\mathbf{y}) &= g_i(\Omega\mathbf{y}) + \sum_{k=q}^{p+q-1} \frac{\zeta_i^{a_k} \prod_{j=1}^s (y_{j1}^{a_{k+n-1}} \dots y_{jn}^{a_k})^{e_{ij}}}{1 - \zeta_i^{a_k} \prod_{j=1}^s (y_{j1}^{a_{k+n-1}} \dots y_{jn}^{a_k})^{e_{ij}}}, \end{aligned}$$

and

$$h_i(\mathbf{y}) = \left( \prod_{k=q}^{p+q-1} \left( 1 - \zeta_i^{a_k} \prod_{j=1}^s (y_{j1}^{a_{k+n-1}} \dots y_{jn}^{a_k})^{e_{ij}} \right) \right) h_i(\Omega\mathbf{y}),$$

since  $a_{k+p} \equiv a_k \pmod{N}$  for any  $k \geq q$ . By Mahler's method improved by Kubota [1], at least one of the following two cases arises:

- (i) There are algebraic numbers  $b_1, \dots, b_\rho, c_1, \dots, c_r$ , not all zero, and  $F(\mathbf{y}) \in \overline{\mathbb{Q}}(\mathbf{y})$  such that

$$\begin{aligned} F(\mathbf{y}) &= F(\Omega\mathbf{y}) + \sum_{k=q}^{p+q-1} \left( \sum_{i=1}^{\rho} b_i \zeta_i^{a_k} \prod_{j=1}^s (y_{j1}^{a_{k+n-1}} \dots y_{jn}^{a_k})^{e_{ij}} \right. \\ &\quad \left. + \sum_{i=1}^r \frac{c_i \zeta_i^{a_k} \prod_{j=1}^s (y_{j1}^{a_{k+n-1}} \dots y_{jn}^{a_k})^{e_{ij}}}{1 - \zeta_i^{a_k} \prod_{j=1}^s (y_{j1}^{a_{k+n-1}} \dots y_{jn}^{a_k})^{e_{ij}}} \right). \end{aligned} \quad (4)$$

- (ii) There are rational integers  $d_1, \dots, d_r$ , not all zero, and  $G(\mathbf{y}) \in \overline{\mathbb{Q}}(\mathbf{y}) \setminus \{0\}$  such that

$$G(\mathbf{y}) = \left( \prod_{k=q}^{p+q-1} \prod_{i=1}^r \left( 1 - \zeta_i^{a_k} \prod_{j=1}^s (y_{j1}^{a_{k+n-1}} \dots y_{jn}^{a_k})^{e_{ij}} \right)^{d_i} \right) G(\Omega\mathbf{y}). \quad (5)$$

Let  $M > 0$  be a sufficiently large integer and let

$$\begin{aligned} F^*(z) &= F(z_1^M, \dots, z_n^M, \dots, z_1^{M^s}, \dots, z_n^{M^s}) \in \overline{\mathbb{Q}}(z_1, \dots, z_n), \\ G^*(z) &= G(z_1^M, \dots, z_n^M, \dots, z_1^{M^s}, \dots, z_n^{M^s}) \in \overline{\mathbb{Q}}(z_1, \dots, z_n) \setminus \{0\}. \end{aligned}$$

Then by (4) and (5), at least one of the following two functional equations holds:

$$\begin{aligned} F^*(z) &= F^*(\Omega z) \\ &+ \sum_{k=q}^{p+q-1} \left( \sum_{i=1}^{\rho} b_i \zeta_i^{a_k} (z_1^{a_{k+n-1}} \dots z_n^{a_k})^{E_i} + \sum_{i=1}^r \frac{c_i \zeta_i^{a_k} (z_1^{a_{k+n-1}} \dots z_n^{a_k})^{E_i}}{1 - \zeta_i^{a_k} (z_1^{a_{k+n-1}} \dots z_n^{a_k})^{E_i}} \right), \end{aligned} \quad (6)$$

$$G^*(z) = \left( \prod_{k=q}^{p+q-1} \prod_{i=1}^r \left( 1 - \zeta_i^{a_k} (z_1^{a_{k+n-1}} \dots z_n^{a_k})^{E_i} \right)^{d_i} \right) G^*(\Omega z), \quad (7)$$

where  $\Omega$  sends  $z_1^{a_{k+n-1}} \dots z_n^{a_k}$  to  $z_1^{a_{k+p+n-1}} \dots z_n^{a_{k+p}}$  and  $E_i = \sum_{j=1}^s e_{ij} M^j > 0$  ( $1 \leq i \leq r$ ) such that  $E_i \neq E_{i'}$  if  $\alpha_i/\alpha_{i'}$  is not a root of unity, or equivalently  $(e_{i1}, \dots, e_{is}) \neq (e_{i'1}, \dots, e_{i's})$ . By Theorems 1 and 2 of [7], at least one of the following two properties are satisfied:

(i) For any  $k$  ( $q \leq k \leq p+q-1$ ),

$$\begin{aligned} \sum_{i=1}^{\rho} b_i \zeta_i^{a_k} X^{E_i} + \sum_{i=1}^r \frac{c_i \zeta_i^{a_k} X^{E_i}}{1 - \zeta_i^{a_k} X^{E_i}} &= \sum_{i=1}^{\rho} b_i \zeta_i^{a_k} X^{E_i} + \sum_{i=1}^r c_i \sum_{h=1}^{\infty} (\zeta_i^{a_k} X^{E_i})^h \\ &\in \overline{\mathbb{Q}}. \end{aligned} \quad (8)$$

(ii) For any  $k$  ( $q \leq k \leq p+q-1$ ),

$$\prod_{i=1}^r (1 - \zeta_i^{a_k} X^{E_i})^{d_i} = \gamma_k \in \overline{\mathbb{Q}}^\times. \quad (9)$$

If (6) is satisfied, then all the coefficients of the right-hand side of (8) must be zero. Therefore, if  $c_i = 0$  ( $1 \leq i \leq r$ ), then  $\alpha_1, \dots, \alpha_\rho$  are  $\{a_k\}_{k \geq 0}$ -dependent, which contradicts the assumption. If  $c_1, \dots, c_r$  are not all zero, then  $\alpha_1, \dots, \alpha_r$  are strongly  $\{a_k\}_{k \geq 0}$ -dependent, which also contradicts the assumption.

If (7) is satisfied, taking the logarithmic derivative of (9), we get

$$\sum_{i=1}^r \frac{-d_i E_i \zeta_i^{a_k} X^{E_i-1}}{1 - \zeta_i^{a_k} X^{E_i}} = 0 \quad (q \leq k \leq p+q-1)$$



and so

$$\sum_{i=1}^r \frac{d_i E_i \zeta_i^{a_k} X^{E_i}}{1 - \zeta_i^{a_k} X^{E_i}} = \sum_{i=1}^r d_i E_i \sum_{h=1}^{\infty} (\zeta_i^{a_k} X^{E_i})^h = 0 \quad (q \leq k \leq p + q - 1).$$

Therefore  $\alpha_1, \dots, \alpha_r$  are strongly  $\{a_k\}_{k \geq 0}$ -dependent in this case by the same way as above. This completes the proof of the theorem.

## References

- [1] K.K. Kubota: *On the algebraic independence of holomorphic solutions of certain functional equations and their values*, Math. Ann. **227** (1977), 9–50.
- [2] J. H. Loxton and A. J. van der Poorten: *Algebraic independence properties of the Fredholm series*, J. Austral. Math. Soc. Ser. A **26** (1978), 31–45.
- [3] K. Mahler: *Arithmetische Eigenschaften der Lösungen einer Klasse von Funktionalgleichungen*, Math. Ann. **101** (1929), 342–366.
- [4] K. Nishioka, *Conditions for algebraic independence of certain power series of algebraic numbers*, Compositio Math. **62** (1987), 53–61.
- [5] K. Nishioka: *Mahler functions and transcendence*, Lecture Notes in Mathematics No. **1631**, Springer, 1996.
- [6] T. Tanaka: *Algebraic independence of the values of power series generated by linear recurrences*, Acta Arith. **74** (1996), 177–190.
- [7] T. Tanaka: *Algebraic independence results related to linear recurrences*, Osaka J. Math. **36** (1999), 203–227.
- [8] T. Tanaka: *Algebraic independence of the values of power series, Lambert series, and infinite products generated by linear recurrences*, Osaka J. Math., to appear.