

On holomorphic framed vertex operator algebras of rank 24

Ching Hung Lam

Department of Mathematics, National Cheng Kung University,

Tainan, Taiwan 701

e-mail: `chlam@mail.ncku.edu.tw`

1 Framed vertex operator algebra

This article is based on an on-going research project with H. Yamauchi and H. Shimakura. We shall discuss our ideas on how to classify holomorphic framed vertex operator algebras of rank 24 using the structure codes and \mathbb{Z}_2 -orbifold constructions.

First, we shall review the notion of a framed vertex operator algebra.

Definition 1.1. A Virasoro vector e is called an *Ising vector* if the subalgebra $\text{Vir}(e) \simeq L(1/2, 0)$. Two Virasoro vectors $u, v \in V$ are called *orthogonal* if $[Y(u, z_1), Y(v, z_2)] = 0$. A decomposition $\omega = e^1 + \dots + e^n$ of the conformal vector ω of V is called *orthogonal* if e^i are mutually orthogonal Virasoro vectors.

Remark 1.2. It is well-known that $L(1/2, 0)$ is rational, C_2 -cofinite and has three irreducible modules $L(1/2, 0)$, $L(1/2, 1/2)$ and $L(1/2, 1/16)$. The fusion rules of $L(1/2, 0)$ -modules are computed in [DMZ]:

$$\begin{aligned} L(1/2, 1/2) \boxtimes L(1/2, 1/2) &= L(1/2, 0), & L(1/2, 1/2) \boxtimes L(1/2, 1/16) &= L(1/2, 1/16), \\ L(1/2, 1/16) \boxtimes L(1/2, 1/16) &= L(1/2, 0) \oplus L(1/2, 1/2). \end{aligned} \tag{1.1}$$

Definition 1.3. ([DGH, M3]) A simple vertex operator algebra (V, ω) is called *framed* if there exists a set $\{e^1, \dots, e^n\}$ of Ising vectors of V such that $\omega = e^1 + \dots + e^n$ is an orthogonal decomposition. The full sub VOA F generated by e^1, \dots, e^n is called an *Ising frame* or simply a *frame* of V . By abuse of notation, we sometimes call the set of Ising vectors $\{e^1, \dots, e^n\}$ a *frame*, also.

Give a framed VOA V with a frame T , one can associate two binary codes C and D to V and T as follows:

Since $T = L(1/2, 0)^{\otimes n}$ is a rational vertex operator algebra, V is a completely reducible T -module. That is,

$$V \cong \bigoplus_{h_i \in \{0, \frac{1}{2}, \frac{1}{16}\}} m_{h_1, \dots, h_n} L(h_1, \dots, h_n),$$

where the nonnegative integer m_{h_1, \dots, h_n} is the multiplicity of $L(h_1, \dots, h_n)$ in V . In particular, all the multiplicities are finite and m_{h_1, \dots, h_n} is at most 1 if all h_i are different from $\frac{1}{16}$.

Let $L = L(1/2, h_1) \otimes \dots \otimes L(1/2, h_n)$ be an irreducible module for T . The τ -word $\tau(L)$ of L is a binary word $\beta = (\beta_1, \dots, \beta_n) \in \mathbb{Z}_2^n$ such that

$$\beta_i = \begin{cases} 0 & \text{if } h_i = 0 \text{ or } 1/2, \\ 1 & \text{if } h_i = 1/16. \end{cases} \quad (1.2)$$

For any $\beta \in \mathbb{Z}_2^n$, define V^β as the sum of all irreducible submodules L of V such that $\tau(L) = \beta$. Denote $D := \{\beta \in \mathbb{Z}_2^n \mid V^\beta \neq 0\}$. Then D is an even linear subcode of \mathbb{Z}_2^n and $V = \bigoplus_{\beta \in D} V^\beta$.

For any $c = (c_1, \dots, c_n) \in \mathbb{Z}_2^n$, denote $V(c) = m_{h_1, \dots, h_n} L(h_1, \dots, h_n)$ where $h_i = \frac{1}{2}$ if $c_i = 1$ and $h_i = 0$ elsewhere. Set

$$C := \{c \in \mathbb{Z}_2^n \mid V(c) \neq 0\}.$$

Then $V^0 = \bigoplus_{c \in C} V(c) \neq 0$.

Summarizing, there exists a pair (C, D) of even linear codes such that V is an D -graded extension of a code VOA V_C associated to C . We call the pair (C, D) the *structure codes* of a framed VOA V associated with the frame F . Since the powers of z in an $L(1/2, 0)$ -intertwining operator of type $L(1/2, 1/2) \times L(1/2, 1/2) \rightarrow L(1/2, 1/16)$ are half-integral, the structure codes (C, D) satisfy $C \subset D^\perp$. Moreover, it is known [DGH, M3] that V is holomorphic if and only if $C = D^\perp$.

Remark 1.4. Let V be a framed VOA with the structure codes (C, D) , where $C, D \subset \mathbb{Z}_2^n$. For a binary codeword $\beta \in \mathbb{Z}_2^n$, we define

$$\tau_\beta(u) := (-1)^{\langle \alpha, \beta \rangle} u \quad \text{for } u \in V^\alpha. \quad (1.3)$$

Then by the fusion rules, τ_β defines an automorphism on V [M1]. Note that the subgroup $P = \{\tau_\beta \mid \beta \in \mathbb{Z}_2^n\}$ is an elementary abelian 2-group and is

isomorphic to \mathbb{Z}_2^n/D^\perp . In addition, the fixed point subspace V^P is equal to V^0 and all $V^\alpha, \alpha \in D$ are irreducible V^0 -modules. Similarly, we can define an automorphism on V^0 by

$$\sigma_\beta(u) := (-1)^{\langle \alpha, \beta \rangle} u \quad \text{for } u \in V(\alpha),$$

where $V^0 = \bigoplus_{\alpha \in C} V(\alpha)$. Note that the group $Q = \{\sigma_\beta \mid \beta \in \mathbb{Z}_2^n\} \cong \mathbb{Z}_2^n/C^\perp$ is elementary abelian and $(V^0)^Q = V(0)$.

The following theorem is very important to our argument and is proved in [LY]

Theorem 1.5. *Let $V = \bigoplus_{\alpha \in D} V^\alpha$ be a framed VOA with structure codes (C, D) . Then*

1. *For every non-zero $\alpha \in D$, the subcode C_α of C contains a doubly even self-dual subcode w.r.t. α .*
2. *C is even, every codeword of D has a weight divisible by 8, and $D \subset C \subset D^\perp$.*

As a corollary, the following theorem is also proved in [LY].

Theorem 1.6. *Let $V = \bigoplus_{\alpha \in D} V^\alpha$ be a framed VOA with structure codes (C, D) . Then $V = \bigoplus_{\alpha \in D} V^\alpha$ is a D -graded simple current extension of the code VOA $V^0 = V_C$.*

The following corollaries follow immediately by the standard arguments for simple current extensions.

Corollary 1.7 ([LY, Y2]). *Let $V = \bigoplus_{\alpha \in D} V^\alpha$ be a framed VOA with structure codes (C, D) . Let W be an irreducible V^0 -module. Then there exists $\eta \in \mathbb{Z}_2^n$, which is unique modulo D^\perp , such that W can be uniquely extended to an irreducible τ_η -twisted V -module which is given by $V \boxtimes_{V^0} W$ as a V^0 -module. In particular, every irreducible untwisted V -module is D -stable.*

Corollary 1.8 ([L1, Y2]). *Let $V = \bigoplus_{\alpha \in D} V^\alpha$ be a holomorphic framed VOA with structure codes (C, D) . For any $\delta \in \mathbb{Z}_2^n$, denote*

$$D^0 = \{\alpha \in D \mid \langle \alpha, \delta \rangle = 0\} \quad \text{and} \quad D^1 = \{\alpha \in D \mid \langle \alpha, \delta \rangle \neq 0\}.$$

Define

$$V(\tau_\delta) = \begin{cases} \left(\bigoplus_{\alpha \in D^0} V^\alpha \right) \oplus \left(\bigoplus_{\alpha \in D^1} M_{\delta+C} \times_{M_C} V^\alpha \right) & \text{if wt } \delta \text{ is odd,} \\ \left(\bigoplus_{\alpha \in D^0} V^\alpha \right) \oplus \left(\bigoplus_{\alpha \in D^0} M_{\delta+C} \times_{M_C} V^\alpha \right) & \text{if wt } \delta \text{ is even.} \end{cases}$$

Then $V(\tau_\delta)$ is also a holomorphic framed VOA. Moreover, the structure codes of $V(\tau_\delta)$ are given by (C, D) if wt δ is odd and $(C \cup (\delta + C), D^0)$ if wt δ is even.

The construction of $V(\tau_\delta)$ is often referred as to a \mathbb{Z}_2 -orbifold construction.

2 Structure codes for holomorphic framed VOAs

If V is a holomorphic framed VOA with the structure codes (C, D) , then C will satisfy the following conditions:

1. The length of C is divisible by 16.
2. C is even, every codeword of C^\perp has a weight divisible by 8, and $C^\perp \subset C$.
3. For any $\alpha \in C^\perp$, the subcode C_α of C contains a doubly even self-dual subcode w.r.t. α .

For simplicity, we shall call a code C *F-admissible* if it satisfies the above conditions (1)- (3). Indeed, one can construct a holomorphic framed VOA starting from an *F-admissible* code.

Theorem 2.1 ([LY]). *There exists a holomorphic framed VOA with structure codes (C, C^\perp) if and only if C is F-admissible, i.e., C satisfies conditions (1)-(3) above.*

Remark 2.2. A linear code C is *F-admissible* if and only if its dual C^\perp satisfies the following three conditions:

- (i) the length of C^\perp is divisible by 16,
- (ii) C^\perp contains the all-one vector,
- (iii) C^\perp is *triply even*, that is, wt(α) is divisible by 8 for any $\alpha \in C^\perp$.

For, let D satisfy the conditions (i), (ii) and (iii) above. Then for any $\alpha, \beta \in D$, the weight of their intersection $\alpha \cdot \beta$ is divisible by 4 and so $\alpha \cdot D$ is doubly even. Then there exists a doubly even code E containing $\alpha \cdot D$ such that E is self-dual w.r.t. α . For any $\delta \in (\alpha \cdot D)^\perp$, we have $\langle \delta, D \rangle = \langle \delta \cdot \alpha, D \rangle = \langle \delta, \alpha \cdot D \rangle = 0$, showing $E \subset (\alpha \cdot D)^\perp \subset (D^\perp)_\alpha$. Therefore, D^\perp is *F-admissible*.

Now let us consider some examples of *F-admissible* codes.

2.1 \mathbb{Z}_4 codes and framed VOAs

Let Z be a self-orthogonal linear code over \mathbb{Z}_4 . Define

$$A_4(Z) = \frac{1}{2} \{(x_1, \dots, x_n) \in \mathbb{Z}^n \mid (x_1, \dots, x_n) \in Z \pmod{4}\}.$$

Then $A_4(Z)$ is an even lattice. It is also well-known that $A_4(Z)$ is unimodular iff Z is self-dual. Note that if $Z = 0$, then $A_4(Z) \cong \sqrt{2}A_1^n$. Note that the lattice VOA $V_{\sqrt{2}A_1}$ is a framed VOA (cf. [DMZ, M2]) and

$$V_{\sqrt{2}A_1} \cong L\left(\frac{1}{2}, 0\right) \otimes L\left(\frac{1}{2}, 0\right) \oplus L\left(\frac{1}{2}, \frac{1}{2}\right) \otimes L\left(\frac{1}{2}, \frac{1}{2}\right)$$

as an $L(\frac{1}{2}, 0) \otimes L(\frac{1}{2}, 0)$ -module. Hence the lattice VOA $V_{A_4(Z)}$ is framed for any Z .

The positive definite even unimodular lattices of rank 24 have been classified by Niemeier. There are exactly 24 such lattices and they are characterized by their root systems. The following theorem by Kitazume-Harada shows that all positive definite even unimodular lattices of rank 24 can be constructed by \mathbb{Z}_4 -codes.

Theorem 2.3 (Kitazume-Harada). *Let N be the Leech lattice or a Niemeier lattice. Then there exists a self-dual \mathbb{Z}_4 code Z such that $N \cong A_4(Z)$. In particular, the lattice VOA V_N is framed.*

Now let us study the structure codes for the lattice VOA V_N .

Let Z be a self-dual \mathbb{Z}_4 code. Denote

$$\begin{aligned} Z_0 &= \{(\alpha_1, \dots, \alpha_n) \in \{0, 1\}^n \mid (2\alpha_1, \dots, 2\alpha_n) \in Z\}, \\ Z_1 &= \{\alpha \in \{0, 1\}^n \mid \alpha \equiv \beta \pmod{2} \text{ for some } \beta \in Z\}. \end{aligned}$$

Then both Z_0 and Z_1 are even binary codes. Moreover, Z_1 is doubly even and $Z_0^\perp = Z_1$.

Proposition 2.4. *Let Z be a self-dual linear \mathbb{Z}_4 -code and Z_0 and Z_1 defined as above. Then the structure codes of the lattice VOA $V = V_{A_4(Z)}$ are given by*

$$D = d(Z_1) \quad \text{and} \quad C = D^\perp,$$

where $d : \mathbb{Z}_2^n \rightarrow \mathbb{Z}_2^{2n}$ is given by $d(a_1, a_2, \dots, a_n) = (a_1, a_1, a_2, a_2, \dots, a_n, a_n)$. Note that the code $\{(0, 0), (1, 1)\}^n$ is contained in C .

Let $\delta = (10)^n$. Then τ_δ defines an automorphism on $V_{A_4(Z)}$. In fact, τ_δ is conjugate to the automorphism θ , which is the lift of the -1 -map on the lattice $N = A_4(Z)$.

By Corollary 1.8, the structure codes for the τ_δ -twisted orbifold

$$V(\tau_\delta) = \left(\bigoplus_{\alpha \in D^0} V^\alpha \right) \oplus \left(\bigoplus_{\alpha \in D^0} M_{\delta+C} \times_{M_C} V^\alpha \right)$$

are given by $D = \langle d(Z_1), (10)^n \rangle$ and $C = D^\perp$. Note that C contains the code E^+ generated by

11110000...
00111100...
00001111...
0000001111...

We believe that this case is the typical case and the following holds.

Conjecture 1. Let D be a indecomposable triply even binary code of length $16k$. Then D is a subcode of $\langle d(C), (01)^{8k} \rangle$, where C is a double even self-dual codes of length $8k$.

A code is said to be *indecomposable* if there does not a partition $I \cup J = \{1, \dots, n\}$ such that $I \cap J = \emptyset$ but $D = D_1 \oplus D_2$ with $\text{supp } D_1 \subset I$ and $\text{supp } D_2 \subset J$.

The conjecture actually holds for $k = 1, 2$ but is not proved for $k \geq 3$. If the conjecture also holds for $k = 3$, then we have the following classification of holomorphic framed VOAs of rank 24.

Theorem 2.5. *Assume that Conjecture 1 holds for $k = 3$. If V is holomorphic framed VOA of rank 24, then V is isomorphic to one of the following:*

- (1) a lattice VOA V_N , or
- (2) the θ -orbifold of a lattice VOA V_N ,

where N is the Leech lattice or a Niemeier lattice.

In particular, V can be characterized by its weight 1 subspace V_1 .

Remark 2.6. Note that if $N \subset A_1^{24}$, then τ_δ -orbifold is again a lattice VOA. There are 9 such cases and hence there are exactly 15 holomorphic framed VOAs of rank 24, which are not lattice VOAs. Therefore, there are totally 39 holomorphic framed VOAs of rank 24 – 24 lattice VOAs and 15 θ -twisted orbifolds.

Sketch of the proof

Let (C, D) be the structure codes of V . If the conjecture holds, then $C \supset E^+$, which is generated by

$$\begin{aligned} &11110000\dots \\ &00111100\dots \\ &00001111\dots \\ &0000001111\dots \end{aligned}$$

If C contains $\{(00), (11)\}^{24}$, then V contains a subVOA isomorphic to $V_{\sqrt{2}A_1}^{\otimes 24}$ and hence V is isomorphic to a lattice VOA associated with a Niemeier lattice.

Otherwise, let $\alpha = (1100\dots 0)$. Then $\alpha \notin C$ and the τ_α -orbifold of V will have the structure code containing $\{(00), (11)\}^{24}$ and hence the τ_δ -orbifold $V(\tau_\delta)$ is isomorphic to a lattice VOA V_N . By reversing the orbifold construction, one can show that V is a θ -twisted orbifold of V_N .

References

- [DGH] C. Dong, R.L. Griess and G. Höhn, Framed vertex operator algebras, codes and the moonshine module, *Comm. Math. Phys.* **193** (1998), 407–448.
- [DL] C. Dong and J. Lepowsky, Generalized vertex algebras and relative vertex operators, *Progress in Math.* **112**, Birkhäuser, Boston, 1993.
- [DLM3] C. Dong, H. Li and G. Mason, Modular-invariance of trace functions in orbifold theory and generalized moonshine, *Comm. Math. Phys.* **214** (2000), 1–56.
- [DLM4] C. Dong, H. Li and G. Mason, Simple currents and extensions of vertex operator algebras, *Comm. Math. Phys.* **180**, 671–707.
- [DM1] C. Dong and G. Mason, On quantum Galois theory, *Duke Math. J.* **86** (1997), 305–321.
- [DM2] C. Dong and G. Mason, Rational vertex operator algebras and the effective central charge, *Internat. Math. Res. Notices* **56** (2004), 2989–3008.
- [DMZ] C. Dong, G. Mason and Y. Zhu, Discrete series of the Virasoro algebra and the moonshine module, *Proc. Symp. Pure. Math.*, American Math. Soc. **56** II (1994), 295–316.
- [FHL] I. Frenkel, Y.-Z. Huang and J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, *Memoirs Amer. Math. Soc.* **104**, 1993.

- [FLM] I.B. Frenkel, J. Lepowsky and A. Meurman, Vertex Operator Algebras and the Monster, Academic Press, New York, 1988.
- [GKO] P. Goddard, A. Kent and D. Olive, Unitary representations of the Virasoro and super-Virasoro algebras, *Comm. Math. Phys.* **103** (1986), 105–119.
- [H1] Y.-Z. Huang, Virasoro vertex operator algebras, (non-meromorphic) operator product expansion and the tensor product theory, *J. Algebra* **182** (1996), 201–234.
- [L1] C.H. Lam, Twisted representations of code vertex operator algebras, *J. Algebra* **217** (1999), 275–299.
- [L2] C.H. Lam, Some twisted module for framed vertex operator algebras, *J. Algebra* **231** (2000), 331–341.
- [LY] C.H. Lam and H. Yamauchi, On the structure framed vertex operator algebras and their pointwise framed stabilizers, preprint.
- [McS] F.J. MacWilliams and N.J.A Sloane, The theory of error-correcting codes, North-Holland, 1998.
- [McST] F.J. MacWilliams, N.J.A Sloane and J.G. Thompson, Good self-dual codes exist, *Discrete Mathematics* **3** (1972), 153–162.
- [M1] M. Miyamoto, Griess algebras and conformal vectors in vertex operator algebras, *J. Algebra* **179** (1996), 528–548.
- [M2] M. Miyamoto, Representation theory of code vertex operator algebras, *J. Algebra* **201** (1998), 115–150.
- [M3] M. Miyamoto, A new construction of the moonshine vertex operator algebra over the real number field, *Ann. of Math* **159** (2004), 535–596.
- [Sh] H. Shimakura, Automorphism group of the vertex operator algebra V_L^+ for an even lattice L without roots, *J. Algebra* **280** (2004), 29–57.
- [Y1] H. Yamauchi, Module categories of simple current extensions of vertex operator algebras, *J. Pure Appl. Algebra* **189** (2004), 315–328.
- [Y2] H. Yamauchi, A theory of simple current extensions of vertex operator algebras and applications to the moonshine vertex operator algebra, Ph.D. thesis, University of Tsukuba, 2004.; available on the author's web site: <http://www.ms.u-tokyo.ac.jp/~yamauchi>
- [Y3] H. Yamauchi, 2A-orbifold construction and the baby-monster vertex operator superalgebra, *J. Algebra* **284** (2005), 645–668.
- [Z] Y.Zhu, Modular invariance of characters of vertex operator algebras, *J. Amer. Math. Soc.* **9** (1996), 237–302.