

# Lyapounov exponents and meromorphic maps

Henry de Thélin  
Université de Paris-Sud

Complex Dynamics and Related Topics

Research Institute for Mathematical Sciences, Kyoto University

September 3–6, 2007

# Lyapounov exponents and meromorphic maps

①

$(X, \omega)$  compact Kähler manifold of dimension  $k$ .

$f: X \rightarrow X$  dominating meromorphic map

$I_f =$  the indeterminacy set of  $f$ .

$C_f =$  the critical set of  $f$ .

## I) Dynamical quantities:

1) Dynamical degrees

(Rusakovskii-Schiffman)  
for  $X = \mathbb{C}P^k$

$$\omega^l = \underbrace{\omega \wedge \dots \wedge \omega}_{l \text{ times}} \quad 0 \leq l \leq k = \dim X$$

$f^*(\omega^l)$  form with  $L^2_{loc}$  coefficients

$$S_e(f) = \int f^*(\omega^l) \wedge \omega^{k-l}$$

$$d_e = \lim_{n \rightarrow +\infty} |S_e(f^n)|^{1/n} \leftarrow \text{this limit exists (Dinh-Sibony)}$$

$d_e$  dynamical degree

Th (Kovanskii - Teissier - Gromov)

(2)

$q \rightarrow \log d_q$  is concave.

It implies that the dynamical degrees look like:

$$d_0 = 1 \leq d_1 \leq \dots \leq d_s \geq d_{s+1} \geq \dots \geq d_k$$

Examples:

•  $f$  holomorphic endomorphism of  $\mathbb{C}P^k$  of degree  $d \geq 2$

$$d_0 = 1 < d_1 = d < d_2 = d^2 < \dots < d_k = d^k$$

•  $f$  birational map of  $\mathbb{C}P^2$  of degree  $d \geq 2$   
 $f$  algebraically stable

$$d_0 = 1 < d_1 = d > d_2 = 1$$

3


2) Entropy:

a) Topological entropy:

$$d_n(x, y) := \max_{0 \leq i \leq n-1} d(f^i(x), f^i(y))$$

$F$  is a  $(n, \delta)$ -separated set if

$$\forall x, y \in F \quad x \neq y \Rightarrow d_n(x, y) \geq \delta.$$



$$h_{\text{top}}(f) := \sup_{\delta > 0} \lim_{n \rightarrow \infty} \frac{1}{n} \log \max \{ \text{Card } F, F \text{ } (n, \delta)\text{-separated set } \}$$

↑  
topological  
entropy

Th (Gromov: holomorphic case

Dinh-Sibony: meromorphic case)

$$h_{\text{top}}(f) \leq \max_{0 \leq p \leq k} \log d_p \leftarrow \text{dynamical degrees}$$

(4)

b) Metric entropy:

$\mu$  measure  $\mu(I_f) = 0$   
 $f_*\mu = \mu$  ( $\mu$  is invariant)

$B_n(x, \delta)$  = ball with center  $x$  and  
 radius  $\delta$  for the metric  $d_n$

$$h_\mu(f) := \sup_{\delta > 0} \lim_{n \rightarrow \infty} -\frac{1}{n} \log \mu B_n(x, \delta)$$

metric  
 entropy

$x$  generic for  $\mu$

(Brin-Katok)

Fact:  $h_\mu(f) \leq h_{\text{top}}(f)$ .

## II) Meromorphic maps:

⑤

$f: X \rightarrow X$  dominating meromorphic map  
 ( $X$  w/ compact Kähler manifold dimension  $k$ )

$$d_0 \leq d_1 \leq \dots \leq d_{s-1} < d_s > d_{s+1} \geq \dots \geq d_k$$

Example: holomorphic endomorphisms of  $\mathbb{C}P^k$   
 $d_p = d^p$  of degree  $d \geq 2$   
 $s = k$  in this case.

We consider a measure  $\mu$  invariant, ergodic  
 with  $\log d(\cdot, \mathbb{I}_f \cup \mathbb{C}_f) \in L^1(\mu)$  also

$$\mathbb{R} \quad \uparrow$$

necessary condition  
 for the existence  
 of the Lyapounov  
 exponents

$$+\infty > \chi_1 \geq \dots \geq \chi_k > -\infty$$

$$\underline{h}_p(\mu) \leq \underline{h}_{\text{top}}(\mu) \leq \log d_s$$

Dinh-Sibony

Th (2.1)

(6)

If  $h_\mu(f) > \max(\log d_{s-1}, \log d_{s+1})$

(or  $h_\mu(f) > \log d_{k-1}$  if  $s=k$ )

Then:

$$x_1 \geq \dots \geq x_s \geq \frac{1}{2} (h_\mu(f) - \log d_{s-1}) > 0$$

$$0 > \frac{1}{2} (\log d_{s+1} - h_\mu(f)) \geq x_{s+1} \geq \dots \geq x_k$$

In particular  $\mu$  is hyperbolic.

$x_1 \geq \dots \geq x_s > 0 \rightsquigarrow s$  directions with expansion  
 $0 > x_{s+1} \geq \dots \geq x_k \rightsquigarrow k-s$  directions with contraction.

Cor:

If  $h_\mu(f) = \log d_s$

$$x_1 \geq \dots \geq x_s \geq \frac{1}{2} \log \frac{d_s}{d_{s-1}} > 0$$

~~0 > \frac{1}{2} \log \frac{d\_{s+1}}{d\_s} \geq x\_{s+1} \geq \dots \geq x\_k~~

$$0 > \frac{1}{2} \log \frac{d_{s+1}}{d_s} \geq x_{s+1} \geq \dots \geq x_k$$

Ab: These bounds are sharp.

Examples of dynamical systems and measure  $\mu$  which satisfy the hypothesis of the corollary: (7)

- Holomorphic endomorphisms of  $\mathbb{C}P^k$  of degree  $d \geq 2$

$$d_0 = 1 < d_1 = d < \dots < d_k = d^k$$

$\mu =$  Green measure (Fornaess - Sibony)  
 $\mu$  is invariant and ergodic,  $\log d(x, c_j) \in L^1(\mu)$

$a =$  a generic point

$$\frac{1}{d^{kn}} \sum_{f^n(a) = a} \delta_{a_i} \rightarrow \mu$$

(Dinh-Sibony  
Briend-Duval)

$d^k =$  topological degree of  $f$

Th: Gromov / Misiurewicz - Przytycki:

$$h_\mu(f) = k \log d = \log d^k = \log d_k.$$

we can apply the corollary

$$\Rightarrow \chi_1 \geq \dots \geq \chi_k \geq \frac{1}{2} \log \frac{d^k}{d^{k-1}} = \frac{1}{2} \log d$$

$\rightarrow$  we found the Briend-Duval's inequality.



- $X$  projective ~~manifold~~ manifold with dimension  $k$  ⑧

$f: X \rightarrow X$  dominating meromorphic map  
with  $d_k > d_{k-1} \geq \dots \geq d_0 = 1$

Guedj constructed an invariant ergodic measure  $\mu$  with  $\log d(x, \mathbb{R}) \in L^2(\mu)$   
and  $h_\mu(f) = \log d_k$ .

Corollary  $(S=k) \rightsquigarrow \chi_1 \geq \dots \geq \chi_k \geq \frac{1}{2} \log \frac{d_k}{d_{k-1}} > 0$

$\rightsquigarrow$  this is the Guedj's inequality.

- $f$  regular birational map of  $\mathbb{C}P^k$   
(Dinh-Sibony)

They constructed an invariant ergodic measure  $\mu$  with  $\log d(x, \mathbb{R}) \in L^1(\mu)$   
and with maximal entropy

$\rightsquigarrow \mu$  is hyperbolic.  
Cor.

- other examples: holomorphic automorphisms in Kähler manifolds (Dinh-Sibony)....

### III) A general inequality:

( $d_0 \leq d_1 \leq \dots \leq d_s \leq \dots \leq d_k$ )

(9)

#### Th (D.1)

Let  $\mu$  be an invariant ergodic measure

$$\log d(x, \mathcal{R}) \in L^2(\mu).$$

$\lambda_1 \geq \dots \geq \lambda_k$  the Lyapounov exponents

Fix  $s$   $1 \leq s \leq k$  and we define

$$p = |s| \quad \text{with:}$$

$$p' = p'(|s|)$$

$$\lambda_1 \geq \dots \geq \lambda_{s-p-1} > \lambda_{s-p} = \dots = \lambda_s = \dots = \lambda_{s+p'} > \lambda_{s+p'+1} \geq \dots \geq \lambda_k$$

$$(\text{with } s-p=1 \text{ if } \lambda_1 = \dots = \lambda_s$$

$$\text{and } s+p'=k \text{ if } \lambda_s = \dots = \lambda_k)$$

Then

$$h_\mu^+(f) \leq \max_{0 \leq q \leq s-p-1} \log d_q + 2X_{s-p}^+ + \dots + 2X_k^+$$

$$X_i^+ = \max(X_i, 0)$$

$$h_\mu^-(f) \leq \max_{s+p' \leq q \leq k} \log d_q - 2X_1^- - \dots - 2X_{s+p'}^-.$$

$$X_i^- = \min(X_i, 0).$$

⑨'

Cor 1:

Let  $\mu$  be an invariant ergodic measure  
 $\log d(\alpha, \mathbb{R}) \in L^1(\mu)$

$$h_\mu(f) \leq 2X_1^+ + \dots + 2X_k^+$$

(Ruelle's inequality)

Proof: take  $s=1$  in the first inequality  
 $d_0 = 1$ .

Cor 2:

Same hypothesis

$$h_\mu(f) \leq \log d_k - 2X_1^- - \dots - 2X_k^-$$

topological  
degree

"inverse Ruelle's inequality"

Proof:  $s=k$  in the second inequality.

Proof of the previous theorem

(10)

$$d_0 = 1 \leq d_1 \leq \dots \leq d_{s-1} < d_s > d_{s+1} \geq \dots \geq d_k$$

measure  $\mu$  invariant ergodic

with  $\log d(x, \mathcal{R}) \in L^1(\mu)$

$$h_\mu(f) > \max(\log d_{s-1}, \log d_{s+1})$$

$\Rightarrow \mu$  is hyperbolic

$$\lambda_1 \geq \dots \geq \lambda_s > 0$$

$$0 > \lambda_{s+1} \geq \dots \geq \lambda_k$$

Proof:

Suppose  $\lambda_s \leq 0$

1<sup>st</sup> formula:

$$\lambda_{s-1}^+ = \dots = \lambda_s^+ = 0 = \dots = \lambda_k^+$$

$$\text{because } 0 \geq \lambda_{s-1} = \dots = \lambda_s \geq \dots \geq \lambda_k$$

$$\log d_{s-1} < h_\mu(f) \leq \log d_{s-1} \leq \log d_{s-1}$$

$\rightarrow$  contradiction.

$$\Rightarrow \lambda_s > 0$$

By using the second formula with  $s = s+1$

$$\rightarrow \lambda_{s+1} < 0.$$

(11)

IV) Ideas for the proof of the inequalities:

an easier case:

$$X = \mathbb{C}P^2$$

$$X_1 \geq X_2 \quad \text{we suppose } X_1 > X_2 \quad S=2$$

the first inequality becomes ( $l=0$ )


$$h_\mu(f) \leq \log d_1 + 2X_2^+$$

$$h_\mu(f) = \lim_{S \rightarrow 0} \underline{\lim} -\frac{1}{n} \log \mu B_n(x, S)$$

$$\mu B_n(x, S) \approx e^{-h_\mu n}$$

 $x$  generic  
for  $\mu$  $\rightarrow$  we can find  $x_1, \dots, x_N \in X$ with  $N \approx e^{h_\mu n}$  and  $d_n(x_i, x_j) \geq S$  if  $i \neq j$ and  $x_i =$  good points for Pesin's theory. $B_n(x_1, S)$  $B_n(x_2, S)$ 

...


 $B_n(x_N, S)$ 

(12)

$x_1 > x_2$  so through each  $x_i$  we can construct an "approximate stable manifold"

$W^s(x_i)$   $\uparrow$   $x_i$   $W^s(x_i)$  dimension 1  
 diameter,  $\bigcup_{i=0, \dots, n-1} W^s(x_i) \leq \frac{\delta}{4}$   
 area  $\approx e^{-2x_2 + n}$

to realize that, we use the graph trans.-form:



if  $x_2 \leq 0 \rightarrow$  we do a cut-off  
 we keep the part in a box of size  $\frac{\delta}{4}$

if  $x_2 > 0 \rightarrow$  we keep all

size  $\approx \frac{\delta}{4} e^{-x_2}$   
 diameter  $\frac{\delta}{4}$

we start again this process

(13)

$\pi \circ \omega^s(x)$


size  
diameter

$\approx \delta/4$  if  $x_2 \leq 0$

$\frac{\delta}{4} e^{-x_2^+ n}$  if  $x_2 > 0$

area  $\approx e^{-2x_2^+ n} c(\delta)$   $x_2^+ = \max(x_2, 0)$

So we have  $\approx e^{h\mu|f|n}$  approximate stable manifold which have area  $\approx e^{-2x_2^+ n} c(\delta)$



$a_n =$  The area of  $\pi \left( \bigcup_{i=1}^n \omega^s(x_i) \right) \geq e^{h\mu|f|n - 2x_2^+ n}$


if  $a_n \leq d_1^n$

$\Rightarrow \log d_1 \geq h\mu|f| - 2x_2^+ \rightarrow \underline{OK.}$


We can find a line  $L$  with

$$\#(L \cap \mathcal{W}^S) \geq e^{h_{\mu} f(n) - 2\chi_2^+ n}$$

(14)



Fundamental remark: Buzzi / Nowhouse




$$d_n(y_i, y_j) \geq \delta/2$$

the diameter of  $f^p(W^S(x_i)) \leq \delta/4$   
 $p=0, \dots, n-1$

and  $d_n(x_i, x_j) \geq \delta \rightarrow \exists p \in [0, n-1]$   
 $d(f^p(x_i), f^p(x_j)) \geq \delta$





$\Rightarrow d(f^e(y_i), f^e(y_j)) \geq \delta/2.$

we proved

$e^{h_{\text{top}}(L) - 2\epsilon + n}$

$\leq$  maximal cardinality of a  $(n, \frac{\delta}{2})$  separated set in  $L$ .

$\leq d^n$

the idea is the same than for (Dinh - Sibony)

$h_{\text{top}}(L) \leq \max_{a \leq i \leq k} \log d_i$

$X \rightsquigarrow L$