

Morita equivalences between blocks of finite group algebras

Department of Mathematics and Informatics,
Graduate School of Science, Chiba University
e-mail `koshitan@math.s.chiba-u.ac.jp`
千葉大学 大学院理学研究科

Shigeo Koshitani 越谷 重夫

1. Introduction and notation

In representation theory of finite groups, particularly, in modular representation theory, studying structure of p -blocks (block algebras) of finite groups G , where p is a prime number, is one of the most important and interesting things.

Notation 1.1. Throughout this note we use the following notation and terminology. We denote by G always a finite group, and let p be a prime. Then, a triple $(\mathcal{K}, \mathcal{O}, k)$ is so-called a p -modular system, which is big enough for all finitely many finite groups which we are looking at, including G . Namely, \mathcal{O} is a complete discrete valuation ring, \mathcal{K} is the quotient field of \mathcal{O} , \mathcal{K} and \mathcal{O} have characteristic zero, and k is the residue field $\mathcal{O}/\text{rad}(\mathcal{O})$ of \mathcal{O} such that k has characteristic p . We mean by "big enough" above that \mathcal{K} and k are both splitting fields for the finite groups mentioned above. Let A be a block of $\mathcal{O}G$ (and sometimes of kG) with a defect group P . We denote by $\text{mod-}kG$ and by $\text{mod-}A$ the categories of finitely generated right kG - and A -modules, respectively. We write $B_0(kG)$ for the principal block algebra of kG . For the notation and terminology we shall not explain precisely, see the books of [2] and [3].

Setup 1.2. Throughout this note all the time except in Theorem 2.1 our situation is the following: Namely, G and H are finite groups which have the same Sylow p -subgroup P , and hence $P \subseteq G \cap H$. Assume that \tilde{G} is a normal subgroup of G and \tilde{H} is a normal subgroup of H such that \tilde{G} and \tilde{H} have the same Sylow p -subgroup \tilde{P} , and hence $\tilde{P} \subseteq \tilde{G} \cap \tilde{H}$, and moreover that $G/\tilde{G} \cong H/\tilde{H}$.

Remark 1.3. If the factor group G/\tilde{G} is p' -groups, then we know essentially by the famous result due to H.Maschke (1898) that the ring extension $k\tilde{G} \subseteq kG$ is a so-called *separable* extension. Then, roughly speaking, $\text{mod-}kG$ and $\text{mod-}k\tilde{G}$ are in some sense *similar* (of course, even the numbers of simples in the two module categories are different, though). Therefore, much more interesting situation should be the case where $|G/\tilde{G}|$ is divisible by p . Then, here comes our situation.

Our situation 1.4. We still keep the setup 1.2. In addition we assume that the factor groups $G/\tilde{G} \cong H/\tilde{H}$ are p -groups. Surely, the factor groups are isomorphic to P/\tilde{P} , too. Then, we naturally come to the following questions.

Questions 1.5. Our main concern in this note is the following:

- (i) If there is a *nice* equivalence between $\text{mod-}k\tilde{G}$ and $\text{mod-}k\tilde{H}$, can we lift it to a *nice* equivalence between $\text{mod-}kG$ and $\text{mod-}kH$?
- (ii) If there is a *nice* equivalence between $\text{mod-}kG$ and $\text{mod-}kH$, can we descend it to a *nice* equivalence between $\text{mod-}k\tilde{G}$ and $\text{mod-}k\tilde{H}$?

2. Results

In this short section we shall list two results which come up from Question 1.5.

Theorem 2.1. *Assume 1.4, however, note that we do not assume that P and \tilde{P} are Sylow p -subgroups of G and \tilde{G} , respectively. Namely, P is just a p -subgroup of G and also of H , and \tilde{P} is just a p -subgroup of \tilde{G} and also of \tilde{H} . We assume then that P is a defect group of A and B , and \tilde{P} is a defect group of \tilde{A} and \tilde{B} . Moreover, we suppose*

that the factor groups $Q := G/\tilde{G} \cong H/\tilde{H} \cong P/\tilde{P}$ are just cyclic group C_p of order p , and that $A, \tilde{A}, B, \tilde{B}$ respectively are block algebras of $kG, k\tilde{G}, kH, k\tilde{H}$, such that A covers \tilde{A} and B covers \tilde{B} . Set $\Delta Q := \{(u, u) \in Q \times Q | u \in Q\}$. We assume, in addition, that \tilde{A} and \tilde{B} are both ΔQ -invariant, that is, they are stable under conjugation action by all elements in Q . Set furthermore that $\Delta := (\tilde{G} \times \tilde{H})\Delta Q = (\tilde{G} \times \tilde{H})\Delta P = (\tilde{G} \times \tilde{H})\Delta G = (\tilde{G} \times \tilde{H})\Delta H$. Then, we get the following: Suppose that there is a bounded complex $\tilde{M}^\bullet \in C^b(\mathcal{O}\tilde{A}\text{-mod-}\mathcal{O}\tilde{B})$ of finitely generated $(\mathcal{O}\tilde{A}, \mathcal{O}\tilde{B})$ -bimodules such that

- (1) $\tilde{M}^\bullet \otimes_{\mathcal{O}} \mathcal{K}$ induces an isometry \tilde{I} from $\mathbb{Z}\text{Irr}(\tilde{A})$ to $\mathbb{Z}\text{Irr}(\tilde{B})$,
- (2) \tilde{M}^\bullet is perfect (exact), that is, all terms in the complex \tilde{M}^\bullet are projective as left $\mathcal{O}\tilde{G}$ -modules and also as right $\mathcal{O}\tilde{H}$ -modules (and hence the isometry \tilde{I} above is perfect),
- (3) the complex \tilde{M}^\bullet extends from $\tilde{G} \times \tilde{H}$ to Δ .

Then, we can define a bounded complex $M^\bullet := \tilde{M}^\bullet_{\tilde{G} \times \tilde{H} \rightarrow \Delta} \uparrow^{G \times H} \in C^b(\mathcal{O}A\text{-mod-}\mathcal{O}B)$, and the new complex M^\bullet induces a perfect isometry from $\mathbb{Z}\text{Irr}(A)$ to $\mathbb{Z}\text{Irr}(B)$. where $M^\bullet := \tilde{M}^\bullet_{\tilde{G} \times \tilde{H} \rightarrow \Delta} \uparrow^{G \times H}$ is an induced complex by applying the functor $-\otimes_{\mathcal{O}\Delta} \mathcal{O}[G \times H]$ to the bounded complex \tilde{M}^\bullet .

Corollary 2.2. *We easily get [1, Example 4.3] in our previous paper by making use of Theorem 2.1.*

Theorem 2.2. *Assume 1.4. Here we assume that P is a Sylow p -subgroup of G and H , and also \tilde{P} is a Sylow p -subgroup of \tilde{G} and \tilde{H} . Moreover, we suppose that the factor groups $Q := G/\tilde{G} \cong H/\tilde{H} \cong P/\tilde{P}$ are isomorphic finite p -groups, and that $A, \tilde{A}, B, \tilde{B}$ respectively are principal block algebras of $kG, k\tilde{G}, kH, k\tilde{H}$. Set $\Delta P := \{(u, u) \in P \times P | u \in P\}$. Moreover, we denote by $\text{Scott}(G \times H, \Delta P)$ the (Alperin-)Scott module in $G \times H$ with respect to a subgroup ΔP of $G \times H$, see [2, Chap.4 Theorem 8.4, Corollary 8.5]. Then, we get the following: If ${}_A M_B := \text{Scott}(G \times H, \Delta P)$ induces a Morita equivalence (and hence it is a Puig equivalence) between A and B , then ${}_{\tilde{A}} \tilde{M}_{\tilde{B}} := \text{Scott}(\tilde{G} \times$*

$\tilde{H}, \Delta\tilde{P}$) induces a Morita equivalence (and hence it is a Puig equivalence) between \tilde{A} and \tilde{B} . (Recall that $A := B_0(kG) = \text{Scott}(G \times G, \Delta\tilde{P})$ and $B := B_0(kH) = \text{Scott}(H \times H, \Delta\tilde{P})$).

Acknowledgment. The author is grateful to Professor Akihide Hanaki for organizing such a wonderful meeting held in Shinshu University as a RIMS meeting during 17–20 November, 2009.

REFERENCES

- [1] M. Holloway, S. Koshitani, N. Kunugi, Blocks with nonabelian defect groups which have cyclic subgroups of index p , *Archiv der Mathematik* **94** (2010) 101–116.
- [2] H. Nagao and Y. Tsushima, *Representations of Finite Groups*, Academic Press, New York, 1988.
- [3] J. Thévenaz, *G-Algebras and Modular Representation Theory*. Clarendon Press, Oxford, 1995.