

The Parents and the Children of Non-Weierstrass semigroups ¹

神奈川工科大学・基礎・教養教育センター 米田 二良 (Jiryo Komeda)
Center for Basic Education and Integrated Learning
Kanagawa Institute of Technology

Abstract

For a numerical semigroup H of genus g Bras-Amorós [1] defined a numerical semigroup $p(H)$ of genus $g-1$ which is called the *parent* of H . She also called H a *child* of $p(H)$. We consider three kinds of non-Weierstrass semigroups. Children of some non-Weierstrass semigroups in each case are investigated.

1 Parents and children of numerical semigroups

Let \mathbb{N}_0 be the additive monoid of non-negative integers. A submonoid H of \mathbb{N}_0 is called a *numerical semigroup* if the complement $\mathbb{N}_0 \setminus H$ is finite. The cardinality of $\mathbb{N}_0 \setminus H$ is called the *genus* of H , denoted by $g(H)$. Let $f(H)$ be the greatest number of $\mathbb{N}_0 \setminus H$, which is called the *Frobenius number* of H .

Example 1.1 Let $H = \langle 3, 5, 7 \rangle$. Then we have $\mathbb{N}_0 \setminus H = \{1, 2, 4\}$, which implies that $g(\langle 3, 5, 7 \rangle) = 3$. Moreover, we get $f(\langle 3, 5, 7 \rangle) = 4$.

Let $p(H) = H \cup \{f(H)\}$, which is called the *parent* of H . Then $p(H)$ is a numerical semigroup of genus $g(H) - 1$. Moreover, H is called a *child* of $p(H)$. These definitions are due to Bras-Amorós [1].

Example 1.2 We have $p(\langle 3, 5, 7 \rangle) = \langle 3, 4, 5 \rangle$. Hence, the semigroup $\langle 3, 4, 5 \rangle$ is the parent of $\langle 3, 5, 7 \rangle$. Conversely, the semigroup $\langle 3, 5, 7 \rangle$ is a child of $\langle 3, 4, 5 \rangle$.

Let $M(H)$ be a unique minimal set of generators for H . An element $\mu \in M(H)$ is called an *effective minimal generator* if $\mu > f(H)$. When

¹This paper is an extended abstract and the details will appear elsewhere.

$\{\mu_1 < \dots < \mu_{r+e}\} = M(H)$ where μ_i 's ($i \geq r+1$) are effective, we denote H by $\langle \mu_1, \dots, \mu_r \mid \mu_{r+1}, \dots, \mu_{r+e} \rangle$.

Remark 1.1 For an effective minimal generator μ we set $H_\mu = H \setminus \{\mu\}$. Then H_μ becomes a numerical semigroup of genus $g(H)+1$. Moreover, the set $p^{-1}(H)$ of children of H is $\{H_\mu \mid \mu \text{ is an effective minimal generator of } H\}$.

Example 1.3 i) Let $H = \langle 2, 3 \rangle$. Then $H_3 = \langle 2 \mid 5 \rangle$ and $H_2 = \langle 3, 4, 5 \rangle$ are the children of H .

ii) Let $H = \langle 3, 4, 5 \rangle$. Then the children of H are $H_5 = \langle 3, 4 \mid \rangle$, $H_4 = \langle 3 \mid 5, 7 \rangle$ and $H_3 = \langle 4, 5, 6, 7 \rangle$.

Example 1.4 $\langle 3, 4 \mid \rangle$ and $\langle 3, 5 \mid \rangle$ have no child.

2 Non-Weierstrass semigroups

In this paper a *curve* means a projective non-singular curve over an algebraically closed field k of characteristic 0. Let $k(C)$ be the field of rational functions on C . A numerical semigroup H is said to be *Weierstrass* if there is a pointed curve (C, P) such that

$$H = H(P) = \{n \in \mathbb{N}_0 \mid \exists f \in k(C) \text{ with } (f)_\infty = nP\}.$$

Remark 2.1 (Buchweitz [3]) For $m \geq 2$, we define

$$L_m(H) := \{a_1 + \dots + a_m \mid a_i \in \mathbb{N}_0 \setminus H\}.$$

If there exists m such that $\#L_m(H) \geq (2m-1)(g(H)-1)+1$, then H is non-Weierstrass.

Example 2.1 (Buchweitz [3]) Let $B_{16} = \langle 13 \rightarrow 18, 20, 22, 23 \rangle$ where for two integers $a < b$ the symbol $a \rightarrow b$ means the consecutive numbers $a, a+1, \dots, b-1, b$. Then $\mathbb{N}_0 \setminus B_{16} = \{1 \rightarrow 12, 19, 21, 24, 25\}$, which implies that $g(B_{16}) = 16$ and $\#L_2(B_{16}) = 46 = (2*2-1)(16-1)+1$. Hence, B_{16} is non-Weierstrass. We note that $B_{16} = \langle 13 \rightarrow 18, 20, 22, 23 \mid \rangle$ has no child.

A numerical semigroup H is said to be *Buchweitz* if there exists $m \geq 2$ such that $\#L_m(H) \geq (2m-1)(g(H)-1)+1$.

Problem 2.1 Are there non-Weierstrass semigroups which are not Buchweitz?

The above problem was solved by Stöhr and Torres [5]. We will describe their method. For a numerical semigroup H we set

$$d_2(H) = \left\{ \frac{h}{2} \mid h \in H \text{ is even} \right\},$$

which is also a numerical semigroup.

Remark 2.2 (Stöhr-Torres) *Let H be a numerical semigroup such that $d_2(H)$ is non-Weierstrass. If $g(H) \geq 6g(d_2(H)) + 4$, then H is non-Weierstrass.*

Example 2.2 (Stöhr-Torres) Let B_{16} be as in Example 2.1. Assume that $g \geq 100 = 6 * 16 + 4$. We set

$$ST_g = 2B_{16} \cup \{2g - 1 - 2t \mid t \in \mathbb{Z} \setminus B_{16}\}.$$

Then we have $d_2(ST_g) = B_{16}$ and $g(ST_g) = g$. By Remark 2.2 we see that ST_g is a non-Weierstrass semigroup, which is non-Buchweitz. Moreover, we obtain

$$ST_{100} = 2\langle 13 \rightarrow 18, 20, 22, 23 \rangle + \langle 149, 151, 157, 161 \mid \rangle$$

with $f(ST_{100}) = 199$. Hence, ST_{100} has no child.

A non-Buchweitz numerical semigroup H is said to be *Stöhr-Torres* if there exists a finite sequence $\{H_i\}_{i=0,1,\dots,n}$ with Buchweitz H_0 , $d_2(H_i) = H_{i-1}$ and $H_n = H$ satisfying $g(H_i) \geq 6g(H_{i-1}) + 4$ for $i = 1, \dots, n$. Here, we pose the following problem:

Problem 2.2 Are there non-Weierstrass semigroups which are neither Buchweitz nor Stöhr-Torres?

The above problem was also solved in [4]. In fact, we have the following:

Example 2.3 ([4]) $H = \langle 8, 12, 18, 22, 51, 55 \rangle$ is a non-Weierstrass semigroup which is neither Buchweitz nor Stöhr-Torres. In fact, the numerical semigroup $d_2(H) = \langle 4, 6, 9, 11 \rangle$ of genus 5 is Weierstrass. Moreover, we get $g(H) = 34$ and $f(H) = 69$. Hence, we obtain $H = \langle 8, 12, 18, 22, 51, 55 \mid \rangle$, which implies that H has no child.

From the results in this section we have the following problem which will be considered in the next section:

Problem 2.3 Are there non-Weierstrass semigroups which have children?

3 Infinite chains

We can find a Buchweitz semigroup which has a child.

Example 3.1 We set

$$H = \langle 112 \longrightarrow 119, 121 \longrightarrow 143, 145 \longrightarrow 215, 217, 218, 219 | 223 \rangle.$$

The semigroup H is Buchweitz. It has a child $H' = H \setminus \{223\}$ which is also Buchweitz. In this case, $f(H) = 222$ and $g(H) = 117$.

We also see that there is a Stöhr-Torres semigroup with a child.

Example 3.2 We set

$$H = 2\langle 13 \longrightarrow 18, 20, 22, 23 \rangle + \langle 151, 153, 169, 163 | 201 \rangle.$$

The semigroup H with $f(H) = 175$ is Stöhr-Torres. It has a child $H' = H \setminus \{201\} = ST_{101}$ which is also Stöhr-Torres. In this case, we have $g(H) = 100$ and $d_2(H) = \langle 13 \longrightarrow 18, 20, 22, 23 \rangle = B_{16}$.

We are interested in numerical semigroups which leave offsprings through all eternity. So, we define the following: A numerical semigroup H has an *infinite chain* if there exists an infinite sequence $\{H^{(i)}\}_{i \geq 0}$ such that $H^{(0)} = H$ and $p(H^{(i)}) = H^{(i-1)}$ for all $i \geq 1$. This definition is due to Bras-Amorós and Bulygin [2].

Example 3.3 i) Let $g \geq 1$. We set $H = \langle 2, 2g+1 \rangle$. Then H has an infinite chain. In fact, we get

$$p(\langle 2, 2(g+i)+1 \rangle) = \langle 2, 2(g+i-1)+1 \rangle$$

for all $i \geq 1$. Namely we have an infinite sequence $\{H^{(i)} = \langle 2, 2(g+i)+1 \rangle\}_{i \geq 0}$ with $H^{(0)} = H$.

ii) Let $g \geq 1$. We set $H = \langle g+1 \longrightarrow 2g+1 \rangle$, which is ordinary. Then H has an infinite chain, because we obtain

$$p(\langle g+i+1 \longrightarrow 2(g+i)+1 \rangle) = \langle g+(i-1)+1 \longrightarrow 2(g+i-1)+1 \rangle$$

for all $i \geq 1$. That is to say, we have an infinite sequence

$$\{H^{(i)} = \langle g + i + 1 \longrightarrow 2(g + i) + 1 \rangle\}_{i \geq 0}$$

with $H^{(0)} = H$.

By the above examples the most general or the most special numerical semigroup has an infinite chain. Moreover, we want to consider the following problem:

Problem 3.1 Are there non-Weierstrass semigroups which have infinite chains?

We can find Stöhr-Torres semigroups which have infinite chains.

Theorem 3.1 Let $n \geq 10g(B_{16}) + 9 = 169$ be an odd number. We set

$$H = 2B_{16} + \langle n, n + 2, \dots, n + 2 \cdot 15 \rangle.$$

Then H is a non-Weierstrass semigroup which has an infinite chain. It is Stöhr-Torres.

Proof. We have

$$\begin{aligned} & p(2B_{16} + \langle n + 2i, n + 2i + 2, \dots, n + 2i + 2 \cdot 15 \rangle) \\ &= 2B_{16} + \langle n + 2(i - 1), n + 2(i - 1) + 2, \dots, n + 2(i - 1) + 2 \cdot 15 \rangle. \end{aligned}$$

Namely we have an infinite sequence

$$\{H^{(i)} = 2B_{16} + \langle n + 2i, n + 2i + 2, \dots, n + 2i + 2 \cdot 15 \rangle\}_{i \geq 0}$$

with Stöhr-Torres $H^{(i)}$'s. □

Finally, the unsolved problems are presented using the following table:

- Problem 3.2**
- i) Is there a non-Weierstrass semigroup with a child which is neither Buchweitz nor Stöhr-Torres?
 - ii) Is there a Buchweitz semigroup with an infinite chain?
 - iii) Is there a non-Weierstrass semigroup with an infinite chain which is neither Buchweitz nor Stöhr-Torres?

Summarizing the results in this paper and the above open problems we get the table.

Properties	Weierstrass	New Type	Stöhr-Torres	Buchweitz
Children	\exists	?	\exists	\exists
Infinite chains	\exists	?	\exists	?

Here, "New Type" means a non-Weierstrass semigroup which is neither Buchweitz nor Stöhr-Torres. We note that Example 2.3 gives a "New Type" semigroup.

References

- [1] M. Bras-Amorós, *Fibonacci-like behavior of the number of numerical semigroups of a given genus*, Semigroup Forum **76** (2008) 379-384.
- [2] M. Bras-Amorós and S. Bulygin, *Towards a better understanding of the semigroup tree*, Semigroup Forum **79** (2009) 561-574.
- [3] R.O. Buchweitz, *On Zariski's criterion for equisingularity and non-smoothable monomial curves*, preprint 113, University of Hannover, 1980.
- [4] J. Kameda, *Double coverings of curves and non-Weierstrass semigroups*, Submitted.
- [5] F. Torres, *Weierstrass points and double coverings of curves with application: Symmetric numerical semigroups which cannot be realized as Weierstrass semigroups*, Manuscripta Math. **83** (1994) 39-58.