

# Geometric means on gyrogroups

Takuro Honma

Graduate School of Science and Technology, Niigata University

## 1 Introduction

Abraham Albert Ungar initiated the theory of gyrogroups in 1989 [1] associated with the study of Einstein’s velocity addition in the theory of special relativity. It is the study of analytic hyperbolic geometry. A gyrogroup has a weak associativity. It is a generalization of a group. The set of all positive invertible elements in a unital  $C^*$ -algebra is an example of a gyrogroup. It is difficult to give an appropriate definition for "the geometric mean" of more than two points on a gyrovector space because of nonassociativity and noncommutativity of the operation. So we define a geometric mean for the Einstein gyrovector space.

## 2 Einstein gyrovector space

Einsteinian velocities with the Einstein’s velocity addition based on the special theory of relativity is a gyrocommutative gyrogroup. Let  $\mathbb{V}$  be a real Hilbert space. Let  $\mathbb{V}_1$  be an open unit ball of  $\mathbb{V}$ , that is,

$$\mathbb{V}_1 = \{\mathbf{v} \in \mathbb{V} : \|\mathbf{v}\| < 1\}.$$

The Einstein addition  $\oplus_E$  on  $\mathbb{V}_1$  is a binary operation on  $\mathbb{V}_1$  given by the equation

$$\mathbf{a} \oplus_E \mathbf{b} = \frac{1}{1 + \mathbf{a} \cdot \mathbf{b}} \left\{ \mathbf{a} + \frac{1}{\gamma_{\mathbf{a}}} \mathbf{b} + \frac{\gamma_{\mathbf{a}}}{1 + \gamma_{\mathbf{a}}} (\mathbf{a} \cdot \mathbf{b}) \mathbf{a} \right\},$$

where  $\gamma_{\mathbf{u}}$  is the Lorentz gamma factor defined by

$$\gamma_{\mathbf{u}} = \frac{1}{\sqrt{1 - \|\mathbf{u}\|^2}}.$$

These  $\cdot$  and  $\|\cdot\|$  denote the usual inner product and the norm of  $\mathbb{V}$  respectively. Einstein scalar multiplication  $\otimes_E$  is given by the form

$$\begin{aligned} r \otimes_E \mathbf{a} &= \frac{(1 + \|\mathbf{a}\|)^r - (1 - \|\mathbf{a}\|)^r}{(1 + \|\mathbf{a}\|)^r + (1 - \|\mathbf{a}\|)^r} \frac{\mathbf{a}}{\|\mathbf{a}\|} \\ &= \tanh(r \tanh^{-1} \|\mathbf{a}\|) \frac{\mathbf{a}}{\|\mathbf{a}\|}, \end{aligned}$$

where  $r \in \mathbb{R}$ ,  $\mathbf{a} \in \mathbb{V}_1 \setminus \{\mathbf{0}\}$  and  $r \otimes_E \mathbf{0} = \mathbf{0}$ . By Theorem 6.84 in [2],  $(\mathbb{V}_1, \oplus_E, \otimes_E)$  is defined by a gyrovector space.

We define the set  $\|\mathbb{V}_1\| = \{\pm\|\mathbf{v}\| : \mathbf{v} \in \mathbb{V}_1\} \subset \mathbb{R}$ , which coincides with the open interval  $(-1, 1)$ .  $\|\mathbb{V}_1\|$  admits addition  $\oplus'_E$  and scalar multiplication  $\otimes'_E$  given by the following:

$$\begin{aligned} a \oplus'_E b &= \frac{a+b}{1+ab} \\ r \otimes'_E a &= \tanh(r \tanh^{-1} a) \end{aligned}$$

where  $a, b \in \|\mathbb{V}_1\|$  and  $r \in \mathbb{R}$ . These two operator is induced by  $\oplus_E$  and  $\otimes_E$ .  $(\|\mathbb{V}_1\|, \oplus'_E, \otimes'_E)$  is a real one dimensional space.

### 3 The metric space on $(\mathbb{V}_1, \oplus_E, \otimes_E)$

The gyrometric is defined by

$$d(\mathbf{a}, \mathbf{b}) = \|\mathbf{a} \ominus_E \mathbf{b}\| \in \|\mathbb{V}_1\|,$$

where  $\mathbf{a}, \mathbf{b} \in \mathbb{V}_1$  and  $\mathbf{a} \ominus_E \mathbf{b} = \mathbf{a} \oplus_E (-\mathbf{b})$ . The gyrometric is not a metric. It satisfies the following properties:

- (1)  $d(\mathbf{a}, \mathbf{b}) \geq 0$  for every  $\mathbf{a}, \mathbf{b} \in \mathbb{V}_1$ ,  $d(\mathbf{a}, \mathbf{b}) = 0 \Leftrightarrow \mathbf{a} = \mathbf{b}$ .
- (2)  $d(\mathbf{a}, \mathbf{b}) = d(\mathbf{b}, \mathbf{a})$  for all  $\mathbf{a}, \mathbf{b} \in \mathbb{V}_1$ .
- (3) The gyrotriangle inequality:

$$d(\mathbf{a}, \mathbf{b}) \leq d(\mathbf{a}, \mathbf{c}) \oplus'_E d(\mathbf{c}, \mathbf{b})$$

for all  $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{V}_1$ .

We define a metric on  $\mathbb{V}_1$  induced the gyrometric;  $f : \|\mathbb{V}_1\| \rightarrow \mathbb{R}$  is  $f(x) = \tanh^{-1}(x)$ . Then  $f$  is monotonic and satisfies the following properties:

- (F1)  $f(a \oplus'_E b) = f(a) + f(b)$  for all  $a, b \in \|\mathbb{V}_1\|$
- (F2)  $f(r \otimes'_E a) = rf(a)$  for all  $a \in \|\mathbb{V}_1\|$  and  $r \in \mathbb{R}$ .

We define  $\delta(\mathbf{a}, \mathbf{b}) = f(d(\mathbf{a}, \mathbf{b}))$ , where  $\mathbf{a}, \mathbf{b} \in \mathbb{V}_1$ .

**Proposition 1.** *The map  $\delta$  give a metric on  $\mathbb{V}_1$ ;  $(\mathbb{V}_1, \delta)$  is a complete metric space.*

### 4 Gyromidpoints and gyrocentroids

Ungar defined the gyromidpoint  $\mathbf{P}_{\mathbf{ab}}^m$  of two elements  $\mathbf{a}, \mathbf{b} \in \mathbb{V}_1$  given by

$$\mathbf{P}_{\mathbf{ab}}^m = \frac{\gamma_{\mathbf{a}} \mathbf{a} + \gamma_{\mathbf{b}} \mathbf{b}}{\gamma_{\mathbf{a}} + \gamma_{\mathbf{b}}},$$

By a natural extension, Ungar [2] define the gyrocentroid  $\mathbb{C}_{abc}^m$  of three elements  $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{V}_1$  written by

$$\mathbb{C}_{abc}^m = \frac{\gamma_a \mathbf{a} + \gamma_b \mathbf{b} + \gamma_c \mathbf{c}}{\gamma_a + \gamma_b + \gamma_c}.$$

The gyromidpoint  $\mathbb{P}_{ab}^m$  satisfies some desirable properties one would expect for means, for example the permutation invariance and the left gyrotranslation invariance which is given by

$$\mathbf{x} \oplus_E \mathbb{P}_{ab}^m = \mathbb{P}_{(x \oplus_E a)(x \oplus_E b)}^m.$$

But the gyrocentroid does not satisfy the left gyrotranslation invariance. In the case of the three points  $\mathbf{0}, \mathbf{0}, \mathbf{c}$ ,  $\mathbb{C}_{abc}^m \neq \frac{1}{3} \otimes_E \mathbf{c}$  by the simple calculation. In this paper we will give a definition of a geometric mean alternatively to remove these difficulties.

## 5 Definition of the geometric mean

We define the geometric mean  $G(\mathbf{a}, \mathbf{b}, \mathbf{c})$  of three elements  $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{V}_1$  as in the following. Starting from  $\mathbf{a}_0 = \mathbf{a}, \mathbf{b}_0 = \mathbf{b}, \mathbf{c}_0 = \mathbf{c}$ , we define  $\mathbf{a}_m, \mathbf{b}_m, \mathbf{c}_m$  by induction on  $m$ . Suppose that  $\mathbf{a}_{m-1}, \mathbf{b}_{m-1}, \mathbf{c}_{m-1}$  are defined. Then

$$\mathbf{a}_m = \mathbf{a}_{m-1} \# \mathbf{b}_{m-1}, \mathbf{b}_m = \mathbf{b}_{m-1} \# \mathbf{c}_{m-1}, \mathbf{c}_m = \mathbf{c}_{m-1} \# \mathbf{a}_{m-1}$$

where  $\mathbf{x} \# \mathbf{y}$  is the gyromidpoint of  $\mathbf{x}$  and  $\mathbf{y}$ . Then  $\lim_{m \rightarrow \infty} \mathbf{a}_m, \lim_{m \rightarrow \infty} \mathbf{b}_m, \lim_{m \rightarrow \infty} \mathbf{c}_m$  exist and they coincide with each other. Define the common limit as  $M_\infty$ . We define  $G(\mathbf{a}, \mathbf{b}, \mathbf{c}) = M_\infty$ .  $G(\mathbf{a}, \mathbf{b}, \mathbf{c})$  is permutation invariant. By a simple calculation,  $G(\mathbf{0}, \mathbf{0}, \mathbf{c}) = \frac{1}{3} \otimes_E \mathbf{c}$  holds.

We define the geometric mean for any finite number of elements as follows. Let  $\Delta_n$  be a  $n$ -points set of  $\mathbb{V}_1$ . We define the geometric mean  $G(\Delta_n)$  by induction of the number of elements  $n$  by generalizing the way as above.

**Definition 1.** (1) Let  $\Delta_2 = \{\mathbf{a}_1, \mathbf{a}_2\} \subset \mathbb{V}_1$ . We define  $G(\Delta_2) = \mathbf{a}_1 \# \mathbf{a}_2$ .

(2) Suppose that we have defined the geometric mean  $G(\Delta_n)$  for any  $\Delta_n$ . Let  $\Delta_{n+1} = \{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_{n+1}\} \subset \mathbb{V}_1$ . Put  $\mathbf{a}_i^0 = \mathbf{a}_i$  for  $i = 1, 2, \dots, n+1$ . For a positive integer  $m$ , we define  $\mathbf{a}_i^m$  for  $i = 1, 2, \dots, n+1$  by induction on  $m$  as follows. Suppose that  $\mathbf{a}_i^{m-1}$  for  $i = 1, 2, \dots, n+1$  is defined, put

$$\mathbf{a}_i^m = G(\{\mathbf{a}_1^{m-1}, \mathbf{a}_2^{m-1}, \dots, \mathbf{a}_{i-1}^{m-1}, \mathbf{a}_{i+1}^{m-1}, \dots, \mathbf{a}_{n+1}^{m-1}\})$$

for  $i = 1, 2, \dots, n+1$ . The point  $\mathbf{a}_i^m$  is well defined since we suppose that  $G(\Delta_n)$  is defined for an  $n$ -point set  $\Delta_n$ .

Put  $\Delta_{n+1}^m = \{\mathbf{a}_1^m, \mathbf{a}_2^m, \dots, \mathbf{a}_{n+1}^m\}$ . Then the limit  $\lim_{m \rightarrow \infty} \mathbf{a}_i^m$  exists for each  $i = 1, 2, \dots, n+1$  and they coincide with each other. Define the common limit by  $M_\infty$ . We define  $G(\Delta_{n+1})$  to be  $M_\infty$ .

This definition is a modification of the definition of the geometric mean for a positive definite matrices by Ando, Chi-Kwong Li and Mathias[5]

## 6 A proof of the existence of and the coincidence of $\lim_{m \rightarrow \infty} \mathbf{a}_i^m$

To prove the existence of  $\lim_{m \rightarrow \infty} \mathbf{a}_i^m$  and the coincidenceness of each other, we need some preparations. We define a gyroline and a gyrosegment in the Einstein gyrovector space.

**Definition 2.** Let  $\mathbf{a}, \mathbf{b}$  be elements of  $\mathbb{V}_1$ . The gyroline  $L(\mathbf{a}, \mathbf{b}) = \{\mathbf{a} \oplus_E t \otimes_E (\ominus_E \mathbf{a} \oplus_E \mathbf{b}) : t \in \mathbb{R}\}$ . The gyrosegment  $S(\mathbf{a}, \mathbf{b}) = \{\mathbf{a} \oplus_E t \otimes_E (\ominus_E \mathbf{a} \oplus_E \mathbf{b}) : 0 \leq t \leq 1\}$ .

$\mathbf{a} \#_t \mathbf{b} = \mathbf{a} \oplus_E t \otimes_E (\ominus_E \mathbf{a} \oplus_E \mathbf{b})$  is called a gyro  $t$ -point on a gyroline or gyrosegment. If  $t = \frac{1}{2}$  then it is the gyromidpoint, that is

$$\mathbf{a} \# \mathbf{b} = \frac{1}{2} \otimes_E (\mathbf{a} \boxplus_E \mathbf{b}),$$

where  $\boxplus_E$  is coaddition on  $(\mathbb{V}_1, \oplus_E, \otimes_E)$  defined by the following;

$$\mathbf{a} \boxplus_E \mathbf{b} = \frac{\gamma_{\mathbf{a}} + \gamma_{\mathbf{b}}}{\gamma_{\mathbf{a}}^2 + \gamma_{\mathbf{b}}^2 + \gamma_{\mathbf{a}} \gamma_{\mathbf{b}} (\mathbf{a} \cdot \mathbf{b}) - 1} (\gamma_{\mathbf{a}} \mathbf{a} + \gamma_{\mathbf{b}} \mathbf{b}).$$

**Proposition 2.** Coaddition  $\boxplus_E$  is commutative, thus  $\mathbf{a} \# \mathbf{b} = \mathbf{b} \# \mathbf{a}$ .

By Proposition2, the gyromidpoint is permutation invariant.

We define a gyroconvex set and a gyroconvex hull in the Einstein gyrovector space.

**Definition 3.** A subset  $C$  of  $\mathbb{V}_1$  is gyroconvex set if for any  $\mathbf{a}, \mathbf{b} \in C$ ,  $S(\mathbf{a}, \mathbf{b}) \subset C$ .

**Definition 4.**  $X \subset \mathbb{V}_1$ . A gyroconvex hull of  $X$  is defined by:

$$\text{conv}(X) = \cap \{C \subset \mathbb{V}_1 : X \subset C \text{ and } C \text{ is convex}\}.$$

These definitions are modifications of definitions of the geometric mean for a positive definite matrices by Bhatia and Holbrook[4]. By a simple calculation, we have the following property.

**Proposition 3.** The set  $C_m$  for every  $m \in \mathbb{N} \cap \{0\}$  is defined by induction. Let  $C_0 = X$ . If  $C_{m-1}$  is defined, then put  $C_m = \cup_{\mathbf{a}, \mathbf{b} \in C_{m-1}} S(\mathbf{a}, \mathbf{b})$ . A gyroconvex hull of  $X$  can be written by

$$\text{conv}(X) = \bigcup_{m=0}^{\infty} C_m.$$

We show an inequality which is related gyromidpoints. It plays a crucial role in the proof of the convergence of the sequence  $\{\mathbf{a}_i^m\}$  we have defined before.

**Theorem 1.** For any  $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{V}_1$  we have

$$d(\mathbf{a} \# \mathbf{b}, \mathbf{a} \# \mathbf{c}) \leq \frac{1}{2} \otimes'_E d(\mathbf{b}, \mathbf{c}).$$

This theorem is proved by a simple calculation. By applying gamma identities, we have

$$\begin{aligned} & \left(\frac{1}{2} \otimes_E d(\mathbf{b}, \mathbf{c})\right)^2 - d^2(\mathbf{a}\#\mathbf{b}, \mathbf{a}\#\mathbf{c}) \\ &= \frac{2\left\{2\gamma_{\mathbf{a}\ominus_E\mathbf{b}}\gamma_{\mathbf{b}\ominus_E\mathbf{c}}\gamma_{\mathbf{c}\ominus_E\mathbf{a}} - (\gamma_{\mathbf{a}\ominus_E\mathbf{b}}^2 + \gamma_{\mathbf{b}\ominus_E\mathbf{c}}^2 + \gamma_{\mathbf{c}\ominus_E\mathbf{a}}^2) + 1\right\}}{(1 + \gamma_{\mathbf{b}\ominus_E\mathbf{c}})(1 + \gamma_{\mathbf{a}\ominus_E\mathbf{b}} + \gamma_{\mathbf{b}\ominus_E\mathbf{c}} + \gamma_{\mathbf{c}\ominus_E\mathbf{a}})^2}. \end{aligned} \quad (6.1)$$

Applying the left gyrotranslation of  $\mathbf{c}$ , put  $\mathbf{A} = \ominus_E \mathbf{c} \oplus_E \mathbf{a}$ ,  $\mathbf{B} = \ominus_E \mathbf{c} \oplus_E \mathbf{b}$ , we calculate the numerator of (6.1),

$$\begin{aligned} & 2\gamma_{\mathbf{A}\ominus_E\mathbf{B}}\gamma_{\mathbf{A}}\gamma_{\mathbf{B}} - (\gamma_{\mathbf{A}\ominus_E\mathbf{B}}^2 + \gamma_{\mathbf{A}}^2 + \gamma_{\mathbf{B}}^2) + 1 \\ &= \frac{\|\mathbf{A}\|^2\|\mathbf{B}\|^2 - (\mathbf{A} \cdot \mathbf{B})^2}{(1 - \|\mathbf{A}\|^2)(1 - \|\mathbf{B}\|^2)} \geq 0. \end{aligned}$$

Note: I would like to thank Professor Akinari Hoshi for his calculation about the numerator by computer. By his calculation I convinced that the numerator is greater than or equal to 0. Finally I succeeded to prove it. Theorem 1 is followed by Corollary 1.

**Corollary 1.**  $\delta(\mathbf{a}\#\mathbf{b}, \mathbf{a}\#\mathbf{c}) \leq \frac{1}{2}\delta(\mathbf{b}, \mathbf{c})$  and hence

$$\delta(\mathbf{a}\#\mathbf{b}, \mathbf{c}\#\mathbf{d}) \leq \frac{1}{2}\delta(\mathbf{b}, \mathbf{d}) + \frac{1}{2}\delta(\mathbf{a}, \mathbf{c})$$

Moreover, since  $g(t) = \delta(\mathbf{a}\#_t\mathbf{b}, \mathbf{c}\#_t\mathbf{d})$  is continuous, then  $g$  is convex, i.e.,

$$\delta(\mathbf{a}\#_t\mathbf{b}, \mathbf{c}\#_t\mathbf{d}) \leq (1-t)\delta(\mathbf{a}, \mathbf{c}) + t\delta(\mathbf{b}, \mathbf{d})$$

especially,

$$\delta(\mathbf{a}\#_t\mathbf{b}, \mathbf{a}\#_t\mathbf{c}) \leq t\delta(\mathbf{b}, \mathbf{c})$$

We define  $\text{diam}(X) = \sup\{\delta(\mathbf{x}, \mathbf{y}) : \mathbf{x}, \mathbf{y} \in X\}$ . We have the following properties. By Corollary 1, we have the following

**Proposition 4.** If  $\text{diam}(\{\mathbf{x}_0, \mathbf{y}_0, \mathbf{x}_1, \mathbf{y}_1\}) \leq M$ , then for arbitrary  $\mathbf{x} \in S(\mathbf{x}_0, \mathbf{x}_1)$  and  $\mathbf{y} \in S(\mathbf{y}_0, \mathbf{y}_1)$ ,  $\delta(\mathbf{x}, \mathbf{y}) \leq M$  holds.

Proposition 5 is proved by applying Proposition 3 and Proposition 4.

**Proposition 5.** Let  $X$  be a subset of  $\mathbb{V}_1$

$$\text{diam}(\text{conv}(X)) = \text{diam}(X).$$

Considering the geometric mean of three elements, by Corollary 1 and Proposition 5, we have

$$\text{diam}(\text{conv}(\Delta_3^m)) \leq \frac{1}{2}\text{diam}(\text{conv}(\Delta_3^{m-1})).$$

Since  $(\mathbb{V}_1, \delta)$  is a complete, there exists  $M_\infty \in \mathbb{V}_1$  such that  $\bigcap_{m=0}^{\infty} \text{conv}(\Delta_3^m) = \{M_\infty\}$  by the Cantor's intersection principle. So the limit of  $\mathbf{a}_i^m$  exists for  $i = 1, 2, 3$  and they coincide with each other.

## 7 Properties of the geometric mean

To prove the existence of the geometric mean, we assume the following inequality by induction.

**Theorem 2.** For any sets of  $n$  elements in  $\mathbb{V}_1$ ,  $D_n = \{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n\}$ ,  $D'_n = \{\mathbf{a}'_1, \mathbf{a}'_2, \dots, \mathbf{a}'_n\}$  the following inequality holds:

$$\delta(G(D_n), G(D'_n)) \leq \frac{1}{n} \sum_{i=1}^n \delta(\mathbf{a}_i, \mathbf{a}'_i).$$

We can prove the existence of the geometric mean of more than three elements by a similar way as above. The geometric mean satisfies following properties which is proved by induction.

**Theorem 3.** The geometric mean  $G(\Delta_n)$  satisfies the permutation invariance and the left gyrotranslation invariance;

$$G(\mathbf{x} \oplus_E \Delta_n) = \mathbf{x} \oplus_E G(\Delta_n),$$

where  $\mathbf{x} \oplus_E \Delta_n = \{\mathbf{x} \oplus_E \mathbf{y} : \mathbf{y} \in \Delta_n\}$ .

## References

- [1] A. A. Ungar, *The relativistic noncommutative nonassociative group of velocities and the Thomas rotation*, Result. Math. **16** (1989), 168–179
- [2] A. A. Ungar, *Analytic Hyperbolic Geometry and Albert Einstein's Special Theory of Relativity*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ. 2005. xviii+463 pp. ISBN: 981-256-457-8
- [3] A. A. Unger, *A Gyrovector Space Approach to Hyperbolic Geometry*, Synthesis Lecture on Mathematics and Statistics, 4. Morgan & Claypool Publishers, Williston, VT, 2009. xii+182 pp. ISBN: 978-1-59829-822-2
- [4] R. Bhatia and J. Holbrook, *Riemannian geometry and matrix geometric means*, Linear Algebra Appl. **413** (2006), 594-618
- [5] T. Ando, Chi-Kwong Li and R.Mathias, *Geometric means*, Linear Algebra Appl. **385** (2004), 305-334