

IRRATIONALITY EXPONENTS OF NUMBERS
RELATED WITH CAHEN'S CONSTANT

IEKATA SHIOKAWA

This is a report on recent results of myself jointly with Duverney [7] on irrationality exponents of numbers related with Cahen's constant.

For a real number α , the irrationality exponent $\mu(\alpha)$ is defined by the greatest lower bound of the set of numbers μ for which the inequality

$$(0.1) \quad \left| \alpha - \frac{p}{q} \right| < \frac{1}{q^\mu}$$

has only finitely many rational solutions p/q , or equivalently the least upper bound of the set of numbers μ for which the inequality (0.1) has infinitely many solutions. If α is irrational, then $\mu(\alpha) \geq 2$. If α is a real algebraic irrationality, then $\mu(\alpha) = 2$ by Roth's theorem. If $\mu(\alpha) = \infty$, then α is called a Liouville number.

The main theorem of this paper, [7] stated below gives lower and upper bounds for the irrationality exponents of continued fractions.

We employ the usual notations for continued fractions :

$$b_0 + \frac{a_1}{b_1 + \frac{a_2}{b_2 + \dots + \frac{a_n}{b_n}}} = b_0 + \frac{a_1}{b_1 + \frac{a_2}{b_2 + \dots + \frac{a_n}{b_n}}} = \frac{A_n}{B_n},$$

and

$$b_0 + \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3} + \dots}} = \lim_{n \rightarrow +\infty} \frac{A_n}{B_n},$$

where $\{A_n\}$ and $\{B_n\}$ are defined by

$$(0.2) \quad \begin{cases} A_{-1} = 1, & A_0 = b_0, & B_{-1} = 0, & B_0 = 1, \\ A_n = b_n A_{n-1} + a_n A_{n-2} & (n \geq 1) \\ B_n = b_n B_{n-1} + a_n B_{n-2} & (n \geq 1) \end{cases}$$

Theorem 1. *Let an infinite continued fraction*

$$(0.3) \quad \alpha = \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3} + \dots}}$$

be convergent, where a_n, b_n ($n \geq 1$) are non zero rational integers. Assume that

$$(0.4) \quad \sum_{n=1}^{+\infty} \left| \frac{a_{n+1}}{b_n b_{n+1}} \right| < \infty,$$

and

$$(0.5) \quad \lim_{n \rightarrow +\infty} \left| \frac{a_1 a_2 \dots a_n}{b_1 b_2 \dots b_n} \right| = 0.$$

2010 *Mathematics Subject Classification.* Primary 11J82; Secondary 11J70.

Key words and phrases. Irrationality exponent, Cahen's constant, Continued fractions, Sylvester sequence, Sierpinski sequence.

Then α is irrational and its irrationality exponent $\mu(\alpha)$ satisfies

$$(0.6) \quad 2 + \sigma \leq \mu(\alpha) \leq 2 + \max(\tau_1, \tau_2),$$

where

$$(0.7) \quad \sigma = \limsup_{n \rightarrow +\infty} \frac{\log |b_{n+1}| - \log |a_1 a_2 \cdots a_{n+1}|}{\log |b_1 b_2 \cdots b_n|},$$

$$(0.8) \quad \tau_1 = \limsup_{n \rightarrow +\infty} \frac{\log |a_1 a_2 \cdots a_{n+1}|}{\log |b_1 b_2 \cdots b_n| - \log |a_1 a_2 \cdots a_n|},$$

and

$$\tau_2 = \limsup_{n \rightarrow +\infty} \frac{\log |b_{n+1}| - \log |a_1 a_2 \cdots a_{n+1}| + 2 \log(A_n, B_n)}{\log |b_1 b_2 \cdots b_n| - \log |a_1 a_2 \cdots a_n|}$$

with (A_n, B_n) the greatest common divisor of A_n and B_n .

We apply Theorem 1 to continued fractions representing numbers related to Cahen's constant and deduce their transcendence from the obtained lower bounds of their irrationality exponents.

In 1880 Sylvester [11] proved that any real number $0 < x < 1$ can be expanded uniquely in the series

$$x = \sum_{n=0}^{+\infty} \frac{1}{t_n},$$

where the t_n are integers satisfying the condition $t_0 \geq 2$, $t_{n+1} \geq t_n^2 - t_n + 1$ ($n \geq 0$), and furthermore that x is irrational if and only if the equality holds for all large n . He examined some of the properties of the (Sylvester) sequence $\{S_n\}_{n \geq 0}$ defined by

$$(0.9) \quad S_0 = 2, \quad S_{n+1} = S_n^2 - S_n + 1 \quad (n \geq 0),$$

which satisfies

$$\sum_{n=0}^{+\infty} \frac{1}{S_n} = 1.$$

Cahen [2] and Sierpinski [9] independently obtained similar results for alternating series; namely, any irrational number $0 < x < 1$ can be uniquely written in the form

$$x = \sum_{n=0}^{+\infty} \frac{(-1)^n}{u_n},$$

where the u_n are integers satisfying $u_0 \geq 1$, $u_{n+1} \geq u_n^2 + u_n$ ($n \geq 0$). As an example, Cahen [2] mentioned that (Cahen's constant)

$$\sum_{n=0}^{+\infty} \frac{(-1)^n}{u_n} = \frac{1}{1} - \frac{1}{2} + \frac{1}{6} - \frac{1}{42} + \frac{1}{1806} - \frac{1}{3263442} + \cdots$$

is an irrational number, where $u_0 = 1$, $u_{n+1} = u_n^2 + u_n$ ($n \geq 0$), and hence $u_n = S_n - 1$ ($n \geq 0$). We note that the sequence $\{s_n\}_{n \geq 0}$ defined by

$$(0.10) \quad s_0 = 2, \quad s_{n+1} = s_n^2 + s_n - 1 \quad (n \geq 0)$$

satisfies

$$\sum_{n=0}^{+\infty} \frac{(-1)^n}{s_n} = \frac{1}{3}.$$

In 1991 Davison and Shallit [4] proved the transcendence of Cahen's constant. Becker [1] generalized the result by Mahler's method.

In this paper we generalize the sequences S_n and s_n defined in (0.9) and (0.10) by introducing the sequences $u_n = u_n(\varepsilon)$ satisfying $u_0 \in \mathbb{N}$, $u_0 > \max(1, \varepsilon)$, and the recurrence

$$(0.11) \quad u_{n+1} = u_n^2 - \varepsilon u_n + \varepsilon \quad (n \geq 0),$$

where ε is a non-zero integer given arbitrary. Next, we define the numbers $\gamma_{l,\varepsilon} = \gamma_{l,\varepsilon}(u_0)$ by

$$(0.12) \quad \gamma_{l,\varepsilon} = \sum_{n=0}^{+\infty} (-1)^n \left(\frac{\varepsilon^n}{u_n - \varepsilon} \right)^l \quad (l = 1, 2, 3, \dots).$$

We expand the numbers $\gamma_{l,\varepsilon}$ in continued fractions whose partial numerators a_n and denominators b_n satisfy the assumptions in Theorem 1. Applying Theorem 1, we obtain the following

Theorem 2. *Let $\gamma_{l,\varepsilon}$ be the numbers defined by (0.12). Assume that u_0 and ε are coprime. Then $\mu(\gamma_{1,\varepsilon}) = 3$ and*

$$(0.13) \quad 2 + \frac{2}{3l - 1} \leq \mu(\gamma_{l,\varepsilon}) \leq 2 + \frac{6(l - 1)}{3l + 1} \quad (l = 2, 3, 4, \dots).$$

Corollary 1. *For every positive integer l , $\gamma_{l,\varepsilon}$ is a non-Liouville transcendental number.*

We give some examples of the numbers $\gamma_{l,\varepsilon}$.

Example 1. When $\varepsilon = 1$ and $u_0 = 2$, we have

$$\gamma_{l,1}(2) = \sum_{n=0}^{+\infty} \frac{(-1)^n}{(S_n - 1)^l} \quad (l = 1, 2, 3, \dots).$$

In particular, $\gamma_{1,1}(2)$ is Cahen's constant.

Example 2. When $\varepsilon = -1$ and $u_0 = 2$, we obtain

$$\begin{aligned} \gamma_{l,-1}(2) &= \sum_{n=0}^{+\infty} \frac{(-1)^n}{(s_n + 1)^l} \quad (l = 2, 4, 6, \dots), \\ \gamma_{l,-1}(2) &= \sum_{n=0}^{+\infty} \frac{1}{(s_n + 1)^l} \quad (l = 1, 3, 5, \dots). \end{aligned}$$

Example 3. When $\varepsilon = 2$ and $u_0 = 3$, u_n is the n -th Fermat number:

$$u_n = F_n = 2^{2^n} + 1.$$

Therefore we have

$$\gamma_{l,2}(3) = \sum_{n=0}^{+\infty} (-1)^n \left(\frac{2^n}{F_n - 2} \right)^l = \sum_{n=0}^{+\infty} (-1)^n \left(\frac{2^n}{2^{2^n} - 1} \right)^l \quad (l = 1, 2, 3, \dots).$$

It should be noted that the irrationality exponent of the sum of the reciprocals of Fermat numbers is equal to 2 (see [3]).

Example 4. Denote by L_n the sequence of Lucas numbers. Define

$$v_n = L_{2^{n+1}} = \Phi^{2^{n+1}} + \Phi^{-2^{n+1}},$$

where $\Phi = \frac{1}{2}(1 + \sqrt{5})$ is the Golden number. Then clearly $v_{n+1} = v_n^2 - 2$. If we put $u_n = v_n + 2$, we see that $u_0 = 5$ and

$$u_{n+1} = u_n^2 - 4u_n + 4$$

for every $n \geq 0$. Therefore

$$\gamma_{l,4}(5) = \sum_{n=0}^{+\infty} (-1)^n \left(\frac{4^n}{L_{2^{n+1}} - 2} \right)^l \quad (l = 1, 2, 3, \dots).$$

As for transcendence, much more general results are obtained by Duverney, Kurosawa, and Shiokawa [6]. They discussed transcendence of the values of the series

$$\sum_{n=0}^{\infty} \frac{a^n}{q(p^n(z))}$$

at algebraic points, where $a \in \overline{\mathbb{Q}}$, $p(z), q(z) \in \overline{\mathbb{Q}}[z]$ with $\deg p(z) \geq 2$ and $\deg q(z) \geq 1$. For example, [6, Example 1.5] states that, if $a \neq 0$ and γ with $S_n \neq \gamma$ for all $n \geq 0$ are algebraic numbers, then

$$\sum_{n=0}^{\infty} \frac{a^n}{(S_n - \gamma)^l},$$

where l is any positive integers, is algebraic if and only if $a = l = 1$ and $\gamma = 0$.

REFERENCES

- [1] P.-G. Becker, *Algebraic independence of the values of certain series by Mahler's method*, Mh. Math. 114 (1992), 183-198.
- [2] E. Cahen, *Note sur un développement des quantités numériques, qui présente quelque analogie avec celui en fraction continue*, Nouv. Ann. Math. 10 (1891), 508-514.
- [3] M. Coons, *On the rational approximation of the sum of the reciprocals of the Fermat numbers*, Ramanujan J. 30, No1 (2013), 39-65; addendum ibid. 37, No1 (2015), 109-111.
- [4] J. L. Davison and J. O. Shallit, *Continued fractions for some alternating series*, Mh. Math. 111 (1991), 119-126.
- [5] D. Duverney, *Number Theory : an elementary introduction through diophantine problems*, *Monographs in Number Theory* 4, World Scientific, 2010.
- [6] D. Duverney, T. Kurosawa, and I. Shiokawa, *Transcendence of numbers related with Cahen's constant*, to appear in *Moscow J. of Cobinatics and Number Theory*.
- [7] D. Duverney and I. Shiokawa, *Irrationality exponents of numbers related with Cahen's constant*, submitted, *Monographs in Number Theory* 4, World Scientific, 2010.
- [8] W. B. Jones and W. J. Thron, *Continued Fractions : analytic theory and applications*, *Encyclopedia of Math. and its applications* 11, Addison-Wesley, 1980.
- [9] W. Sierpinski, *Sur un algorithme pour développer les nombres réels en séries rapidement convergentes*, Bull. Intern. Acad. Sci. Cracovie (1911), 508-514.
- [10] J. Sondow, *Irrationality measures, irrationality bases and a theorem of Jarník*, <https://arxiv.org/abs/math/0406300v1> (2004).
- [11] J. J. Sylvester, *On a point in the theory of vulgar functions*, Amer. J. Math. 3 (1880), 332-335.

IEKATA SHIOKAWA,

E-mail address: shiokawa@beige.ocn.ne.jp