

# Superprime Rings

Hisaya Tsutsui

Department of Mathematics  
Embry-Riddle Aeronautical University  
Prescott, AZ USA  
E-mail address: [Hisaya.Tsutsui@erau.edu](mailto:Hisaya.Tsutsui@erau.edu)

The talk presented was a preliminary report and an introduction to the subject.

A ring in which every (two sided) ideal is an idempotent is called a fully idempotent ring. An example of such a ring includes the class of Von Neumann regular rings. In fact, it is well known and easy to show that every one sided ideal of Von Neumann regular rings is an idempotent.

If a ring  $R$  is fully idempotent, then for any ideals  $J, K$  of  $R$ ,  $J \cap K = JK$ . Hence an ideal  $P$  in a fully idempotent ring  $R$  is prime if and only if  $J \cap K \subseteq P$  implies  $J \subseteq P$ , or  $K \subseteq P$ . On the other hand, in the ring  $\mathbb{Z}_8$ ,  $\langle 2 \rangle \cap \langle 4 \rangle = \langle 4 \rangle$  but  $\langle 4 \rangle$  is not a prime ideal of  $\mathbb{Z}_8$ .

We define a prime ideal  $P$  in an arbitrary ring  $R$  to be superprime if

$\bigcap_{i \in I} J_i \subseteq P \Rightarrow J_i \subseteq P$  for some  $i$ , where  $J_i$  is an ideal of  $R$ . A ring in which  $0$  is superprime will

be called a superprime ring.

The speaker has long been investigated the structure of rings in which every ideal is prime.

An example of such rings includes the ring  $R$  of all linear transformations  $f: V \rightarrow V$  of a vector space  $V$  over a field  $F$ . We are mainly interested in the structure of fully prime rings with a superprime ideal.

**Theorem 1 [1, Theorem 1.2]:** A ring  $R$  is fully prime if and only if  $R$  is fully idempotent and ideals in  $R$  is linearly ordered.

**Proposition 2:** A superprime ring is primitive if and only if it is semiprimitive.

*Proof:* By definition, the intersection of all nonzero ideals of a superprime ring is nonzero, and

hence it is the minimal nonzero two sided ideal. If  $0$  is not a primitive ideal, the ring cannot be semiprimitive since the Jacobson radical must then contain the minimal nonzero two sided ideal.

A commutative fully prime ring is a field. Since a superprime ring is in particular prime, the minimal nonzero ideal is an idempotent. Hence, a commutative superprime ring is also a field. The center of a fully prime ring is either a field or zero ([1, Theorem 1.3]). We ask: what can we say about the center of a superprime ring?

It is wellknown that a prime ring with a minimal right ideal is primitive.

**Theorem 1:** A right Noetherian fully prime superprime ring  $R$  is primitive. Further, if  $R$  is not simple, then  $R$  contains no minimal right ideals.

*Sketch of a proof:* By Nakayama's lemma and Theorem 1,  $R$  is semiprimitive. Hence by Proposition 2,  $R$  is primitive. It can be shown that  $\text{Soc}(R)$  is either  $0$  or  $R$ . Suppose that  $\text{Soc}(R) \neq 0$ . Then since  $R$  is prime,  $\text{Soc}(R)$  is the intersection of nonzero ideals of  $R$ . Since  $R$  is not simple (but fully prime right Noetherian), we have a contradiction.

A prime semiprimitive but not a primitive ring is not superprime. The ring of integers is an obvious example. A Von Neumann regular ring is semiprimitive but there is a wellknown example of prime Von Neumann regular ring that is not primitive. We ask: is a semiprimitive fully prime ring superprimitive?

We conclude this preliminary report with the following conjecture: Let  $R$  be a fully prime ring. The following statements are equivalent:

- (a)  $R$  is primitive.
- (b)  $R$  is semiprimitive
- (c)  $R$  is superprime.

#### Reference

[1] W.D. Blair and H. Tsutsui, Fully Prime Rings, Comm. Algebra 22 (1994), no. 13, 5389-5400.