

# Isometries on a Banach space of analytic functions on the open unit disk

Takeshi Miura

Department of Mathematics, Niigata University

## 1 Introduction

Let  $(N, \|\cdot\|_N)$  be a normed linear space over  $\mathbb{R}$  or  $\mathbb{C}$ . A mapping  $T$  on  $(N, \|\cdot\|_N)$  is an *isometry* if

$$\|T(f) - T(g)\|_N = \|f - g\|_N \quad (\forall f, g \in N).$$

Here, we don't assume linearity of  $T$ . Let  $\mathbb{D}$  be the open unit disc and  $\mathbb{T}$  the unit circle in  $\mathbb{C}$ . We denote by  $H(\mathbb{D})$  the complex linear space of all analytic functions on  $\mathbb{D}$ . Let  $H^p$  be the Hardy space defined by

$$H^p = \left\{ f \in H(\mathbb{D}) : \|f\|_p = \sup_{0 < r < 1} \left[ \frac{1}{2\pi} \int_0^{2\pi} |f(re^{it})|^p dt \right]^{1/p} < \infty \right\} \quad (1 \leq p < \infty),$$

$$H^\infty = \left\{ f \in H(\mathbb{D}) : \|f\|_\infty = \sup_{z \in \mathbb{D}} |f(z)| < \infty \right\}.$$

Complex linear isometries on the Hardy spaces were characterized in 1960's.

**Theorem** (deLeeuw, Rudin and Wermer [1]). *1. Let  $T$  be a surjective, complex linear isometry on  $(H^\infty, \|\cdot\|_\infty)$ . Then there exist a constant  $\alpha \in \mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$  and a conformal map  $\phi: \mathbb{D} \rightarrow \mathbb{D}$  such that*

$$T(f)(z) = \alpha f(\phi(z)) \quad (\forall f \in H^\infty, z \in \mathbb{D}).$$

*2. Let  $T$  be a surjective, complex linear isometry on  $(H^1, \|\cdot\|_1)$ . Then there exist a constant  $\alpha \in \mathbb{T}$  and a conformal map  $\phi: \mathbb{D} \rightarrow \mathbb{D}$  such that*

$$T(f)(z) = \alpha \phi'(z) f(\phi(z)) \quad (\forall f \in H^1, z \in \mathbb{D}).$$

In 1959, Nagasawa [8] gave the characterization of surjective complex linear isometry on uniform algebras. The characterization of isometries on  $H^\infty$  by deLeeuw, Rudin and Wermer is a special case of the result by Nagasawa.

Forelli [3] investigated complex linear, not necessarily surjective, isometries on  $H^p$ . Here, I will introduce the result of surjective case.

**Theorem** (Forelli, [3]). *Let  $p$  be a real number with  $1 \leq p < \infty$  and  $p \neq 2$ , and let  $T$  be a surjective complex linear isometry on  $(H^p, \|\cdot\|_p)$ . There exist a constant  $\alpha \in \mathbb{T}$  and a conformal map  $\phi: \mathbb{D} \rightarrow \mathbb{D}$  such that*

$$T(f)(z) = \alpha(\phi'(z))^{1/p} f(\phi(z)) \quad (\forall f \in H^p, z \in \mathbb{D}).$$

Novinger and Oberlin [9] considered Banach spaces of analytic functions

$$\mathcal{S}^p = \{f \in H(\mathbb{D}) : f' \in H^p\} \quad (1 \leq p < \infty)$$

with the following norms:

$$\|f\|_\sigma = |f(0)| + \|f'\|_p, \quad \|f\|_\Sigma = \|f\|_\infty + \|f'\|_p \quad (f \in \mathcal{S}^p).$$

Here, it should be mentioned that  $\|f\|_\infty$  is well-defined; in fact, if a function  $f \in H(\mathbb{D})$  satisfies  $f' \in H^p$  for some  $p, 1 \leq p$  then  $f$  is extended to a continuous function on the closed unit ball  $\bar{\mathbb{D}}$  (see, for example [2, Theorem 3.11]). Novinger and Oberlin [9] characterized complex linear isometries on  $\mathcal{S}^p$  without assuming surjectivity. For the sake of simplicity, I will show you a surjective case of their results.

**Theorem** (Novinger and Oberlin [9]). *Let  $p$  be a real number with  $1 \leq p < \infty$  and  $p \neq 2$ .*

1. *If  $T$  is a surjective complex linear isometry on  $(\mathcal{S}^p, \|\cdot\|_\sigma)$ , then there exist a constant  $c \in \mathbb{T}$  and a conformal map  $\phi: \mathbb{D} \rightarrow \mathbb{D}$  such that*

$$T(f)(z) = cf(0) + \int_{[0,z]} (\phi'(\zeta))^{1/p} f'(\phi(\zeta)) d\zeta \quad (\forall f \in \mathcal{S}^p, z \in \mathbb{D}).$$

2. *If  $T$  is a surjective complex linear isometry on  $(\mathcal{S}^p, \|\cdot\|_\Sigma)$ , then there exist a constant  $c \in \mathbb{T}$  and a conformal map  $\phi: \mathbb{D} \rightarrow \mathbb{D}$  such that*

$$T(f)(z) = cf(\phi(z)) \quad (\forall f \in \mathcal{S}^p, z \in \mathbb{D}).$$

Novinger and Oberlin excluded the case when  $p = \infty$  in the above result. But  $\mathcal{S}^\infty$  is well-defined, and I believe the characterization of isometries on  $\mathcal{S}^\infty$  is important to the theory of analytic functions. The purpose of this note is to give an answer to the above problem.

## 2 Main results

We define  $\mathcal{S}^\infty = \{f \in H(\mathbb{D}) : f' \in H^\infty\}$ . As is mentioned above, if  $f \in \mathcal{S}^\infty$ , then it can be extended to a continuous function on  $\bar{\mathbb{D}}$ . Thus,  $\|f'\|_\infty$  is well-defined. We consider the following two norms on  $\mathcal{S}^\infty$ :

$$\|f\|_\sigma = |f(0)| + \|f'\|_\infty, \quad \|f\|_\Sigma = \|f\|_\infty + \|f'\|_\infty \quad (f \in \mathcal{S}^\infty).$$

We see that  $(\mathcal{S}^\infty, \|\cdot\|_\sigma)$  and  $(\mathcal{S}^\infty, \|\cdot\|_\Sigma)$  are both Banach spaces. Noviger and Oberlin characterized complex linear isometries on  $\mathcal{S}^p$  ( $1 \leq p < \infty$ ) without assuming surjectivity. Here we investigate *surjective*, not necessarily linear, isometries on  $\mathcal{S}^\infty$ . The main results of this note is as follows.

**Theorem 1.** *A map  $T$  is a surjective isometry on  $(\mathcal{S}^\infty, \|\cdot\|_\Sigma)$  if and only if there exist constants  $c, \lambda \in \mathbb{T}$  such that*

$$\begin{aligned} T(f)(z) &= T(0)(z) + cf(\lambda z) & (\forall f \in \mathcal{S}^p, z \in \mathbb{D}) \quad \text{or} \\ T(f)(z) &= T(0)(z) + c\overline{f(\overline{\lambda z})} & (\forall f \in \mathcal{S}^p, z \in \mathbb{D}). \end{aligned}$$

**Outline of proof.** By the Mazur-Ulam theorem [5], the map  $T_0 = T - T(0)$ , which sends  $f \in \mathcal{S}^\infty$  to  $T(f) - T(0)$ , is real linear. In addition, we see that  $T_0$  is a surjective isometry. Let  $\hat{f}'$  be the Gelfand transform of  $f' \in H^\infty$  and let  $\partial_{H^\infty}$  be the Shilov boundary for  $H^\infty$ . Then  $\sup_{\zeta \in \mathbb{D}} |f'(\zeta)| = \sup_{z \in \partial_{H^\infty}} |\hat{f}'(\zeta)|$  for  $f \in \mathcal{S}^\infty$ . We denote by  $\hat{f}$  the unique continuous extension of  $f \in \mathcal{S}^\infty$  to  $\mathbb{D}$ . By the maximal modulus principle,  $\sup_{z \in \mathbb{D}} |f(z)| = \sup_{z \in \mathbb{T}} |\hat{f}(z)|$  for  $f \in \mathcal{S}^\infty$ . Therefore

$$\|f\|_\Sigma = \sup_{z \in \mathbb{D}} |f(z)| + \sup_{\zeta \in \mathbb{D}} |f'(\zeta)| = \sup_{z \in \mathbb{T}} |\hat{f}(z)| + \sup_{\zeta \in \partial_{H^\infty}} |\hat{f}'(\zeta)| = \sup_{(z, w, \zeta) \in \mathbb{T}^2 \times \partial_{H^\infty}} |\hat{f}(z) + w\hat{f}'(\zeta)|.$$

We now define a map  $U: \mathcal{S}^\infty \rightarrow C(\mathbb{T}^2 \times \partial_{H^\infty})$  by

$$U(f)(z, w, \zeta) = \hat{f}(z) + w\hat{f}'(\zeta) \quad (\forall f \in \mathcal{S}^\infty, (z, w, \zeta) \in \mathbb{T}^2 \times \partial_{H^\infty}).$$

Set  $B = U(\mathcal{S}^\infty)$ , and then  $U$  is a surjective complex linear isometry from  $(\mathcal{S}^\infty, \|\cdot\|_\Sigma)$  onto  $(B, \|\cdot\|_\infty)$ .

$$\begin{array}{ccc} \mathcal{S}^\infty & \xrightarrow{T_0} & \mathcal{S}^\infty \\ U \downarrow & & \downarrow U \\ B & \xrightarrow[V]{} & B \end{array}$$

We set  $V = UT_0U^{-1}$ . Then  $V$  is a surjective *real linear* isometry on  $(B, \|\cdot\|_\infty)$ .

By a modified arguments of [10, Proof of Lemma 3.1], we can prove that

$$V_*(\{\lambda\delta_x : \lambda \in \mathbb{T}, x \in \mathbb{T}^2 \times \partial_{H^\infty}\}) = \{\lambda\delta_x : \lambda \in \mathbb{T}, x \in \mathbb{T}^2 \times \partial_{H^\infty}\},$$

where  $V_*: B^* \rightarrow B^*$  is a map defined by

$$V_*(\eta)(a) = \operatorname{Re} \eta(V(a)) - i\operatorname{Re} \eta(V(ia)) \quad (\forall \eta \in B^*, a \in B),$$

and  $\delta_x: B \rightarrow \mathbb{C}$  is a point evaluation functional with  $\delta_x(a) = a(x)$  for  $a \in B$ . Using the form of  $V$ , we can describe  $T_0$  with extra variables, say  $w \in \mathbb{T}$  and  $\zeta \in \partial_{H^\infty}$ . By straightforward, but complicated arguments, we obtain the desired form of  $T$ . The reader may refer to [7] for the detail.  $\square$

**Theorem 2.** *Let  $T$  be a surjective isometry on  $(\mathcal{S}^\infty, \|\cdot\|_\sigma)$ . Then there exist constants  $c_0, c_1, \lambda \in \mathbb{T}$  and  $a \in \mathbb{D}$  such that*

$$T(f)(z) = T(0)(z) + c_0 f(0) + \int_{[0,z]} c_1 f' \left( \lambda \frac{z-a}{1-\bar{a}\zeta} \right) d\zeta \quad (\forall f \in \mathcal{S}^p, z \in \mathbb{D}) \quad \text{or}$$

$$T(f)(z) = T(0)(z) + c_0 \overline{f(0)} + \int_{[0,z]} c_1 f' \left( \lambda \frac{z-a}{1-\bar{a}\zeta} \right) d\zeta \quad (\forall f \in \mathcal{S}^p, z \in \mathbb{D}) \quad \text{or}$$

$$T(f)(z) = T(0)(z) + c_0 f(0) + \int_{[0,z]} c_1 \overline{f' \left( \lambda \frac{z-a}{1-\bar{a}\zeta} \right)} d\zeta \quad (\forall f \in \mathcal{S}^p, z \in \mathbb{D}) \quad \text{or}$$

$$T(f)(z) = T(0)(z) + c_0 \overline{f(0)} + \int_{[0,z]} c_1 \overline{f' \left( \lambda \frac{z-a}{1-\bar{a}\zeta} \right)} d\zeta \quad (\forall f \in \mathcal{S}^p, z \in \mathbb{D}).$$

*Conversely, if  $T$  is one of the above four, then it is a surjective isometry on  $(\mathcal{S}^\infty, \|\cdot\|_\sigma)$ .*

**Outline of proof.** The idea of this proof is quite similar to that of Theorem 1. We need the characterization of surjective, real linear isometries on uniform algebras (see [4, 6]).  $\square$

## References

- [1] K. deLeeuw, W. Rudin and J. Wermer, *The isometries of some function spaces*, Proc. Amer. Math. Soc. **11** (1960), 694–698.
- [2] P.L. Duren, *The theory of  $H^p$  spaces*, Pure and Applied Mathematics, Vol. 38 Academic Press, New York-London, 1970.
- [3] F. Forelli, *The isometries of  $H^p$* , Canad. J. Math. **16** (1964), 721–728.
- [4] O. Hatori and T. Miura, *Real linear isometries between function algebras. II*, Cent. Eur. J. Math. **11** (2013), 1838–1842.
- [5] S. Mazur and S. Ulam, *Sur les transformations isométriques d'espaces vectoriels normés*, C. R. Acad. Sci. Paris **194** (1932), 946–948.
- [6] T. Miura, *Real-linear isometries between function algebras*, Cent. Eur. J. Math. **9** (2011), 778–788.
- [7] T. Miura, *Surjective isometries on a Banach space of analytic functions on the open unit disc*, preprint.
- [8] M. Nagasawa, *Isomorphisms between commutative Banach algebras with an application to rings of analytic functions*, Kōdai Math. Sem. Rep. **11** (1959), 182–188.

- [9] W.P. Novinger and D.M. Oberlin, *Linear isometries of some normed spaces of analytic functions*, Can. J. Math. **37** (1985), 62–74.
- [10] N.V. Rao and A.K. Roy, *Linear isometries of some function spaces*, Pacific J. Math. **38** (1971), 177–192.

Department of Mathematics  
Niigata University  
Niigata 950-2181  
JAPAN  
E-mail address: miura@math.sc.niigata-u.ac.jp