

On the Feigin-Tipunin conjecture

SHOMA SUGIMOTO*†

Research Institute For Mathematical Sciences, Kyoto University

1 Introduction

The triplet W -algebra (= type A_1 logarithmic W -algebra) ([AM1]-[AM3], [FGST1]-[FGST3], [NT], [TW], ...) is one of the most famous examples of C_2 -cofinite but irrational vertex operator algebra, and it relates to many interesting objects such as the tails of colored Jones polynomials and false theta functions [BM1, CCFGH, MN], quantum groups at root of unity [CGR, FGR, NT], and the quantum geometric Langlands program [CG, Cr1]. We can immediately generalize the definition of the triplet W -algebra to type ADE cases, and we call them the type ADE logarithmic W -algebras $W(p)_Q$. However, very little is known about the properties and the representation theory of the higher rank generalizations of the triplet W -algebra.

In [FT], without detailed proofs, they claimed that $W(p)_Q$ and its irreducible modules are constructed as the spaces of global sections of some homogeneous vector bundles over the flag variety, and we call it Feigin-Tipunin conjecture. In [S1, S2], the author proved it partially and obtained some of new results on the type ADE logarithmic W -algebras.

In this paper, with some comments and remarks, we gather results that will given in [S1, S2]. We give the geometric construction of the type ADE logarithmic W -algebra $W(p)_Q$ that claimed in [FT]. This construction reveals us the G -module structure and the character formula of $W(p)_Q$. Moreover, under the assumption of simpleness of $W(p)_Q$, we also completely determine the $\mathcal{W}^k(\mathfrak{g})$ -module structure of $W(p)_Q$. Finally, applying this result to the cases of type A_2 with small $p \in \mathbb{Z}_{\geq 2}$, we prove the C_2 -cofiniteness of $W(p)_Q$ in these cases under the assumption of simpleness.

1.1 Setting

Let \mathfrak{g} be a simply-laced simple Lie algebra of rank l , and $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$ the triangular decomposition, \mathfrak{h} the Cartan subalgebra, $\mathfrak{b} = \mathfrak{n}_- \oplus \mathfrak{h}$ the Borel subalgebra, G , H , and B the semisimple, simply-connected, complex algebraic groups corresponding to \mathfrak{g} , \mathfrak{h} , \mathfrak{b} , respectively. We adopt the standard numbering for the simple roots $\{\alpha_1, \dots, \alpha_l\}$ of \mathfrak{g} as in [B] and denote by $\{\omega_1, \dots, \omega_l\}$ the corresponding fundamental weights, and denote by Π denotes the set of simple roots. Let Q be the root lattice of \mathfrak{g} , P the weight lattice of \mathfrak{g} , P_+ the set of dominant integral weights of \mathfrak{g} . Denote by (\cdot, \cdot) the normalized invariant form of \mathfrak{g} , W the Weyl group of \mathfrak{g} generated by the simple reflections $\{\sigma_i\}_{i=1}^l$, (c_{ij}) the Cartan matrix of \mathfrak{g}

* Research Institute For Mathematical Sciences, Kyoto University, Kyoto 606-8502 JAPAN

† shoma@kurims.kyoto-u.ac.jp

and (c^{ij}) the inverse matrix to (c_{ij}) , ρ the half sum of positive roots, h the Coxeter number of \mathfrak{g} , Ω the abelian group P/Q . We choose the representatives of generators of Ω in P in the following way: for A_l , D_l , E_6 , E_7 , E_8 , we choose $\{0, \omega_1, \dots, \omega_l\}$, $\{0, \omega_1, \omega_{l-1}, \omega_l\}$, $\{0, \omega_1, \omega_3\}$, $\{0, \omega_2\}$, $\{0\}$, respectively. We fix an integer $p \in \mathbb{Z}_{\geq 2}$.

Let $V_{\sqrt{p}Q} = \bigoplus_{\alpha \in Q} \mathcal{F}_{\sqrt{p}\alpha}$ be the lattice vertex operator algebra associated to the rescaled root lattice $\sqrt{p}Q$, where $\mathcal{F}_{\sqrt{p}\alpha} = \mathcal{U}(\hat{\mathfrak{h}}^{<0}) \otimes |\sqrt{p}\alpha\rangle$ is the Fock module of the Heisenberg vertex operator algebra $\mathcal{F}_0 = \mathcal{U}(\hat{\mathfrak{h}}^{<0}) \otimes |0\rangle$.

We choose the shifted conformal vector ω of $V_{\sqrt{p}Q}$ as

$$\omega = \frac{1}{2} \sum_{1 \leq i, j \leq l} c^{ij} (\alpha_i)_{(-1)} \alpha_j + Q_0(\rho)_{(-2)} |0\rangle \in \mathcal{F}_0 \subseteq V_{\sqrt{p}Q}, \quad (1)$$

where $Q_0 = \sqrt{p} - \frac{1}{\sqrt{p}}$. The central charge c of ω is given by

$$c = l + 12(\rho, \rho)(2 - p - \frac{1}{p}) = l + h \dim \mathfrak{g}(2 - p - \frac{1}{p}). \quad (2)$$

For $n \in \mathbb{Z}$, we use the traditional notation L_n for the Virasoro operator $\omega_{(n+1)}$.

Irreducible modules over $V_{\sqrt{p}Q}$ are classified by elements of the abelian group $\Lambda = \frac{1}{\sqrt{p}}P/\sqrt{p}Q$ ([D]). For each equivalence class $\langle \lambda \rangle \in \Lambda$, we choose the unique representative $\lambda \in \frac{1}{\sqrt{p}}P$ of $\langle \lambda \rangle \in \Lambda$ as

$$\lambda = -\sqrt{p}\hat{\lambda} + \bar{\lambda} = -\sqrt{p}\hat{\lambda} + \sum_{j=1}^l \frac{s_j}{\sqrt{p}} \omega_j, \quad (3)$$

where $0 \leq s_j \leq p-1$, $\hat{\lambda} \in \Omega$ and the representatives of generators of Ω are given in above: for A_l , D_l , E_6 , E_7 , E_8 , we have $\{0, \omega_1, \dots, \omega_l\}$, $\{0, \omega_1, \omega_{l-1}, \omega_l\}$, $\{0, \omega_1, \omega_3\}$, $\{0, \omega_2\}$, $\{0\}$, respectively. For $\lambda \in \frac{1}{\sqrt{p}}P$, denote by $V_{\sqrt{p}Q+\lambda}$ the irreducible $V_{\sqrt{p}Q}$ -module

$$V_{\sqrt{p}Q+\lambda} = \bigoplus_{\alpha \in Q} \mathcal{F}_{\sqrt{p}\alpha+\lambda}. \quad (4)$$

corresponding to $\langle \lambda \rangle \in \Lambda$, where $\mathcal{F}_{\sqrt{p}\alpha+\lambda}$ is the Fock module over \mathcal{F}_0 with the highest weight vector $|\sqrt{p}\alpha + \lambda\rangle$. For $\mu \in \frac{1}{\sqrt{p}}P$, the conformal weight Δ_μ of $|\mu\rangle$ is

$$\Delta_\mu = \frac{1}{2} |\mu - Q_0\rho|^2 + \frac{c-l}{24} = \frac{1}{2} |\mu|^2 - Q_0(\mu, \rho). \quad (5)$$

1.2 Screening and narrow screening

For $1 \leq i \leq l$, $\alpha \in Q$ and $\lambda \in \Lambda$, we consider the *screening operators*

$$f_i = |\sqrt{p}\alpha_i\rangle_{(0)} \in \text{Hom}(\mathcal{F}_{-\sqrt{p}\alpha+\lambda}, \mathcal{F}_{-\sqrt{p}(\alpha+\alpha_i)+\lambda}). \quad (6)$$

For $\sigma \in W$ and $\mu \in \frac{1}{\sqrt{p}}P$, set

$$\sigma \star \mu = \sigma(\mu + \frac{1}{\sqrt{p}}\rho) - \frac{1}{\sqrt{p}}\rho. \quad (7)$$

Then we have the following W -action on Λ :

$$\sigma \star \lambda = -\sqrt{p}\hat{\lambda} + \sigma \star \bar{\lambda}. \quad (8)$$

In order to define the *narrow screening operators* $F_{i,\lambda} \in \text{Hom}(\mathcal{F}_{-\sqrt{p}\alpha+\lambda}, \mathcal{F}_{-\sqrt{p}\alpha+\sigma_i*\lambda})$ for $1 \leq i \leq l$, we consider the following element in $\text{Hom}(V_{\sqrt{p}Q}, V_{\sqrt{p}Q-\frac{\alpha_i}{\sqrt{p}}}) \otimes \mathbb{C}[[z^\pm]]$:

$$F_i(z) = e^{-\frac{\alpha_i}{\sqrt{p}}z} z^{-\frac{(\alpha_i)_0}{\sqrt{p}}} \exp\left(\sum_{n<0} \frac{z^{-n}}{n} \frac{(\alpha_i)_{(n)}}{\sqrt{p}}\right) \exp\left(\sum_{n>0} \frac{z^{-n}}{n} \frac{(\alpha_i)_{(n)}}{\sqrt{p}}\right) c_{-\frac{\alpha_i}{\sqrt{p}}}. \quad (9)$$

Here the element $e^{-\frac{\alpha_i}{\sqrt{p}}} \in \text{Hom}(V_{\sqrt{p}Q}, V_{\sqrt{p}Q-\frac{\alpha_i}{\sqrt{p}}})$ is defined by

$$\begin{cases} e^{-\frac{\alpha_i}{\sqrt{p}}} |\sqrt{p}\mu\rangle = |-\frac{\alpha_i}{\sqrt{p}} + \sqrt{p}\mu\rangle, \\ [h_{(n)}, e^{-\frac{\alpha_i}{\sqrt{p}}}] = \delta_{n,0}(h, -\frac{\alpha_i}{\sqrt{p}}) e^{-\frac{\alpha_i}{\sqrt{p}}}, \end{cases} \quad (10)$$

for $\mu \in Q$ and $h_{(n)} \in \mathcal{U}(\hat{\mathfrak{h}})$, and the element $c_{-\frac{\alpha_i}{\sqrt{p}}} \in \text{Hom}(V_{\sqrt{p}Q}, V_{\sqrt{p}Q-\frac{\alpha_i}{\sqrt{p}}})$ is defined by

$$c_{-\frac{\alpha_i}{\sqrt{p}}} s |\sqrt{p}\mu\rangle = \epsilon'(-\alpha_i, \mu) s |\sqrt{p}\mu\rangle. \quad (11)$$

Here $s \in \mathcal{U}(\hat{\mathfrak{h}}^{<0})$ and $\epsilon' : Q \times Q \rightarrow \mathbb{C}^\times$ is the 2-cocycle defined by

$$\epsilon(\alpha_i, \alpha_j) = \begin{cases} (-1) & i = j, \\ (-1)^{(\alpha_i, \alpha_j)} & i < j, \\ 1 & i > j. \end{cases} \quad (12)$$

For $\alpha \in Q$, the narrow screening operator is given by the z^{-1} coefficient of $F_i(z)$

$$F_{i,0} = \int F_i(z) dz \in \text{Hom}(\mathcal{F}_{-\sqrt{p}\alpha}, \mathcal{F}_{-\sqrt{p}\alpha-\frac{\alpha_i}{\sqrt{p}}}). \quad (13)$$

Denote by \mathcal{F}_0^i the rank 1 Heisenberg vertex algebra generated by α_i , and $\mathcal{F}_0^{i,\perp}$ the rank $l-1$ Heisenberg vertex algebra generated by $\{\omega_j\}_{1 \leq j \neq i \leq l}$, respectively. Then for $a \in \mathcal{F}_0^{i,\perp}$ and $n \in \mathbb{Z}$, we have $F_{i,0} a_{(n)} = a_{(n)} F_{i,0}$. By applying the multiplication of narrow screening operators in the case of type A_1 (see [CRW, NT]) to $F_{i,0}|_{\mathcal{F}_0^i}$, for $\alpha \in Q$ and $\lambda \in \Lambda$ such that $0 \leq s_i \leq p-2$, we have the non-trivial map

$$F_{i,\lambda} = \int_{[\Gamma_{s_i+1}]} F_i(z_1) \dots F_i(z_{s_i+1}) dz_1 \dots dz_{s_i+1} \in \text{Hom}(\mathcal{F}_{-\sqrt{p}\alpha+\lambda}, \mathcal{F}_{-\sqrt{p}\alpha+\sigma_i*\lambda}),$$

where the cycle $[\Gamma_{s_i+1}]$ such that $F_{i,\lambda}$ to be non-trivial is uniquely determined up to normalization. For convenience, we set $F_{i,\lambda} = 0$ for $\lambda \in \Lambda$ such that $s_i = p-1$.

Clearly, the screening and narrow screening operators are differential operators on $V_{\sqrt{p}Q}$ because they are zero modes. In other words, they satisfy the Leibniz rule

$$f_i u_{(n)} v = (f_i u)_{(n)} v + u_{(n)} f_i v, \quad (14)$$

$$\begin{aligned} F_{i,0} a_{(n)} b &= (F_{i,0} a)_{(n)} b + a_{(n)} F_{i,0} b \\ &= \sum_{m \geq 0} \frac{(-1)^{-n-m-1}}{m!} T^m b_{(n+m)} F_{i,0} a + a_{(n)} F_{i,0} b, \end{aligned} \quad (15)$$

where $n \in \mathbb{Z}$, $u, v \in V_{\sqrt{p}Q+\lambda}$ and $a, b \in V_{\sqrt{p}Q}$. Moreover, a straightforward calculation shows that

$$[f_i, L_n] = [F_{i,\lambda}, L_n] = 0 \quad (16)$$

and

$$[f_i, F_{j,\lambda}] = 0 \quad (17)$$

for $1 \leq i, j \leq l$ and $n \in \mathbb{Z}$. In particular, (16) means that f_i and $F_{i,\lambda}$ preserve the conformal grading.

1.3 Logarithmic W -algebra

Since every $F_{i,0}$ satisfies (15), we have the vertex operator subalgebra

$$W(p)_Q = \bigcap_{i=1}^l \ker F_{i,0}|_{V_{\sqrt{p}Q}} \subseteq V_{\sqrt{p}Q}. \quad (18)$$

By (16), ω is a conformal vector of $W(p)_Q$. This vertex operator algebra is called the *logarithmic W -algebra* associated to Q and p . In particular, in the case of type A_1 , $W(p)_Q$ is the *triplet W -algebra* ([AM1]-[AM3], [FGST1]-[FGST3], [NT], [TW], ...).

For $1 \leq i \leq l$, we consider the following operator $h_{i,\lambda}$ acting on $V_{\sqrt{p}Q+\lambda}$:

$$h_{i,\lambda} = -\frac{1}{\sqrt{p}}(\alpha_i)_{(0)} + \frac{1}{\sqrt{p}}(\alpha_i, \bar{\lambda}) \text{id}. \quad (19)$$

Theorem 1 ([FT, Theorem 4.1]).

1. The operators $\{f_i, h_{i,\lambda}\}_{i=1}^l$ give rise to an action of \mathfrak{b} on $V_{\sqrt{p}Q+\lambda}$.
2. The action of \mathfrak{b} in (1) is integrable.

For $\lambda \in \Lambda$, we consider the homogeneous vector bundle

$$\xi_\lambda = G \times_B V_{\sqrt{p}Q+\lambda} \quad (20)$$

over the flag variety G/B , where the action of B on G is given by the right multiplication and that on $V_{\sqrt{p}Q,\lambda}$ is given by Theorem 1. We can easily show that the space of global sections $H^0(\xi_0)$ inherits the vertex operator algebra structure from $V_{\sqrt{p}Q}$, and each $H^0(\xi_\lambda)$ is a $H^0(\xi_0)$ -module as in the same way.

1.4 Results

Definition 1. For $\lambda \in \Lambda$ and $\sigma \in W$, set

$$\epsilon_\lambda(\sigma) = \frac{1}{\sqrt{p}}(\sigma \star \bar{\lambda} - \overline{\sigma \star \lambda}). \quad (21)$$

Let J be a subset of nodes of the Dynkin diagram of G and $\lambda \in \Lambda$. The pairing (J, λ) is *good* if $\epsilon_\lambda(\sigma_j) = -\alpha_j$ for any $j \in J$ or $(\epsilon_\lambda(\sigma_j), \alpha_i) = -\delta_{i,j}$ for any $i, j \in J$. In particular, when $J = \Pi$, we call λ is good if (Π, λ) is good.

Remark 1. If $|J| = 1$, (J, λ) is good.

The following three theorems will be given in [S1].

Theorem 2. 1. For $p \in \mathbb{Z}_{\geq 2}$, we have the vertex operator algebra isomorphism

$$H^0(\xi_0) \simeq W(p)_Q.$$

In particular, the group G acts on $W(p)_Q$ as an automorphism group.

2. More generally, if λ is good, we have the $W(p)_Q$ -module and G -module isomorphism

$$H^0(\xi_\lambda) \simeq \bigcap_{i=1}^l \ker F_{i,\lambda}|_{V_{\sqrt{p}Q+\lambda}}.$$

3. If λ is not good, $H^0(\xi_\lambda)$ is properly embedded into $\bigcap_{i=1}^l \ker F_{i,\lambda}|_{V_{\sqrt{p}Q+\lambda}}$.

Theorem 3. We have the vertex operator algebra isomorphism

$$\mathcal{W}^k(\mathfrak{g}) \simeq \bigcap_{i=1}^l \ker f_i|_{\mathcal{F}_0},$$

where $\mathcal{W}^k(\mathfrak{g})$ is the affine W -algebra $[FF]$ of level $k = p - h$

Theorem 4.

1. Let \mathcal{R}_μ be the irreducible \mathfrak{g} -module with the highest weight $\mu \in P_+$. Then we have the $W(p)_Q$ -module and G -module isomorphism

$$H^0(\xi_\lambda) \simeq \bigoplus_{\alpha \in P_+ \cap Q} \mathcal{R}_{\alpha+\tilde{\lambda}} \otimes \mathcal{W}_{-\sqrt{p}\alpha+\lambda} \subseteq V_{\sqrt{p}Q+\lambda} \quad (22)$$

where $\mathcal{W}_{-\sqrt{p}\alpha+\lambda} = \bigcap_{i=1}^l (\ker f_i|_{\mathcal{F}_0} - \sqrt{p}\alpha + \lambda)$ and $\ker f_i|_{\mathcal{F}_0} - \sqrt{p}\alpha + \lambda$ is the $\ker f_i|_{\mathcal{F}_0}$ -module generated by the highest weight vector $|\!-\sqrt{p}\alpha + \lambda\rangle$. In particular, we have

$$W(p)_Q \simeq \bigoplus_{\alpha \in P_+ \cap Q} \mathcal{R}_\alpha \otimes \mathcal{W}_{-\sqrt{p}\alpha},$$

and $\mathcal{W}^k(\mathfrak{g}) \simeq \mathcal{R}_0 \otimes \mathcal{W}_0$ is the vertex operator full subalgebra of $W(p)_Q$.

2. Let us fix $\lambda \in \Lambda$, and a minimal expression of $w_0 = \sigma_{i_N} \dots \sigma_{i_1}$. If $(\epsilon_\lambda(\sigma_{i_k} \dots \sigma_{i_1}), \alpha_{i_{k+1}}) = 0$ for $1 \leq k \leq N - 1$, then we have $H^k(\xi_\lambda) = 0$ for $k \geq 1$. In particular, if $\tilde{\lambda} = 0$, then $H^k(\xi_\lambda) = 0$ for $k \geq 1$. Moreover, we have the character formula

$$\begin{aligned} \mathrm{Tr}_{H^0(\xi_\lambda)}(q^{L_0 - \frac{c}{24}} z_1^{h_{1,\lambda}} \dots z_l^{h_{l,\lambda}}) &= \sum_{\alpha \in P_+ \cap Q} \chi_{\alpha+\tilde{\lambda}}^{\mathfrak{g}}(z) \left(\sum_{\sigma \in W} (-1)^{l(\sigma)} \frac{q^{\frac{1}{2}|\sqrt{p}\sigma(\alpha+\rho+\tilde{\lambda})-\tilde{\lambda}-\frac{1}{\sqrt{p}}\rho|^2}}{\eta(q)^l} \right) \\ &= \sum_{\alpha \in P_+ \cap Q} \chi_{\alpha+\tilde{\lambda}}^{\mathfrak{g}}(z) \mathrm{Tr}_{H^0_{DS,\alpha+\tilde{\lambda}}(\mathbb{V}_{p,\sqrt{p}\tilde{\lambda}})}(q^{L_0 - \frac{c}{24}}), \end{aligned}$$

where $\chi_{\beta}^{\mathfrak{g}}(z)$ be the Weyl character of \mathcal{R}_β , $l(\sigma)$ the length of $\sigma \in W$, $\eta(q)$ the Dedekind eta function, and $H^0_{DS,\alpha+\tilde{\lambda}}(\mathbb{V}_{p,\sqrt{p}\tilde{\lambda}})$ is the $\mathcal{W}^k(\mathfrak{g})$ -module defined in [ArF].

Remark 2. The author believe that the assumption $(\epsilon_\lambda(\sigma_{i_k} \dots \sigma_{i_1}), \alpha_{i_{k+1}}) = 0$ for $1 \leq k \leq N - 1$ in Theorem 4.2 is not necessary: i.e. he expect that $H^k(\xi_\lambda) = 0$ and the character formula above hold for all $\lambda \in \Lambda$ and $k \geq 1$. However, because of some technical difficulty in the proof of vanishing of higher cohomologies, he proved them on the restricted cases.

The following three theorems will be given in [S2].

Theorem 5. If $H^0(\xi_\lambda)$ is an irreducible $W(p)_Q$ -module, then $\mathcal{W}_{-\sqrt{p}\alpha+\lambda} \simeq \mathcal{W}^k(\mathfrak{g})|\!-\sqrt{p}\alpha + \lambda$. In other words, $\bigcap_{i=1}^l (\ker f_i|_{\mathcal{F}_0} - \sqrt{p}\alpha + \lambda) = (\bigcap_{i=1}^l \ker f_i|_{\mathcal{F}_0})|\!-\sqrt{p}\alpha + \lambda$. Moreover, when $\lambda = 0$, $\mathcal{W}_{-\sqrt{p}\alpha}$ is the irreducible $\mathcal{W}^k(\mathfrak{g})$ -module.

Definition 2. 1. For $\alpha \in P_+ \cap Q$, let H_α be a nonzero element of $\mathcal{R}_{\alpha,0} \otimes \mathbb{C}|\!-\sqrt{p}\alpha\rangle$, where $\mathcal{R}_{\alpha,0}$ is the space of zero-weight vectors of \mathcal{R}_α .

2. Let $\{W_i\}_{i=2}^{l+1}$ be strong generators of $\mathcal{W}^k(\mathfrak{g})$ such that $\Delta_{W_i} = i$. We use the notation

$$(W_i)_n = (W_i)_{(n+i-1)} \quad (23)$$

for $n \in \mathbb{Z}$. However, we often use the notation $(W_2)_n$ but L_n for traditional reason. Moreover, for a fixed $\alpha \in P_+ \cap Q$, we can assume that

$$(W_i)_0 | - \sqrt{p}\alpha = 0 \quad (24)$$

for $3 \leq i \leq l+1$ by considering the new strong generators

$$\{\omega\} \cup \{W_i - \nabla_{\alpha, i}\omega\}_{i=3}^{l+1} \quad (25)$$

of $\mathcal{W}^k(\mathfrak{g})$, where $\nabla_{\alpha, i} \in \mathbb{C}$ is defined by $(W_i)_0 | - \sqrt{p}\alpha = \nabla_{\alpha, i} | - \sqrt{p}\alpha$.

3. For $a \in W(p)_Q \simeq \bigoplus_{\alpha \in P_+ \cap Q} \mathcal{R}_\alpha \otimes W_{-\sqrt{p}\alpha}$, denote by $\tilde{a} \in \mathcal{W}^k(\mathfrak{g})$ be the $\mathcal{W}^k(\mathfrak{g})$ -component of a .

Theorem 6. 1. For the projection to the C_2 -algebra $\pi : W(p)_Q \rightarrow R_{W(p)_Q} = W(p)_Q / C_2(W(p)_Q)$, we have $\dim \pi(W(p)_Q \setminus \mathcal{W}^k(\mathfrak{g})) < \infty$. In other words, if $a \in W(p)_Q \setminus \mathcal{W}^k(\mathfrak{g})$, then $\pi(a)$ is nilpotent. In particular, for $\alpha \in P_+ \cap Q$, $\alpha \neq 0$, $\pi(H_\alpha)$ is nilpotent.

2. If $W(p)_Q$ is simple, then $W(p)_Q$ is strongly generated by $\{W_i\}_{i=2}^{l+1}$ and finitely many elements in $W(p)_Q \setminus \mathcal{W}^k(\mathfrak{g})$. In particular, if $W(p)_Q$ is simple and all $\pi(W_i)$ are nilpotent, then $W(p)_Q$ is C_2 -cofinite.

3. For $\alpha \in P_+ \cap Q$, $\{\widetilde{(H_\alpha)_{(N)} H_{\alpha'}}\}_{N \in \mathbb{Z}}$ satisfy the following conditions:

(a) For $m \geq 0$, we have

$$(\widetilde{(H_\alpha)_{(N+m)} H_{\alpha'}}) = \frac{L_m(\widetilde{(H_\alpha)_{(N)} H_{\alpha'}})}{(m+1)(\Delta_{-\sqrt{p}\alpha} - 1) - N + \delta_{m,0}\Delta_{-\sqrt{p}\alpha}}. \quad (26)$$

(b) Moreover, for $3 \leq i \leq l+1$, $n \geq i-1$ and $N \in \mathbb{Z}$, we have

$$\sum_{k=0}^{i-1} (-1)^k \binom{i-1}{k} \frac{(W_i)_{n-k} L_k(\widetilde{(H_\alpha)_{(N)} H_{\alpha'}})}{(k+1)(\Delta_{-\sqrt{p}\alpha} - 1) - N + \delta_{k,0}\Delta_{-\sqrt{p}\alpha}} = 0. \quad (27)$$

4. In the cases of types A_1 or A_2 , the conditions (26) and (27) determines $\{\widetilde{(H_\alpha)_{(N)} H_{\alpha'}}\}_{N \in \mathbb{Z}}$ uniquely up to scalar. Moreover, if $W(p)_Q$ is simple, then the conditions (26) and (27) determines $\{\widetilde{(H_\alpha)_{(N)} H_{\alpha'}}\}_{N \in \mathbb{Z}}$ uniquely up to nonzero scalar.

Remark 3. Theorem 6 claims that if $W(p)_Q$ is simple, the conditions (26) and (27) give an algorithm that enables us to calculate the nilpotent ideal in $\pi(\mathcal{W}^k(\mathfrak{g}))$ much easier than direct calculation. Applying it to the cases of type A_2 with small p , we obtain the following:

Theorem 7. Let us consider the cases when $\mathfrak{g} = \mathfrak{sl}_3$ and $p = 2, 3, 4$. If $W(p)_Q$ is simple, then $W(p)_Q$ is C_2 -cofinite.

参考文献

- [AM1] D. Adamovic and A. Milas. On the triplet vertex algebra $W(p)$, *Advances in Mathematics* 217 (2008), 2664-2699.
- [AM2] D. Adamovic and A. Milas. The $N=1$ triplet vertex operator superalgebras, *Communications in Mathematical Physics*, 288 (2009), 225-270.
- [AM3] D. Adamovic and A. Milas. The structure of Zhu's algebras for certain W -algebras, *Advances in Mathematics*, 227 (2011) 2425-2456.

- [ArF] T. Arakawa, E. Frenkel. Quantum Langlands duality of representations of \mathcal{W} -algebras, arXiv:1807.01536 [math.QA].
- [B] N. Bourbaki. Lie Groups and Lie Algebras. Chapter 4-6, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 2002, Translated from the 1968 French original by Andrew Pressly. MR 1890629
- [BM1] K. Bringmann and A. Milas. W -algebras, false theta functions and quantum modular forms I, International Mathematical Research Notices 21 (2015), 11351-11387.
- [CG] T. Creutzig, D. Gaiotto. Vertex Algebras for S-duality, arXiv:1708.00875.
- [CGR] T. Creutzig, A. M. Gainutdinov, I. Runkel. A quasi-Hopf algebra for the triplet vertex operator algebra, arXiv:1712.07260.
- [Cr1] T. Creutzig. Logarithmic W -algebras and Argyres-Douglas theories at higher rank, arXiv:1809.01725.
- [CRW] T. Creutzig, D. Ridout, S. Wood. Coset Construction of Logarithmic $(1, p)$ Models, *Letters in Mathematical Physics*, May(2014), Vol.104, 553-583.
- [CCFGH] M.C.N. Cheng, S. Chun, F. Ferrari, S. Gukov, S.M. Harrison. 3d Modularity, arXiv:1809.10148
- [D] C. Dong. Vertex algebras associated with even lattices, *J. Algebra* 161 (1993) 245-265.
- [FF] B. Feigin and E. Frenkel. Quantization of the Drinfeld-Sokolov reduction. *Phys. Lett. B*, 246(1-2): 75-81, 1990.
- [FGR] V. Farsad, A. M. Gainutdinov, I. Runkel. $SL(2, \mathbb{Z})$ -action for ribbon quasi-Hopf algebras, *Journal of Algebras*, March(2019), Vol.522, 243-308.
- [FGST1] B. L. Feigin, A.M. Gainutdinov, A. M. Semikhatov, and I. Yu Tipunin. The Kazhdan-Lusztig correspondence for the representation category of the triplet W -algebra in logarithmic conformal field theories. (Russian) *Teoret. Mat. Fiz.* 148 (2006), no. 3, 398-427.
- [FGST2] B. L. Feigin, A.M. Gainutdinov, A. M. Semikhatov, and I. Yu Tipunin. Logarithmic extensions of minimal models: characters and modular transformations. *Nuclear Phys. B* 757 (2006), 303-343.
- [FGST3] B. L. Feigin, A. M. Gainutdinov, A. M. Semikhatov, and I. Yu Tipunin. Modular group representations and fusion in logarithmic conformal field theories and in the quantum group center. *Comm. Math. Phys.* 265 (2006), 47-93.
- [FT] B. L. Feigin, I. Yu. Tipunin. Logarithmic CFTs connected with simple Lie algebras, arXiv:1002.5047.
- [MN] J. Murakami, K. Nagatomo. Logarithmic knot invariants arising from restricted quantum groups, *International Journal of Mathematics*, Vol.19, No.10, 1203-1213, 2008.
- [NT] K. Nagatomo and A. Tsuchiya. The Triplet Vertex Operator Algebra $W(p)$ and the Restricted Quantum Group at Root of Unity, Exploring new structures and natural constructions in mathematical physics, 149, *Adv. Stud. Pure Math.*, 61, Math. Soc. Japan, Tokyo, 2011., arXiv:0902.4607.
- [S1] S. Sugimoto. On the Feigin-Tipunin conjecture, In preparation.
- [S2] S. Sugimoto. In preparation.
- [TW] A. Tsuchiya. S. Wood. On the extended W -algebra of type \mathfrak{sl}_2 at positive rational level, *International Mathematics Research Notices*, Vol.2015, 14, 5357-5435.