

A NOTE ON LÖWENHEIM-SKOLEM CARDINALS

TOSHIMICHI USUBA

ABSTRACT. In this note we provide some applications of Löwenheim-Skolem cardinals introduced in [3].

1. INTRODUCTION

Throughout this note, our base theory is **ZF** unless otherwise specified. In Usuba [3], we introduced the notion of *Löwenheim-Skolem cardinal*, which corresponds to the downward Löwenheim-Skolem theorem in the context of **ZFC**.

Definition 1.1. Let κ be an uncountable cardinal.

- (1) κ is *weakly Löwenheim-Skolem* (weakly LS, for short) if for every $\gamma < \kappa$, $\alpha \geq \kappa$, and $x \in V_\alpha$, there is $X \prec V_\alpha$ such that $V_\gamma \subseteq X$, $x \in X$, and the transitive collapse of X belongs to V_κ .
- (2) κ is *Löwenheim-Skolem* (LS, for short) if for every $\gamma < \kappa$, $\alpha \geq \kappa$, and $x \in V_\alpha$, there is $\beta \geq \alpha$ and $X \prec V_\beta$ such that $V_\gamma \subseteq X$, $x \in X$, $V_\gamma(X \cap V_\alpha) \subseteq X$, and the transitive collapse of X belongs to V_κ .

Note that if κ is a limit of LS (weakly LS, respectively) cardinals, then κ is LS (weakly LS, respectively). In [3], we proved that some non-trivial consequences of the Axiom of Choice **AC** holds on the successor of a singular weakly LS cardinal:

Theorem 1.2. *Suppose κ is a singular weakly LS cardinal (e.g., a singular limit of weakly LS cardinals).*

- (1) *There is no cofinal map from V_κ into κ^+ , hence κ^+ is regular.*
- (2) *For every function f from V_κ into the club filter over κ^+ , the intersection $\bigcap f \restriction V_\kappa$ contains a club in κ^+ . Hence the club filter is κ^+ -complete.*
- (3) *For every regressive function $f : \kappa^+ \rightarrow \kappa^+$, there is $\alpha < \kappa^+$ such that the set $\{\eta < \kappa^+ \mid f(\eta) = \alpha\}$ is stationary.*

Date: April 30, 2020.

2010 Mathematics Subject Classification. Primary 03E10, 03E25.

Key words and phrases. Axiom of Choice, Löwenheim-Skolem cardinal.

See [3] for more information, and applications to choiceless set-theoretic geology.

In this note, we provide related results and other applications of LS cardinals, and we show that various consequences of AC follows from LS cardinals.

We will use the following notion in this note. For a set x , let $\|x\|$ be the least ordinal α such that there is a surjection from V_α onto x . Note that:

- $\|V_\alpha\| = \alpha$ for every infinite ordinal α .
- If $x \subseteq y$, or there is a surjection from y onto x , then $\|x\| \leq \|y\|$.
- For every ordinal α , there is a cardinal κ with $\|\kappa\| > \alpha$.
- If κ is weakly LS, then for every $\gamma < \kappa$ there is no surjection from V_γ onto κ ([3]), so $\|\kappa\| = \kappa$.
- Suppose $\kappa = \|\kappa\|$. Let $\alpha \geq \kappa$ and $X \prec V_\alpha$ be with $\|X\| < \kappa$. Then the transitive collapse of X is in V_κ ; Let \bar{X} be the transitive collapse of X , and δ the rank of \bar{X} . There is a surjection from \bar{X} onto δ , so $\|\delta\| \leq \|\bar{X}\|$. Since the collapsing map is a bijection from X onto \bar{X} , we have $\|\delta\| \leq \|\bar{X}\| = \|X\| < \kappa$. If $\delta \geq \kappa$, then we know $\|\delta\| \geq \|\kappa\| = \kappa$, but this is impossible and we obtain $\delta < \kappa$.

Using this notion, we can reformulate LS and weakly LS cardinals as follows:

- κ is weakly LS if and only if $\|\kappa\| = \kappa$, and for every $\alpha \geq \kappa$ and set $A \subseteq V_\alpha$ with $\|A\| < \kappa$, there is $X \prec V_\alpha$ such that $A \subseteq X$ and $\|X\| < \kappa$.
- κ is LS if and only if $\|\kappa\| = \kappa$, and for every $\gamma < \kappa$, $\alpha \geq \kappa$ and set $A \subseteq V_\alpha$ with $\|A\| < \kappa$, there is $\beta \geq \alpha$ and $X \prec V_\beta$ such that $A \subseteq X$, $\|X\| < \kappa$, and for every $y \subseteq X \cap V_\alpha$, if $\|y\| \leq \gamma$ then $y \in X$.

2. ON ELEMENTARY SUBMODELS

First let us present somewhat trivial characterization.

Proposition 2.1. *Let κ be an uncountable cardinal. Then the following are equivalent:*

- (1) κ is weakly LS.
- (2) $\|\kappa\| = \kappa$ and for every first order structure $\mathcal{M} = \langle M; \dots \rangle$ with countable language and $A \subseteq M$ with $\|A\| < \kappa$, there is an elementary submodel $\mathcal{N} = \langle N; \dots \rangle \prec \mathcal{M}$ such that $A \subseteq N$ and $\|N\| < \kappa$.

Proof. (2) \Rightarrow (1). Take $\gamma < \kappa$, $\alpha \geq \kappa$, and $x \in V_\alpha$. Identifying x as a constant, consider the structure $\langle V_\alpha; \in, x \rangle$. By (2), we can find an elementary submodel $X \prec V_\alpha$ such that $x \in X$, $V_\gamma \subseteq X$, and $\|X\| < \kappa$. If Y is the transitive collapse of X , then Y must be in V_κ .

(1) \Rightarrow (2). $\|\kappa\| = \kappa$ is already noted above. Fix a structure $\mathcal{M} = \langle M; \dots \rangle$ and $A \subseteq M$ such that there is a surjection f from some $x \in V_\kappa$ onto A .

Take a large $\alpha > \kappa$ with $f, \mathcal{M} \in V_\alpha$, and take also a large $\gamma < \kappa$ with $x \in V_\gamma$. By (1), we can find $X \prec V_\alpha$ such that $f, \mathcal{M} \in V_\alpha$, $V_\gamma \subseteq X$, and the transitive collapse of X is in V_κ . Since $x \in V_\gamma \subseteq X$, we have $A = f^{\omega}x \subseteq X$. Let $\mathcal{N} = \langle M \cap X; \dots \rangle$. Since $A \subseteq X \prec V_\alpha$, it is easy to see that $\mathcal{N} \prec \mathcal{M}$ and $A \subseteq M \cap X$. If $\pi : X \rightarrow Y$ is the collapsing map, then $\pi \upharpoonright M \cap X : M \cap X \rightarrow \pi(M)$ is a bijection and $\pi(M) \in V_\kappa$. Then the inverse map gives a surjection from $\pi(M)$ onto $M \cap X$. Hence $\|M \cap X\| < \kappa$. \square

3. ON SUCCESSORS OF WEAKLY LS CARDINALS

In ZFC, for every cardinal κ and $\alpha \geq \kappa^+$, we can find an elementary submodel $X \prec V_\alpha$ with $|X| = \kappa \subseteq X$ and $X \cap \kappa^+ \in \kappa^+$. We can prove a similar result for singular weakly LS cardinals.

Lemma 3.1. *Let κ be a singular weakly LS cardinal. Then for every $\alpha \geq \kappa^+$ and $x \in V_\alpha$, there is $X \prec V_\alpha$ such that $x \in X$, $V_\kappa \subseteq X$, $X \cap \kappa^+ \in \kappa^+$, and $\|X\| = \kappa$.*

Proof. Fix a large $\beta > \alpha$, and take $Y \prec V_\beta$ such that $\alpha, \kappa, x \in Y$, $\text{cf}(\kappa) \subseteq Y$, and the transitive collapse of Y belongs to V_κ .

Let $\mathcal{F} = \{Z \prec V_\alpha \mid x \in Z \text{ and the transitive collapse of } Z \text{ is in } V_\kappa\}$. We have $\mathcal{F} \in Y$.

Claim 3.2. (1) $\mathcal{F} \cap Y$ is upward directed, that is, for every $Z_0, Z_1 \in \mathcal{F} \cap Y$, there is $Z_2 \in \mathcal{F} \cap Y$ with $Z_0 \cup Z_1 \subseteq Z_2$.

(2) For every $\gamma < \kappa$, there is $Z \in \mathcal{F} \cap Y$ with $V_\gamma \subseteq Z$.

Proof of Claim. (1) Take a large $\gamma < \kappa$ such that the transitive collapses of Z_0 and Z_1 are in V_γ , and take surjections $f_0 : V_\gamma \rightarrow Z_0$ and $f_1 : V_\gamma \rightarrow Z_1$. Since $Z_0, Z_1 \in Y \prec V_\beta$, we may assume that $\gamma, f_0, f_1 \in Y$. In Y , we can choose $Z'_2 \prec V_{\alpha+\omega}$ such that $V_\gamma \subseteq Z'_2$, $x, \alpha, f_0, f_1 \in Z'_2$, and the transitive collapse of Z'_2 is in V_κ . Let $Z_2 = Z'_2 \cap V_\alpha \in Y$. We know $Z_2 \prec V_\alpha$, so $Z_2 \in \mathcal{F} \cap Y$. Since $V_\gamma \subseteq Z_2$, we also have $Z_0 \cup Z_1 \subseteq Z_2$.

(2) Since $\text{cf}(\kappa) \subseteq Y$, we have that $Y \cap \kappa$ is cofinal in κ . Hence for a given $\gamma < \kappa$, there is $\delta \in Y \cap \kappa$ with $\gamma \leq \delta$. Then in Y we can choose $Z \in \mathcal{F}$ with $V_\delta \subseteq Z$. \square

Let $X = \bigcup(\mathcal{F} \cap Y)$. By the claim above, we have that $x \in X \prec V_\alpha$ and $V_\kappa \subseteq X$.

Claim 3.3. *There is a surjection from V_κ onto X .*

Proof of Claim. Let \bar{Y} be the transitive collapse of Y , and $\pi : Y \rightarrow \bar{Y}$ the collapsing map. For each $Z \in \mathcal{F} \cap Y$, let \bar{Z} be the transitive collapse of Z , and π_z be the collapsing map.

Define $f : V_\kappa \times V_\kappa \rightarrow X$ as follows: For $\langle a, b \rangle \in V_\kappa \times V_\kappa$, if $\pi^{-1}(a) \in \mathcal{F} \cap Y$ and $b \in \overline{Z}$ (where $Z = \pi^{-1}(a)$), then $f(a, b) = \pi_Z^{-1}(b)$. Otherwise, $f(a, b) = \emptyset$. This f is an surjection; Take $c \in X$. Then $c \in Z$ for some $Z \in \mathcal{F} \cap Y$, and $f(\pi(Z), \pi_Z(c)) = c$. We can easily take a surjection from V_κ onto $V_\kappa \times V_\kappa$, hence we obtain a surjection from V_κ onto X . \square

Finally, since there is no cofinal map from V_κ into κ^+ by Theorem 1.2, we have that $\sup(X \cap \kappa^+) < \kappa^+$. In addition, since $\kappa \subseteq X$, we have $\sup(X \cap \kappa^+) \subseteq X$ and $\sup(X \cap \kappa^+) = X \cap \kappa^+ \in \kappa^+$. \square

Let κ be a singular weakly LS cardinal. While we already knew that the club filter over κ^+ is κ^+ -complete, we do not know if it is normal. Among this, we can construct a normal filter over κ^+ which is definable with parameter κ^+ .

Note 3.4. Let κ be a cardinal, and F a filter over κ . Then the following are equivalent:

- (1) For every $X_\alpha \in F$ ($\alpha < \kappa$), the diagonal intersection $\Delta_{\alpha < \kappa} X_\alpha = \{\beta < \kappa \mid \beta \in X_\alpha \text{ for all } \alpha < \beta\}$ is in F .
- (2) For every $X \in F^+$ and regressive function $f : X \rightarrow \kappa$, there is $\alpha < \kappa$ with $\{\beta \in X \mid f(\beta) = \alpha\} \in F^+$.

Where $F^+ = \{X \in \mathcal{P}(\kappa) \mid X \cap C \neq \emptyset \text{ for every } C \in F\}$. An element of F^+ is an F -positive set. We say that a filter F is *normal* if F is proper, contains all co-bounded subsets of κ , and satisfies the above conditions (1) and/or (2).

Proposition 3.5. Let κ be a singular weakly LS cardinal. Let $F \subseteq \mathcal{P}(\kappa^+)$ be the set such that: $D \in F \iff$ there is $\alpha \geq \kappa^+$ and $x \in V_\alpha$ such that D contains the set $\{\eta < \kappa^+ \mid \text{there is } X \prec V_\alpha \text{ with } x \in X \text{ and } \eta = X \cap \kappa^+\}$. Then F is a normal filter over κ^+ .

Proof. By Lemma 3.1, we have that $\emptyset \notin F$ and $\kappa^+ \setminus \eta \in F$ for every $\eta < \kappa^+$. One can check that F is a filter over κ^+ . For the normality, take a family $\{D_\gamma \in F \mid \gamma < \kappa^+\}$, and let $D = \Delta_{\gamma < \kappa^+} D_\gamma$. Suppose to the contrary that $D \notin F$. Now fix a large $\beta > \kappa^+$ such that for every $\gamma < \kappa^+$, there is $\alpha < \beta$ and $x \in V_\alpha$ such that $\{\eta < \kappa^+ \mid \text{there is } X \prec V_\alpha \text{ with } x \in X \text{ and } \eta = X \cap \kappa^+\} \subseteq D_\gamma$. Since $D \notin F$, we can find $Y \prec V_\beta$ such that Y contains all relevant objects, $Y \cap \kappa^+ \in \kappa^+$, and $Y \cap \kappa^+ \notin D$. Let $\delta = Y \cap \kappa^+$. We see that $\delta \in D_\gamma$ for every $\gamma < \delta$, this is a contradiction.

Take $\gamma < \delta$. Then $\gamma \in Y$. Hence we can find $\alpha \in Y$ and $x \in Y \cap V_\alpha$ such that $\{\eta < \kappa^+ \mid \text{there is } X \prec V_\alpha \text{ with } x \in X \text{ and } \eta = X \cap \kappa^+\} \subseteq D_\gamma$. However, since $\alpha, x \in Y$, we have $x \in Y \cap V_\alpha \prec V_\alpha$ and $(Y \cap V_\alpha) \cap \kappa^+ = \delta$, hence $\delta \in D_\gamma$. \square

For a non-empty set S , let $\text{Col}(S)$ be the poset of all finite partial functions from ω to S with the reverse inclusion order. The forcing with $\text{Col}(S)$ adds a surjection from ω onto S .

Proposition 3.6. *Let κ be a singular weakly LS cardinal. Then $\text{Col}(V_\kappa)$ forces $(\kappa^+)^V = \omega_1$ and the Dependent Choice DC.*

Proof. For the equality $(\kappa^+)^V = \omega_1$, take $p \in \text{Col}(V_\kappa)$ and a name \dot{f} for a function from κ to κ^+ . Define $F : \text{Col}(V_\kappa) \times \kappa \rightarrow \kappa^+$ as follows: For $\langle q, \alpha \rangle \in \text{Col}(V_\kappa) \times \kappa$, if $q \leq p$ and $q \Vdash \dot{f}(\alpha) = \eta$ for $\eta < \kappa^+$, then $F(q, \alpha) = \eta$. Otherwise, let $F(q, \alpha) = 0$. By Theorem 1.2, F is not a cofinal map, hence $\gamma = \sup(F''(\text{Col}(V_\kappa) \times \kappa)) < \kappa^+$. Then it is clear that $p \Vdash \dot{f}''\kappa \subseteq \gamma$, so $p \Vdash (\kappa^+)^V$ is regular.

To show that $\text{Col}(V_\kappa)$ forces DC, take $p \in \text{Col}(V_\kappa)$, $\text{Col}(V_\kappa)$ -names \dot{S} and \dot{R} such that $p \Vdash \dot{S}$ is non-empty, $\dot{R} \subseteq \dot{S}^2$, and for every $x \in \dot{S}$ there is $y \in \dot{S}$ with $\langle x, y \rangle \in \dot{R}$.

Fix a large limit $\alpha > \kappa^+$ with $\dot{S}, \dot{R} \in V_\alpha$. By Lemma 3.1, we can find $X \prec V_\alpha$ such that $V_\kappa \subseteq X$, X contains all relevant objects, $X \cap \kappa^+ \in \kappa^+$, and $\|X\| = \kappa$. Note that $\text{Col}(V_\kappa) \subseteq X$.

Take a $(V, \text{Col}(V_\kappa))$ -generic G , and let $X[G] = \{\dot{x}_G \mid \dot{x} \in X \text{ is a } \text{Col}(V_\kappa)\text{-name}\}$ (where \dot{x}_G is the interpretation of \dot{x} by G). Since $\text{Col}(V_\kappa) \subseteq X$, we may assume $X[G] \prec V_\alpha[G] = V[G]_\alpha$. There is a canonical surjection from X onto $X[G]$, namely $\dot{x} \mapsto \dot{x}_G$, hence we can take a surjection from V_κ onto $X[G]$. Since V_κ is countable in $V[G]$, we have that $X[G]$ is countable as well. Let $S = \dot{S}_G$ and $R = \dot{R}_G$. We know $S, R \in X[G]$. By the elementarity of $X[G]$, for every $x \in S \cap X[G]$, there is $y \in S \cap X[G]$ with $\langle x, y \rangle \in R$. Now $X[G]$ is well-orderable, thus we can take a map $f : \omega \rightarrow S \cap X[G]$ such that $\langle f(n), f(n+1) \rangle \in R$ for every $n < \omega$. \square

When κ is regular weakly LS, we can obtain a similar result to Lemma 3.1. Note that in ZFC, κ is regular weakly LS if and only if κ is inaccessible.

Lemma 3.7. *Let κ be a regular weakly LS cardinal. Then for every $\gamma < \kappa$, $\alpha > \kappa$, and $x \in V_\alpha$, there is $X \prec V_\alpha$ such that $x \in X$, $\gamma < X \cap \kappa \in \kappa$, $V_{X \cap \kappa} \subseteq X$, and the transitive collapse of X is in V_κ .*

Proof. Similar to Lemma 3.1. Take a large $\beta > \alpha$ and $Y \prec X_\beta$ such that Y contains all relevant objects, and the transitive collapse of Y is in V_κ . We know $\sup(Y \cap \kappa) < \kappa$; If $\sup(Y \cap \kappa) = \kappa$, then $\text{ot}(Y \cap \kappa) = \kappa$ since κ is regular. However then the transitive collapse of Y cannot be in V_κ .

Let $\mathcal{F} = \{Z \prec V_\alpha \mid x \in Z, \text{ the transitive collapse of } Z \text{ is in } V_\kappa\}$, and $X = \bigcup(\mathcal{F} \cap Y)$. We have $X \prec V_\alpha$. For every $Z \in \mathcal{F} \cap Y$, we have that $Z \cap \kappa \leq \sup(Y \cap \kappa)$, and for every $\delta \in Y \cap \kappa$, there is $Z \in \mathcal{F} \cap Y$ with $V_\delta \subseteq Z$. Hence $X \cap \kappa = \sup(Y \cap \kappa)$ and $V_{X \cap \kappa} \subseteq X$.

Take a large $\delta < \kappa$ such that the transitive collapse of Y is in V_δ . Note that $\sup(Y \cap \kappa) \leq \delta$. For every $Z \in \mathcal{F} \cap Y$, the transitive collapse of Z is also in V_δ . Then as in Lemma 3.1 we can define a surjection from V_δ onto X . Therefore $\|X\| \leq \delta < \kappa$, and the transitive collapse of X is in V_κ . \square

4. ON SUCCESSORS OF LS CARDINALS

If κ is a singular LS cardinal and there is a weakly LS cardinal $\leq \text{cf}(\kappa)$, we can take a small elementary submodel with certain closure property. We use the following fact:

Theorem 4.1 ([3]). *Let κ be a weakly LS cardinal. Then for every cardinal $\lambda \geq \kappa$ and $x \in V_\kappa$, there is no cofinal map from x into λ^+ .*

Lemma 4.2. *Let α be a limit ordinal α and a set x , if $\text{cf}(\alpha) > \|x\|$ and there is a weakly LS cardinal κ with $\text{cf}(\alpha) \geq \kappa \geq \|x\|$, then there is no cofinal map from x into α .*

Proof. First suppose $\|x\| < \kappa < \text{cf}(\alpha)$. In this case we may assume $x \in V_\kappa$. If $\text{cf}(\alpha)$ is a successor cardinal, then we have done by Theorem 4.1. Suppose $\text{cf}(\alpha)$ is a limit cardinal. Then $\text{cf}(\alpha) > \kappa^+$, so $\{\eta < \alpha \mid \text{cf}(\eta) = \text{cf}(\kappa^+)\}$ is stationary in α . If $f : x \rightarrow \alpha$ is a cofinal map, then $C = \{\eta < \alpha \mid \eta \cap f''x \text{ is cofinal in } \eta\}$ is a club in α . Hence we can find $\eta < \alpha$ such that $\eta \cap f''x$ is cofinal in η and $\text{cf}(\eta) = \text{cf}(\kappa^+)$. However then there is a cofinal map from x into κ^+ , This is a contradiction.

Next suppose $\|x\| = \kappa < \text{cf}(\alpha)$. Then we may assume $x = V_\kappa$. Take $f : V_\kappa \rightarrow \alpha$. By the previous case, for every $\gamma < \kappa$, we have $\sup(f''V_\gamma) < \alpha$. Since $\text{cf}(\alpha) > \kappa$, we have $\sup(f''V_\kappa) = \sup_{\gamma < \kappa} \sup(f''V_\gamma) < \alpha$.

Finally suppose $\|x\| < \kappa = \text{cf}(\alpha)$. Then κ is a regular weakly LS cardinal. If there is a cofinal map from x into α , we can find a cofinal map f from x into κ . Take an elementary submodel $X \prec V_{\alpha+\omega}$ such that $V_{\|x\|} \subseteq X$, X contains all relevant objects, and the transitive collapse of X is in V_κ . We know $f''x \subseteq X$, hence $\sup(X \cap \kappa) = \kappa$. Then $\text{ot}(X \cap \kappa) = \kappa$ because κ is regular, so the transitive collapse of X cannot be in V_κ . This is a contradiction. \square

Lemma 4.3. *Let κ be a singular LS cardinal and $\nu \leq \text{cf}(\kappa)$ a weakly LS cardinal. Then for every $\alpha > \kappa$ and $x \in V_\alpha$, there is $\beta > \alpha$ and $X \prec V_\beta$ such that $V_\kappa \subseteq X$, $x \in X$, $X \cap \kappa^+ \in \kappa^+$, $V_\gamma(X \cap V_\alpha) \subseteq X$ for every $\gamma < \nu$, and $\|X\| = \kappa$.*

Proof. Take a large β and $Y \prec V_\beta$ such that $\alpha, x, \dots \in Y$, $V_\nu(Y \cap V_{\alpha+\omega+\omega}) \subseteq Y$, $\text{cf}(\kappa) \subseteq Y$, and the transitive collapse of Y is in V_κ . Let $\mathcal{F} = \{Z \prec V_{\alpha+\omega} \mid x \in Z, \text{ the transitive collapse of } Z \text{ is in } V_\kappa\}$. Let $X = \bigcup(\mathcal{F} \cap Y)$. As in the

proof of Lemma 3.1, we have $X \prec V_{\alpha+\omega}$, $x \in X$, $V_\kappa \subseteq X$, $X \cap \kappa^+ \in \kappa^+$, and $\|X\| = \kappa$.

We have to see that $V_\gamma(X \cap V_\alpha) \subseteq X$ for every $\gamma < \nu$. Take $f : V_\gamma \rightarrow X \cap V_\alpha$. Since ν is weakly LS, we can find $Y' \prec V_\beta$ such that Y contains all relevant objects, $V_\gamma \subseteq Y'$, and the transitive collapse of Y' is in V_ν . We have $f \subseteq Y'$. Let $\mathcal{F}' = \mathcal{F} \cap Y \cap Y' \subseteq Y \cap V_{\alpha+\omega+\omega}$. Since $V_\nu(Y \cap V_{\alpha+\omega+\omega}) \subseteq Y$ and $\|\mathcal{F}'\| \leq \|Y\| < \nu$, we have $\mathcal{F}' \in Y$. Note that $\|Z\| < \kappa$ for every $Z \in \mathcal{F}'$. Since $\|\mathcal{F}'\| < \nu \leq \text{cf}(\kappa)$, there is no cofinal map from \mathcal{F}' into κ by Lemma 4.2. Thus we have $\delta = \sup\{\|Z\| \mid Z \in \mathcal{F}'\} < \kappa$. Then, as in the proof of Lemma 3.1, we can define a surjection from V_δ onto $\bigcup \mathcal{F}'$. In Y , we can find $Z' \in \mathcal{F} \cap Y$ such that $\mathcal{P}(V_\alpha \cap \bigcup \mathcal{F}') \subseteq Z'$. Since $f \in Y'$, for each $a \in V_\gamma$ there is $Z \in (\mathcal{F} \cap Y) \cap Y'$ with $f(a) \in Z$. Hence $f''V_\gamma \subseteq V_\alpha \cap \bigcup \mathcal{F}'$, and $f''V_\gamma \in Z' \subseteq X$. \square

Note 4.4. In Lemma 4.5, if $\nu < \text{cf}(\kappa)$ is a singular weakly LS cardinal, then we can require that $V_\nu(X \cap V_\alpha) \subseteq X$.

If κ is regular LS cardinal, we have the following parallel result:

Lemma 4.5. *Let κ be an LS cardinal and $\nu < \kappa$ a weakly LS cardinal. Then for every $\alpha > \kappa$, $x \in V_\alpha$, and $\gamma < \kappa$, there is $\beta \geq \alpha$ and $X \prec V_\beta$ such that $x \in X$, $\gamma < X \cap \kappa \in \kappa$, $V_{X \cap \kappa} \subseteq X$, $V_\delta(X \cap V_\alpha) \subseteq X$ for every $\delta < \nu$, and the transitive collapse of X is in V_κ .*

For an ordinal γ , let DC_γ be the assertion that for every non-empty set S and $G : {}^{<\gamma}S \rightarrow \mathcal{P}(S) \setminus \{\emptyset\}$, there is $f : \gamma \rightarrow S$ such that $f(\alpha) \in G(f \upharpoonright \alpha)$ for every $\alpha < \gamma$. Let $\text{DC}_{<\gamma}$ be the assertion that DC_δ holds for every $\delta < \gamma$.

DC_ω is the Dependent Choice, and it is known that AC is equivalent to that DC_γ for every γ . Note also that the following:

- (1) $\text{DC}_\gamma \Rightarrow \text{DC}_{\gamma+1}$.
- (2) If γ is a singular ordinal and $\text{DC}_{<\gamma}$ holds, then DC_γ holds as well.
- (3) $\text{DC}_\gamma \Rightarrow \text{DC}_{<\gamma^+}$.

Woodin [4] proved that collapsing a supercompact cardinal yields DC_κ for some large κ . Where κ is *supercompact* if for every $\alpha > \kappa$, there is $\beta \geq \alpha$, a transitive set N with $V_\alpha N \subseteq N$, and an elementary embedding $j : V_\beta \rightarrow N$ with critical point κ and $\alpha < j(\kappa)$. It is known that a supercompact cardinal is a limit of LS cardinals ([3]).

We prove a similar result using LS cardinals.

Definition 4.6. Let κ be a regular cardinal and S a non-empty set. Let $\text{Col}(\kappa, S)$ be the poset of all partial functions p from κ to S with $|p| < \kappa$ (so p is assumed to be well-orderable). The ordering of $\text{Col}(\kappa, S)$ is the reverse inclusion.

If G is $(V, \text{Col}(\kappa, S))$ -generic, then $\bigcup G$ is a surjection from κ onto S .

Proposition 4.7. *Let κ be a regular uncountable cardinal, and suppose $\text{DC}_{<\kappa}$ holds. Let $\lambda > \kappa$ be a singular LS cardinal such that $\text{cf}(\lambda) > \kappa$ and there is a weakly LS cardinal ν with $\kappa \leq \nu \leq \text{cf}(\lambda)$. Then $\text{Col}(\kappa, V_\lambda)$ does not add new $< \kappa$ -sequences, and forces $(\lambda^+)^V = (\kappa^+)^{V^{\text{Col}(\kappa, V_\lambda)}}$ and DC_κ .*

Proof. By $\text{DC}_{<\kappa}$, we know that $\text{Col}(\kappa, V_\lambda)$ is κ -closed and does not add new $< \kappa$ -sequences. We can check that $\text{Col}(\kappa, V_\lambda)$ preserves $\text{DC}_{<\kappa}$ as well.

To see $\text{Col}(\kappa, V_\lambda)$ forces $(\lambda^+)^V = (\kappa^+)^{V^{\text{Col}(\kappa, V_\lambda)}}$, it is clear that $(\lambda^+)^V \leq (\kappa^+)^{V^{\text{Col}(\kappa, V_\lambda)}}$. For the converse, take $p \in \text{Col}(\kappa, V_\lambda)$ and a name \dot{f} for a function from λ to $(\lambda^+)^V$. Define $F : \lambda \times \text{Col}(\kappa, V_\lambda) \rightarrow \lambda^+$ by $F(\delta, q) = \eta \iff q \leq p$ and $q \Vdash \dot{f}(\delta) = \eta$. F is not cofinal by Theorem 1.2, hence the image of F has an upper bound in λ^+ , say η . It is clear that $p \Vdash \dot{f} \text{``}\lambda \subseteq \eta\text{''}$, so $\Vdash \text{``}(\lambda^+)^V \text{ is regular''}$.

For DC_κ , take $p \in \text{Col}(\kappa, V_\lambda)$ and names \dot{S} and \dot{F} with $p \Vdash \text{``}\dot{F} : <^\kappa \dot{S} \rightarrow \mathcal{P}(\dot{S}) \setminus \{\emptyset\}\text{''}$. Fix a sufficiently large $\alpha > \lambda^+$. By Lemma 4.5, we can find $\beta > \alpha$ and $X \prec V_\beta$ such that X contains all relevant objects, $V_\lambda \subseteq X$, $X \cap \lambda^+ \in \lambda^+$, $V_\gamma(X \cap V_{\alpha+\omega}) \subseteq X$ for every $\gamma < \nu$, and $\|X\| = \lambda$. Take a $(V, \text{Col}(\kappa, V_\lambda))$ -generic G and work in $V[G]$. We may assume $X[G] \prec V[G]_\beta$. We see that $X[G] \cap V[G]_\alpha$ is closed under $< \kappa$ -sequences in $V[G]$. Take $\eta < \kappa$ and $f : \eta \rightarrow X[G] \cap V[G]_\alpha$. By $\text{DC}_{<\kappa}$ in $V[G]$, we can find a sequence of $\text{Col}(\kappa, V_\lambda)$ -names $\langle \dot{x}_i \mid i < \eta \rangle$ such that $\dot{x}_i \in X$ and $f(i) = (\dot{x}_i)_G$ for every $i < \eta$. We may assume $\dot{x}_i \in V_{\alpha+\omega}$ for $i < \eta$. Since $\text{Col}(\kappa, V_\lambda)$ does not add new $< \kappa$ -sequences, we have $\langle \dot{x}_i \mid i < \eta \rangle \in V$, hence $\langle \dot{x}_i \mid i < \eta \rangle \in X$ and $\{(\dot{x}_i)_G \mid i < \eta\} \in X[G]$.

Since there is a surjection from κ onto V_λ , from V_λ onto X , and from X onto $X[G]$, we can obtain a bijection from κ onto $X[G]$. Let $S = \dot{S}_G$ and $F = \dot{F}_G$. Since $X[G] \prec V[G]_\beta$, $X[G] \cap V[G]_\alpha$ is closed under $< \kappa$ -sequences, and $X[G]$ is well-orderable, we can take $f : \kappa \rightarrow S \cap X[G]$ such that $f(i) \in F(f \upharpoonright i)$ for every $i < \kappa$. \square

If there are proper class many LS cardinals, then for every cardinal κ , there is a singular LS cardinal $\lambda > \kappa$ such that there is an LS cardinal ν with $\kappa < \nu < \text{cf}(\lambda)$; Take a singular LS cardinal $\nu > \kappa$, and let λ be the ν^+ -th LS cardinal above ν . The cofinality of λ is ν^+ . In addition, in [3] we proved that if κ is a limit of LS cardinals, then every poset with rank $< \kappa$ forces that κ is LS. Using these facts, we can obtain the following corollary:

Corollary 4.8. *Suppose there are proper class many LS cardinals. Then there is a definable class forcing which forces ZFC.*

5. LS CARDINALS IN $\text{HOD}(V_\lambda)$

For a set X , let $\text{HOD}(X)$ be the class of all hereditarily definable sets with parameters from $\text{ON} \cup \text{trcl}(\{X\})$. $\text{HOD}(X)$ is a transitive model of ZF with $X \in \text{HOD}(X)$. By the definition of $\text{HOD}(X)$, for every $x \in \text{HOD}(X)$ there is an ordinal θ and a surjection $\sigma : \theta \times {}^{<\omega}(\text{trcl}(\{X\})) \rightarrow x$ with $\sigma \in \text{HOD}(X)$. If $X = V_\alpha$ and α is limit, then V_α is transitive, ordinal definable, and ${}^{<\omega}V_\alpha \subseteq V_\alpha$. Hence the domain of σ can be the set $\theta \times V_\alpha$ in this case.

Lemma 5.1. *Let δ be a limit ordinal. Then for every limit ordinals $\alpha > \beta \geq \delta$ and $x \in V_\alpha \cap \text{HOD}(V_\delta)$, there is $X \prec V_\alpha \cap \text{HOD}(V_\delta)$ such that $x \in X \in \text{HOD}(V_\delta)$, $V_\beta \cap \text{HOD}(V_\delta) \subseteq X$, and $\|X\|^{\text{HOD}(V_\delta)} = \beta$.*

Proof. For given $\alpha > \beta \geq \delta$, we can take a large ordinal θ and a surjection $\sigma : \theta \times V_\delta \rightarrow V_\alpha \cap \text{HOD}(V_\delta)$ with $\sigma \in \text{HOD}(V_\delta)$. By induction on $n < \omega$, we define X_n and f_n as follows. First, let $X_0 = (V_\beta \cap \text{HOD}(V_\delta)) \cup \{x\}$, and take a surjection $f_0 : V_\beta \cap \text{HOD}(V_\delta) \rightarrow X_0$ with $f_0 \in \text{HOD}(V_\delta)$. Suppose X_n and f_n are defined and f_n is a surjection from $V_\beta \cap \text{HOD}(V_\delta)$ onto X_n . For a formula $\varphi(v_1, \dots, v_k, w)$ and $x_1, \dots, x_k \in X_n$, if $\exists w \varphi(x_1, \dots, x_k, w)$ holds in $V_\alpha \cap \text{HOD}(V_\delta)$ then we can find the least $\alpha_{\varphi, x_1, \dots, x_k} \in \theta$ such that $\exists w \in \sigma^{-1}(\{\alpha_{\varphi, x_1, \dots, x_k}\} \times V_\delta) \varphi(x_1, \dots, x_k, w)$ holds in $V_\alpha \cap \text{HOD}(V_\delta)$. Set $X_{n+1} = X_n \cup \bigcup \{\sigma^{-1}(\{\alpha_{\varphi, x_1, \dots, x_k}\} \times V_\delta) : \varphi \text{ is a formula, } x_1, \dots, x_k \in X_n\}$. There is a canonical surjection from $V_\beta \cap \text{HOD}(V_\delta)$ onto $\bigcup \{\sigma^{-1}(\{\alpha_{\varphi, x_1, \dots, x_k}\} \times V_\delta) : \varphi \text{ is a formula, } x_1, \dots, x_k \in X_n\}$, namely $\langle [\varphi], y_1, \dots, y_k, z \rangle \mapsto \sigma(\alpha_{\varphi, f_n(y_1), \dots, f_n(y_k)}, z)$ (where $[\varphi] \in \omega$ is the Gödel number of φ). Hence we can define $f_{n+1} : V_\beta \cap \text{HOD}(V_\delta) \rightarrow X_{n+1}$ canonically using this surjection and f_n . We can carry out this construction in $\text{HOD}(V_\delta)$, hence $\langle X_n, f_n \mid n < \omega \rangle \in \text{HOD}(V_\delta)$. Thus we have $X = \bigcup_n X_n \in \text{HOD}(V_\delta)$. We know $V_\beta \cap \text{HOD}(V_\delta) \subseteq X$ and $x \in X$. By Tarski-Vaught criterion, we have $X \prec V_\alpha \cap \text{HOD}(V_\delta)$. Moreover, there is a surjection from $\omega \times (V_\beta \cap \text{HOD}(V_\delta))$ onto X in $\text{HOD}(V_\delta)$ constructed from $\langle f_n \mid n < \omega \rangle$, so we can construct a surjection $f : V_\beta \cap \text{HOD}(V_\delta) \rightarrow X$ in $\text{HOD}(V_\delta)$, hence $\|X\|^{\text{HOD}(V_\delta)} = \beta$. \square

The following is immediate from the previous lemma:

Corollary 5.2. *Let δ be a limit ordinal. Then for every cardinal $\kappa > \delta$ with $\|\kappa\| = \kappa$, κ is weakly LS in $\text{HOD}(V_\delta)$.*

Proof. Take $\gamma < \kappa$, $\alpha > \kappa$ and $x \in V_\alpha \cap \text{HOD}(V_\delta)$. We may assume α and γ are limit ordinals and $\gamma \geq \delta$. By the previous lemma, there is $X \prec V_\alpha \cap \text{HOD}(V_\delta)$ such that $x \in X \in \text{HOD}(V_\delta)$, $V_\gamma \cap \text{HOD}(V_\delta) \subseteq X$,

and $\|X\|^{\text{HOD}(V_\delta)} = \gamma$. Then $\|X\| \leq \|X\|^{\text{HOD}(V_\delta)} = \gamma < \kappa$, so the transitive collapse of X must be in V_κ . \square

An uncountable cardinal κ is said to be *inaccessible* if for every $\gamma < \kappa$, there is no cofinal map from V_γ into κ . Every inaccessible cardinal is regular, and in ZFC, this definition is equivalent to the standard one. One can check that every regular weakly LS cardinal is inaccessible.

The following corollary is already proved in Schlutzenberg [2].

Corollary 5.3. *Let κ be an uncountable cardinal, and suppose κ is inaccessible in $\text{HOD}(V_\kappa)$. Then κ is a weakly LS cardinal in $\text{HOD}(V_\kappa)$. In particular, if κ is inaccessible then κ is regular weakly LS in $\text{HOD}(V_\kappa)$.*

Proof. Take $\gamma < \kappa$, $\alpha > \kappa$ and $x \in V_\alpha \cap \text{HOD}(V_\kappa)$. By Lemma 5.1, we can find $X \prec V_\alpha \cap \text{HOD}(V_\kappa)$ such that $x \in X \in \text{HOD}(V_\kappa)$, $V_\kappa \subseteq X$, and $\|X\|^{\text{HOD}(V_\kappa)} = \kappa$. Take a surjection $\sigma : V_\kappa \rightarrow X$ in $\text{HOD}(V_\kappa)$. Since κ is inaccessible in $\text{HOD}(V_\kappa)$, the set $\{\eta < \kappa \mid \sigma \upharpoonright V_\eta \prec V_\alpha \cap \text{HOD}(V_\kappa)\}$ contains a club in κ , hence we can find $\eta < \kappa$ such that $V_\eta \subseteq \sigma \upharpoonright V_\eta \prec V_\alpha \cap \text{HOD}(V_\kappa)$. \square

Note 5.4. For an uncountable cardinal κ , if κ is not regular in $\text{HOD}(V_\kappa)$ then $\text{cf}(\kappa)^{\text{HOD}(V_\kappa)} = \text{cf}(\kappa)$.

Corollary 5.5. *Let δ be a limit ordinal and $\nu \leq \text{cf}(\delta)$ a weakly LS cardinal. Then ν is weakly LS in $\text{HOD}(V_\delta)$.*

Proof. For $\gamma < \nu$, $\alpha > \delta$ and $x \in V_\alpha \cap \text{HOD}(V_\delta)$, we can take $X \prec V_\alpha \cap \text{HOD}(V_\delta)$ such that $x \in X \in \text{HOD}(V_\delta)$, $V_\delta \subseteq X$, and there is a surjection $\sigma : V_\delta \rightarrow X$ in $\text{HOD}(V_\delta)$. Define $R_0, R_1 \subseteq V_\delta^2$ by $a R_0 b \iff \sigma(a) \in \sigma(b)$, and $a R_1 b \iff \sigma(a) = \sigma(b)$. We know $R_0, R_1 \in \text{HOD}(V_\delta)$. Since ν is weakly LS, we can find $\langle Y; \in, R_0 \cap Y, R_1 \cap Y \rangle \prec \langle V_\delta; \in, R_0, R_1 \rangle$ such that $V_\gamma \subseteq Y$, $\|Y\| < \nu$, and there is $a \in Y$ with $\sigma(a) = x$. Since $\|Y\| < \nu \leq \text{cf}(\delta)$, there is no cofinal map from Y to δ . In particular $\sup\{\text{rank}(y) \mid y \in Y\} < \delta$, so $Y \in V_\delta$ and $Y \in \text{HOD}(V_\delta)$. Then one can check that $V_\gamma \subseteq \sigma \upharpoonright Y \prec V_\alpha \cap \text{HOD}(V_\delta)$, as required. \square

When $\|\kappa\| = \kappa$ and κ is singular in $\text{HOD}(V_\kappa)$, we do not know whether κ is weakly LS in $\text{HOD}(V_\kappa)$.

Note 5.6. $\text{HOD}(V_\delta)$ and $\text{HOD}(V_\kappa)$ in this section can be replaced by $L(V_\delta)$ and $L(V_\kappa)$.

6. ON ELEMENTARY EMBEDDINGS

Woodin [4] proved that if κ is a singular limit of supercompact cardinals and there is a set A of ordinals with $\kappa^+ = (\kappa^+)^{L[A]}$, then there is no non-trivial elementary embedding $j : V_{\kappa+2} \rightarrow V_{\kappa+2}$. We can prove the same

result using weakly LS cardinals. Note that whenever A is a set of ordinals, $\text{HOD}(A)$ is a transitive model of ZFC and $A \in \text{HOD}(A)$.

Proposition 6.1. *Suppose κ is a singular weakly LS cardinal. If there is a set A of ordinals with $\kappa^+ = (\kappa^+)^{\text{HOD}(A)}$, then there is no non-trivial elementary embedding $j : V_{\kappa+2} \rightarrow V_{\kappa+2}$.*

Proof. Suppose such an elementary $j : V_{\kappa+2} \rightarrow V_{\kappa+2}$ exists. Let F be the club filter over κ^+ restricted to the set $\{\alpha < \kappa^+ \mid \text{cf}(\alpha) = \omega\}$, it is κ^+ -complete. F is ordinal definable, hence $F \cap \text{HOD}(A) \in \text{HOD}(A)$, and $F \cap \text{HOD}(A)$ is a κ^+ -complete filter in $\text{HOD}(A)$. Since $\text{HOD}(A)$ computes κ^+ correctly and is a model of ZFC, we have that $F \cap \text{HOD}(A)$ is not κ^+ -saturated in $\text{HOD}(A)$, hence is not $\text{crit}(j)$ -saturated. Take pairwise disjoint F -positive sets $\{E_\alpha \mid \alpha < \text{crit}(j)\} \in \text{HOD}(A)$. Each E_α is a stationary set in κ^+ , and we may assume that $E_\alpha \subseteq \{\eta < \kappa^+ \mid \text{cf}(\eta) = \omega\}$. Hence we obtain $\text{crit}(j)$ -many pairwise disjoint stationary subsets of $\{\eta < \kappa^+ \mid \text{cf}(\eta) = \omega\}$. This partition can be coded in $V_{\kappa+2}$, and using this coded partition we can derive the contradiction (e.g., see Kanamori [1]). \square

The assumption in the previous proposition can be weakened as follows.

Proposition 6.2. *Suppose κ is a singular weakly LS cardinal. If there is a set A of ordinals such that κ^+ is not measurable in $\text{HOD}(A)$, then there is no non-trivial elementary embedding $j : V_{\kappa+2} \rightarrow V_{\kappa+2}$.*

Proof. First note that κ is strong limit in $\text{HOD}(A)$. Let F be the filter defined as in Proposition 6.1. By Tarski's theorem, if $F \cap \text{HOD}(A)$ is $\text{crit}(j)$ -saturated in $\text{HOD}(A)$, then κ^+ is measurable in $\text{HOD}(A)$, this contradicts to the assumption. Hence F cannot be $\text{crit}(j)$ -saturated in $\text{HOD}(A)$, and we can take a large stationary partition of $\{\eta < \kappa^+ \mid \text{cf}(\eta) = \omega\}$. The rest is the same to before. \square

Corollary 6.3. *Suppose there are proper class many weakly LS cardinals. If there is a non-trivial elementary embedding $j : V \rightarrow V$, then $\{\kappa \mid \kappa \text{ is singular and } \kappa^+ \text{ is measurable in } \text{HOD}\}$ forms a proper class.*

7. ON ULTRAPOWERS

A cardinal κ is said to be *critical* if there is a transitive class M and an elementary embedding $j : V \rightarrow M$ with critical point κ (Schlutzenberg [2]). While the definition of critical cardinal is a second-order statement, in ZFC it is equivalent to a first order statement: There is a κ -complete ultrafilter over κ , that is, κ is measurable. Recently Schlutzenberg [2] showed that under the existence of proper class many weakly LS cardinals, the definition of critical cardinal is equivalent to a certain first order sentence. Actually,

under the assumption, he showed that the ultrapower by certain extender sequence satisfies Los' theorem, so the critical cardinals can be formalized by the existence of suitable extenders. Moreover his argument can formalize strong cardinal as well. See [2] for the proof of the following proposition.

Proposition 7.1. *Suppose there are proper class many weakly LS cardinals. Then for every cardinal κ the following are equivalent:*

- (1) *For every $\alpha > \kappa$, there is a weakly LS cardinal $\lambda > \alpha$ of uncountable cofinality, a transitive set N , and an elementary embedding $j : V_\lambda \rightarrow N$ such that the critical point of j is κ , $\alpha < j(\kappa)$, and $V_\alpha \subseteq N$.*
- (2) *For every α , there is a transitive class M and an elementary embedding $j : V \rightarrow M$ such that the critical point of j is κ , $\alpha < j(\kappa)$, and $V_\alpha \subseteq M$.*
- (3) *For every α , there is a definable transitive class M and a definable elementary embedding $j : V \rightarrow M$ such that the critical point of j is κ , $\alpha < j(\kappa)$, and $V_\alpha \subseteq M$.*

Other large cardinals defined by extenders, such as Woodin cardinal, superstrong cardinal, and I_2 -embedding, can be formalized by a similar way.

Acknowledgments. This research was supported by JSPS KAKENHI Grant Nos. 18K03403 and 18K03404.

REFERENCES

- [1] A. Kanamori, *The Higher Infinite: Large Cardinals in Set Theory from Their Beginnings*. Springer-Verlag, 1994.
- [2] F. Schlutzenberg, *Reinhardt cardinals and non-definability*. Preprint. Available at <https://arxiv.org/abs/2002.01215>
- [3] T. Usuba, *Choiceless Löwenheim-Skolem property and uniform definability of grounds*. To appear in the Proceedings of the Symposium on Advances in Mathematical Logic 2018. Available at <https://arxiv.org/abs/1904.00895>
- [4] W. H. Woodin, *Suitable extender models I*. J. Math. Log. 10 (2010), no. 1-2, 101–339.

(T. Usuba) FACULTY OF SCIENCE AND ENGINEERING, WASEDA UNIVERSITY, OKUBO 3-4-1, SHINJYUKU, TOKYO, 169-8555 JAPAN
Email address: usuba@waseda.jp