

**Classification of generic random
holomorphic dynamical systems
associated with
analytic families of rational maps**

Hiroki Sumi

Graduate School of

Human and Environmental Studies

Kyoto University

Japan

E-mail: sumi@math.h.kyoto-u.ac.jp

<http://www.math.h.kyoto-u.ac.jp/~sumi/index.html>

August 17, 2020

Definition 1.

- (1) Let $\hat{\mathbb{C}} := \mathbb{C} \cup \{\infty\} \cong S^2$ be the Riemann sphere endowed with the spherical distance d .
- (2) Let $\text{Rat} := \{f : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}} \mid f \text{ is non-constant and holomorphic}\}$ endowed with the distance η , where $\eta(f, g) = \sup_{z \in \hat{\mathbb{C}}} d(f(z), g(z))$. Note that (Rat, η) is a complete separable metric space.
- (3) For a metric space Y , we denote by $\mathfrak{M}_1(Y)$ the space of all Borel probability measures on Y .
- (4) For a subset Y of Rat , we set

$$\mathfrak{M}_{1,c}(Y) := \{\tau \in \mathfrak{M}_1(Y) \mid \text{supp } \tau \text{ is a compact subset of } Y\}.$$

(5) For a $\tau \in \mathfrak{M}_{1,c}(\text{Rat})$, we set

$$G_\tau := \{\gamma_n \circ \cdots \circ \gamma_1 \mid n \in \mathbb{N}, \gamma_j \in \text{supp } \tau(\forall j)\}.$$

Note that this is a **semigroup** whose product is the composition of maps.

(6) We say that an element $\tau \in \mathfrak{M}_{1,c}(\text{Rat})$ is **weakly mean stable** if there exist an $n \in \mathbb{N}$, an $m \in \mathbb{N}$, non-empty open subsets U_1, \dots, U_m of $\hat{\mathbb{C}}$, a non-empty compact subset K of $\hat{\mathbb{C}}$ with $K \subset \cup_{j=1}^m U_j$, and a constant c with $0 < c < 1$ such that the following (a) (b) (c) hold.

(a) For each $(\gamma_1, \dots, \gamma_n) \in (\text{supp } \tau)^n$, we have

$$\gamma_n \circ \cdots \circ \gamma_1(\cup_{j=1}^m U_j) \subset K.$$

Moreover, for each $j = 1, \dots, m$, for all $x, y \in U_j$ and for each $(\gamma_1, \dots, \gamma_n) \in (\text{supp } \tau)^n$, we have

$$d(\gamma_n \circ \cdots \circ \gamma_1(x), \gamma_n \circ \cdots \circ \gamma_1(y)) \leq cd(x, y).$$

(b) Let $D_\tau := \bigcap_{h \in G_\tau} h^{-1}(\hat{\mathbb{C}} \setminus \cup_{j=1}^m U_j)$. Then $\#D_\tau < \infty$.

(c) For each minimal set L of τ with $L \subset D_\tau$, there exist a $z \in L$ and an $\alpha \in G_\tau$ such that $\alpha(z) = z$ and $|\alpha'(z)| > 1$ (if $z = \infty$ then we consider the condition $|(\varphi \circ \alpha \circ \varphi^{-1})'(0)| > 1$ instead of the condition $|\alpha'(z)| > 1$, where $\varphi(z) = 1/z$).

Here, a non-empty compact subset L of $\hat{\mathbb{C}}$ is said to be a **minimal set of τ** if for each $z \in L$, $\overline{\cup_{h \in G_\tau} \{h(z)\}} = L$.

(7) For each $Y \subset \text{Rat}$, we endow $\mathfrak{M}_{1,c}(Y)$ with the topology such that a sequence $\{\tau_n\}_{n \in \mathbb{N}}$ in $\mathfrak{M}_{1,c}(Y)$ tends to an element $\tau \in \mathfrak{M}_{1,c}(Y)$ if and only if

(*) for each bounded continuous function $\varphi : Y \rightarrow \mathbb{R}$, we have

$$\int_Y \varphi d\tau_n \rightarrow \int_Y \varphi d\tau \text{ as } n \rightarrow \infty,$$

and

(**) $\text{supp } \tau_n \rightarrow \text{supp } \tau$ as $n \rightarrow \infty$ with respect to the Hausdorff metric in the space of all non-empty compact subsets of Y .

Theorem 2 ([4]). *Let Y be one of the following (1)–(4).*

(1) $\{f \in \text{Rat} \mid f \text{ is a polynomial with } \deg(f) \geq 2\}$.

(2) $\{“z \mapsto \lambda z(1 - z)” \in \text{Rat} \mid \lambda \in \mathbb{C} \setminus \{0\}\}$.

(3) $\{“z \mapsto z - \lambda \frac{f(z)}{f'(z)}” \in \text{Rat} \mid \lambda \in \mathbb{C}, |\lambda - 1| < 1\}$
where f is a polynomial with $\deg(f) \geq 2$.

Remark: *This family is related to “random relaxed Newton’s methods for f ” in which we can find roots of any polynomial f more easily than deterministic Newton’s method ([4]).*

(4) $\{“z \mapsto z + \lambda f(z)” \in \text{Rat} \mid \lambda \in \mathbb{C} \setminus \{0\}\}$
*where f is a polynomial with $\deg(f) \geq 2$
such that for each $z_0 \in \mathbb{C}$ with $f(z_0) = 0$, we have $f'(z_0) \neq 0$.*

Then, there exists an open and dense subset A of $\mathfrak{M}_{1,c}(Y)$ such that for each $\tau \in A$, we have the following (I)(II)(III).

(I) τ is *weakly mean stable*.

(II) *There exists $c_\tau < 0$ s.t.*
for all but countably many $z \in \hat{\mathbb{C}}$, for $(\otimes_{n=1}^\infty \tau)$ -a.e. $(\gamma_1, \gamma_2, \dots) \in Y^\mathbb{N}$,
we have

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log \|D(\gamma_n \circ \dots \circ \gamma_1)_z\| \leq c_\tau < 0.$$

(III) *For all but countably many $z \in \hat{\mathbb{C}}$,*
for $(\otimes_{n=1}^\infty \tau)$ -a.e. $\gamma = (\gamma_1, \gamma_2, \dots) \in Y^\mathbb{N}$,
there exists a minimal set $L = L(z, \gamma)$ of τ
which is either

(a) *“attracting for τ ”, or*

(b) *included in D_τ with $\chi(\tau, L) < 0$,*

where $\chi(\tau, L)$ denotes the Lyapunov exponent of (τ, L) ,

such that

$$d(\gamma_n \circ \dots \circ \gamma_1(z), L) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Theorem 3. Let f be a polynomial with $\deg(f) \geq 2$ and let $Q := \{x \in \mathbb{C} \mid f(x) = 0\}$. Suppose that $f'(x) \neq 0$ ($\forall x \in Q$). Suppose also that for each $a, b \in Q$ with $a \neq b$, we have $f'(a) \neq f'(b)$. Let $Y = \{“z \mapsto z + \lambda f(z)” \in \text{Rat} \mid \lambda \in \mathbb{C} \setminus \{0\}\}$. For each $\tau \in \mathfrak{M}_{1,c}(Y)$ and $x \in Q$, let $\chi(\tau, x) := \int_Y \log |h'(x)| d\tau(h)$. Let A be the set of elements $\tau \in \mathfrak{M}_{1,c}(Y)$ satisfying that

- τ is weakly mean stable with $D_\tau \subset Q$,
- for each $x \in Q$, we have $\chi(\tau, x) \neq 0$, and
- if $x \in Q$ and $\chi(\tau, x) > 0$, then for each $h \in \text{supp } \tau$, we have $h'(x) \neq 0$.

Then, we have the following (i)(ii).

- (i) A is *open and dense* in $\mathfrak{M}_{1,c}(Y)$ and statements (II) and (III) in Theorem 2 hold for each $\tau \in A$.
- (ii) For any two subsets Q_1, Q_2 of Q , let $A_{Q_1, Q_2} = \{\tau \in A \mid \chi(\tau, x) > 0 (\forall x \in Q_1) \text{ and } \chi(\tau, x) < 0 (\forall x \in Q_2)\}$. Then A_{Q_1, Q_2} is a *non-empty open* subset of A . Moreover, we have $A = \coprod_{(Q_1, Q_2)} A_{Q_1, Q_2}$ (disjoint union).

Theorem 4. Let $a, x_1, x_2 \in \mathbb{C}$ with $x_1 \neq x_2$.

Let $f(z) = a(z - x_1)(z - x_2)$.

Let $Y = \{“z \mapsto z + \lambda f(z)” \in \text{Rat} \mid \lambda \in \mathbb{C} \setminus \{0\}\}$.

Let A be the set of elements $\tau \in \mathfrak{M}_{1,c}(Y)$ satisfying that

- τ is weakly mean stable with $D_\tau \subset \{x_1, x_2\}$,
- for each $i = 1, 2$, we have $\chi(\tau, x_i) \neq 0$, and
- if $i \in \{1, 2\}$ and $\chi(\tau, x_i) > 0$, then for each $h \in \text{supp } \tau$, we have $h'(x_i) \neq 0$.

Then, we have the following (i)–(iii).

- (i) A is *open and dense* in $\mathfrak{M}_{1,c}(Y)$ and statements (II) and (III) in Theorem 2 hold for each $\tau \in A$.
- (ii) For any $\tau \in A$, we have $D_\tau \neq \emptyset$.

- (iii) For each $\gamma = (\gamma_1, \gamma_2, \dots) \in Y^{\mathbb{N}}$, let F_γ be the set of points $z \in \hat{\mathbb{C}}$ satisfying that there exists a neighborhood U of z in $\hat{\mathbb{C}}$ such that $\{\gamma_n \circ \dots \circ \gamma_1\}_{n=1}^\infty$ is equicontinuous on U . Then for $(\otimes_{n=1}^\infty \tau)$ -a.e. γ , we have $\text{Leb}_2(\hat{\mathbb{C}} \setminus F_\gamma) = 0$ (Leb_2 denotes the Lebesgue meas. on $\hat{\mathbb{C}}$).

Theorem 5. Let f, Y, A, F_γ be as in Theorem 4.

For each $\tau \in \mathfrak{M}_{1,c}(Y)$, let $\text{Min}(\tau)$ be the set of all minimal sets of τ , and we set $\chi(\tau, x_i) := \int_Y \log |h'(x_i)| d\tau(h)$ ($i = 1, 2$).

Then, there exist non-empty open subsets A_1, A_2, \dots, A_5 of A with $A = \coprod_{i=1}^5 A_i$ (disjoint union) such that all of the following (1)–(5) hold.

- (1) Let $\tau \in A_1$. Then we have the following (i)–(iv).

- (i) $\text{Min}(\tau) = \{\{x_1\}, \{x_2\}, \{\infty\}, L_\tau\}$, where L_τ is an “attracting minimal set” of τ with $L_\tau \subset \mathbb{C} \setminus \{x_1, x_2\}$.
- (ii) For each $i = 1, 2$ and each $h \in \text{supp } \tau$, we have $|h'(x_i)| > 1$ and $D_\tau = \{x_1, x_2\}$.
- (iii) For all but countably many $z \in \hat{\mathbb{C}}$, for $(\otimes_{n=1}^\infty \tau)$ -a.e. $\gamma = (\gamma_1, \gamma_2, \dots) \in Y^{\mathbb{N}}$, we have

$$d(\gamma_n \circ \dots \circ \gamma_1(z), L_\tau \cup \{\infty\}) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

- (iv) For each $\gamma = (\gamma_1, \gamma_2, \dots) \in (\text{supp } \tau)^{\mathbb{N}}$, we have $L_\tau \cup \{\infty\} \subset F_\gamma$ and $\{x_1, x_2\} \subset \hat{\mathbb{C}} \setminus F_\gamma$.

- (2) Let $\tau \in A_2$. Then we have the following (i)–(v).

- (i) $\text{Min}(\tau) = \{\{x_1\}, \{x_2\}, \{\infty\}\}$.
- (ii) For each $i = 1, 2$, we have $\chi(\tau, x_i) > 0$.
- (iii) $D_\tau = \{x_1, x_2\}$.
- (iv) For all but countably many $z \in \hat{\mathbb{C}}$, for $(\otimes_{n=1}^\infty \tau)$ -a.e. $\gamma = (\gamma_1, \gamma_2, \dots) \in Y^{\mathbb{N}}$, we have

$$\gamma_n \circ \dots \circ \gamma_1(z) \rightarrow \infty \text{ as } n \rightarrow \infty.$$

(v) For $(\otimes_{n=1}^{\infty} \tau)$ -a.e. $\gamma = (\gamma_1, \gamma_2, \dots) \in Y^{\mathbb{N}}$,
we have

$$\infty \in F_{\gamma} \quad \text{and} \quad \{x_1, x_2\} \subset \hat{\mathbb{C}} \setminus F_{\gamma}.$$

(3) Let $\tau \in A_3$. Then we have the following (i)–(v).

(i) $\text{Min}(\tau) = \{\{x_1\}, \{x_2\}, \{\infty\}\}$.

(ii) $\chi(\tau, x_1) < 0$ and $\chi(\tau, x_2) > 0$.

(iii) Let $A_{3,a} := \{\tau \in A_3 \mid D_{\tau} = \{x_2\}\}$ and
 $A_{3,b} = \{\tau \in A_3 \mid D_{\tau} = \{x_1, x_2\}\}$.

Then $A_{3,a}, A_{3,b}$ are *non-empty open* subsets of A_3 and

$$A_3 = A_{3,a} \coprod A_{3,b} \quad (\text{disjoint union}).$$

(iv) For *all but countably many* $z \in \hat{\mathbb{C}}$,
for $(\otimes_{n=1}^{\infty} \tau)$ -a.e. $\gamma = (\gamma_1, \gamma_2, \dots) \in Y^{\mathbb{N}}$, we have

$$d(\gamma_n \circ \dots \circ \gamma_1(z), \{x_1, \infty\}) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

(v) For $(\otimes_{n=1}^{\infty} \tau)$ -a.e. $\gamma \in Y^{\mathbb{N}}$, we have $\{x_1, \infty\} \subset F_{\gamma}$ and $x_2 \in \hat{\mathbb{C}} \setminus F_{\gamma}$.

(4) Let $\tau \in A_4$. Then we have the following (i)–(v).

(i) $\text{Min}(\tau) = \{\{x_1\}, \{x_2\}, \{\infty\}\}$.

(ii) $\chi(\tau, x_1) > 0$ and $\chi(\tau, x_2) < 0$.

(iii) Let $A_{4,a} := \{\tau \in A_4 \mid D_{\tau} = \{x_1\}\}$ and
 $A_{4,b} = \{\tau \in A_4 \mid D_{\tau} = \{x_1, x_2\}\}$.

Then $A_{4,a}, A_{4,b}$ are *non-empty open* subsets of A_4 and

$$A_4 = A_{4,a} \coprod A_{4,b} \quad (\text{disjoint union}).$$

(iv) For *all but countably many* $z \in \hat{\mathbb{C}}$,
for $(\otimes_{n=1}^{\infty} \tau)$ -a.e. $\gamma = (\gamma_1, \gamma_2, \dots) \in Y^{\mathbb{N}}$, we have

$$d(\gamma_n \circ \dots \circ \gamma_1(z), \{x_2, \infty\}) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

(v) For $(\otimes_{n=1}^{\infty} \tau)$ -a.e. $\gamma \in Y^{\mathbb{N}}$, we have $\{x_2, \infty\} \subset F_{\gamma}$ and $x_1 \in \hat{\mathbb{C}} \setminus F_{\gamma}$.

(5) Let $\tau \in A_5$. Then we have the following (i)–(vi).

(i) $\text{Min}(\tau) = \{\{x_1\}, \{x_2\}, \{\infty\}\}$.

(ii) For each $i = 1, 2$, we have $\chi(\tau, x_i) < 0$.

(iii) $D_{\tau} = \{x_1, x_2\}$.

(iv) For **each** $z \in \hat{\mathbb{C}}$,
for $(\otimes_{n=1}^{\infty} \tau)$ -a.e. $\gamma = (\gamma_1, \gamma_2, \dots) \in Y^{\mathbb{N}}$, we have

$$d(\gamma_n \circ \dots \circ \gamma_1(z), \{x_1, x_2, \infty\}) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

(v) For $(\otimes_{n=1}^{\infty} \tau)$ -a.e. $\gamma \in Y^{\mathbb{N}}$, we have that $\{x_1, x_2, \infty\} \subset F_{\gamma}$.

(vi) For $(\otimes_{n=1}^{\infty} \tau)$ -a.e. $\gamma \in Y^{\mathbb{N}}$, for **each** $i = 1, 2$, for **each** point z in the connected component U_i of F_{γ} with $x_i \in U_i$, we have

$$\gamma_n \circ \dots \circ \gamma_1(z) \rightarrow x_i \text{ as } n \rightarrow \infty.$$

Remark. For any *deterministic iteration dynamics* of a single quadratic map f , we **CANNOT** have a phenomenon such as (vi). In fact, we **CANNOT** have two attracting minimal sets of f in \mathbb{C} .

Remark 6. Statements of Theorems 2, 3, 4, 5 **cannot hold for deterministic dynamics of a single $f \in \text{Rat}$ with $\deg(f) \geq 2$.**

In fact, in the Julia set $J(f)$ of f , we have a chaotic phenomenon. See Mañé's paper (1988)[1] etc.

Thus Theorems 2, 3, 4, 5 describe **randomness-induced phenomena** (new phenomena in random dynamical systems which cannot hold for deterministic dynamical systems).

Idea of Proofs of Theorems 2, 3, 4, 5.

- (1) We use complex analysis, [Montel's theorem](#) (a family of uniformly bounded holomorphic functions on a domain is equicontinuous on the domain), [hyperbolic metric](#).
- (2) We [classify minimal sets](#) and analyze the [bifurcation of minimal sets](#). etc. By using these, enlarging the support of the original τ a little bit, we [destroy non-attracting minimal sets](#) which do not meet D_τ .

Summary

- (1) We introduce the notion of [weak mean stability](#) in i.i.d. random (holomorphic) 1-dimensional dynamical systems.
- (2) If a random holomorphic dynamical system on $\hat{\mathbb{C}}$ is weakly mean stable and satisfies some mild assumptions, then for [all but countably many](#) $z \in \hat{\mathbb{C}}$, for [a.e.](#) orbit starting with z , [the Lyapunov exponent is negative](#). Note that this statement [cannot hold for deterministic dynamics of a single holo. map \$f\$ on \$\hat{\mathbb{C}}\$ with \$\deg\(f\) \geq 2\$](#) .
- (3) Given an analytic family Y of rational maps (with some mild conditions), [generic](#) random holomorphic dynamical systems (with multiplicative noise) of elements of Y are [weakly mean stable](#). Also, we can [classify](#) such generic random holomorphic dynamical systems of elements of Y in terms of [averaged behavior](#) and [quenched dynamics](#).

References:

- [1] R. Mañé, *The Hausdorff dimension of invariant probabilities of rational maps*, Dynamical Systems (Valparaiso, 1986) (Lecture Notes in Mathematics vol 1331) (Berlin: Springer) pp 86-117, 1988.
- [2] H. Sumi, *Random complex dynamics and semigroups of holomorphic maps*, Proc. London Math. Soc. (2011) 102(1), pp 50–112.
- [3] H. Sumi, *Cooperation principle, stability and bifurcation in random complex dynamics*, Adv. Math., 245 (2013) pp 137–181.

- [4] H. Sumi, *Negativity of Lyapunov Exponents and Convergence of Generic Random Polynomial Dynamical Systems and Random Relaxed Newton's Methods*, 61 pages, <https://arxiv.org/abs/1608.05230>.
Some contents of this talk are included in this paper.