

Hyperbolicity and Primitivity of Group Rings *

Brent B. Solie
Embry-Riddle Aeronautical University
solieb@erau.edu

May 28, 2020

A ring R is said to be (*right*) *primitive* if it contains a faithful irreducible (right) R -module. In order to show the primitivity of a group ring KG , Alexander and Nishinaka develop the following useful Property (*) for the group G [1].

(*) For each subset M of G consisting of a finite number of elements not equal to 1, and for any positive integer $m \geq 2$, there exist distinct $a, b, c \in G$ so that if $(x_1^{-1}g_1x_1)(x_2^{-1}g_2x_2) \cdots (x_m^{-1}g_mx_m) = 1$, where $g_i \in M$ and $x_i \in \{a, b, c\}$ for all $i = 1, \dots, m$, then $x_i = x_{i+1}$ for some i .

Equipped with Property (*), Alexander and Nishinaka obtain the following:

Theorem 1 ([1, Theorem 1.1]). *Let G be a group which has a non-Abelian free subgroup whose cardinality is the same as that of G , and suppose that G satisfies Property (*). Then, if R is a domain with $|R| \leq |G|$, the group ring RG of G over R is primitive. In particular, the group algebra KG is primitive for any field K .*

As seen in [1], Theorem 1 immediately implies the primitivity of group rings for a large class of groups, generalizing many classical results and obtaining several new ones. The present result, published in [9], shows that Theorem 1 also applies to the non-elementary torsion-free hyperbolic groups. Thus the class of groups satisfying Property (*) is indeed quite large and encompasses “almost all” groups in a particular statistical sense [7].

Let G be a group with finite generating set X . Recall that the *Cayley graph* $\Gamma_X(G)$ of G with respect to X is an X -digraph with vertex set G and an x -labelled edge directed from g to gx for all $g \in G$ and $x \in X$. We may promote $\Gamma_X(G)$ to a geodesic metric space by assigning each edge a length of one.

When there exists $\delta \geq 0$ such that each side of a geodesic triangle in $\Gamma_X(G)$ is contained in the δ -neighborhood of the remaining two sides, we say that $\Gamma_X(G)$ has *δ -thin triangles*. This thin triangle property is independent of the choice of

*A detailed version of this paper appears in Journal of Algebra 493 (2018) [9].

finite generating set, though δ may vary. When $\Gamma_X(G)$ has δ -thin triangles for some finite generating set X and $\delta \geq 0$, we say G is *hyperbolic*. A hyperbolic group is *non-elementary* if it is not virtually cyclic.

Hyperbolic groups enjoy a close relationship between their algebraic and geometric properties; see, for instance, [2, 4, 5]. For our purposes, we require the so called “big powers property” of torsion-free hyperbolic groups. The version stated here follows immediately from a more general version for certain relatively hyperbolic groups given in [6].

Theorem 2 (The big powers property [6]). *Let G be a torsion-free hyperbolic group. Let $u \in G$ be nontrivial and not a proper power. Let g_1, \dots, g_k be elements of G which do not commute with u . Then there exists $N > 0$ such that if $|n_i| \geq N$ for $i = 0, \dots, k$ then*

$$u^{n_0} g_1 u^{n_1} g_2 \cdots u^{n_{k-1}} g_k u^{n_k} \neq 1.$$

The big powers property allows one to programmatically generate large sets of nontrivial elements of G , and has seen useful application towards residual properties, logic, and algebraic geometry [3, 8, 6].

Proposition 3. *If G is a non-elementary torsion-free hyperbolic group, then G satisfies Property (*).*

Proof. Let M be a finite subset of G not containing the identity. A classical result due to Gromov asserts that the subgroup generated by sufficiently high powers of elements of M must be free. Since nontrivial elements of a non-elementary torsion free hyperbolic group have maximal infinite cyclic centralizers, one can therefore find an element $u \in G$ which generates its own centralizer and commutes with no $g \in M$.

Let $m \geq 2$ be an integer and consider a finite sequence g_1, \dots, g_m of elements from M . Since u commutes with none of the g_i and generates its own centralizer, the big powers property gives $N(g_1, \dots, g_m) > 0$ such that

$$u^{n_0} g_1 u^{n_1} g_2 \cdots g_{m-1} u^{n_{m-1}} g_m u^{n_m} \neq 1$$

whenever $|n_i| \geq N$ for all $i = 0, \dots, m$.

Since M is a finite set, there are finitely many m -tuples (g_1, \dots, g_m) of elements from M . Therefore, let $N > \max \{N(g_1, \dots, g_m) \mid g_1, \dots, g_m \in M\}$.

We now define $a = u^N, b = u^{2N}$, and $c = u^{3N}$. Since G is torsion-free, these elements are distinct. Consider a product

$$w = (x_1^{-1} g_1 x_1)(x_2^{-1} g_2 x_2) \cdots (x_m^{-1} g_m x_m)$$

where $x_1, \dots, x_m \in \{u^N, u^{2N}, u^{3N}\}$. We then have

$$w = u^{n_0} g_1 u^{n_1} g_2 \cdots g_{m-1} u^{n_{m-1}} g_m u^{n_m},$$

where $u^{n_0} = x_1^{-1}, u^{n_m} = x_m, u^{n_i} = x_i x_{i+1}^{-1}$ and $n_i \in \{0, \pm N, \pm 2N\}$ for $i = 1, \dots, m-1$. Note that by choice of x_1 and x_m , we have $n_0 \neq 0$ and $n_m \neq 0$.

By the big powers property and choice of N , if $n_i \neq 0$ for all $i = 0, \dots, m$, then $w \neq 1$. Therefore, if $w = 1$, then some $n_i = 0$. Since we cannot have $n_0 = 0$ or $n_{m+1} = 0$, we have $n_i = 0$ for some $i \in \{1, \dots, m-1\}$, in which case we must have $1 = u^{n_i} = x_i x_{i+1}^{-1}$, and so $x_i = x_{i+1}$. \square

We immediately obtain the following main result as a corollary to Theorem 1.

Theorem 4. *If G is a non-elementary torsion-free hyperbolic group, then for any countable domain R , the group ring RG of G over R is primitive. In particular, the group ring KG is primitive for any field K .*

References

- [1] James Alexander and Tsunekazu Nishinaka. Non-noetherian groups and primitivity of their group algebras. *J. Algebra*, 473:221–246, 2017.
- [2] J. M. Alonso, T. Brady, D. Cooper, V. Ferlini, M. Lustig, M. Mihalik, M. Shapiro, and H. Short. Notes on word hyperbolic groups. In *Group theory from a geometrical viewpoint (Trieste, 1990)*, pages 3–63. World Sci. Publ., River Edge, NJ, 1991. Edited by Short.
- [3] Benjamin Baumslag. Residually free groups. *Proc. London Math. Soc. (3)*, 17:402–418, 1967.
- [4] Étienne Ghys and Pierre de la Harpe. La propriété de Markov pour les groupes hyperboliques. In *Sur les groupes hyperboliques d’après Mikhael Gromov (Bern, 1988)*, volume 83 of *Progr. Math.*, pages 165–187. Birkhäuser Boston, Boston, MA, 1990.
- [5] M. Gromov. Hyperbolic groups. In *Essays in group theory*, volume 8 of *Math. Sci. Res. Inst. Publ.*, pages 75–263. Springer, New York, 1987.
- [6] O. Kharlampovich and A. Myasnikov. Limits of relatively hyperbolic groups and lyndon’s completions. arXiv:0904.2423v3 [math.GR], 2009.
- [7] A. Yu. Ol’shanskiĭ. Almost every group is hyperbolic. *Internat. J. Algebra Comput.*, 2(1):1–17, 1992.
- [8] A. Yu. Ol’shanskiĭ. On residualizing homomorphisms and G -subgroups of hyperbolic groups. *Internat. J. Algebra Comput.*, 3(4):365–409, 1993.
- [9] Brent B. Solie. Primitivity of group rings of non-elementary torsion-free hyperbolic groups. *J. Algebra*, 493:438–443, 2018.