

On formalization of logic puzzles *à la* Smullyan*

Ken-etsu Fujita (Gunma University)

February 18, 2021

Abstract

George Boolos (1996) posed the puzzle “The hardest logic puzzle ever” which had been devised by Raymond Smullyan, and gave a solution in the style of if-and-only-if. Later, Roberts (2001) and Rabern-Rabern (2008) provided another solution in the style of embedded questions as a simpler solution. Here, we introduce a simple formalization of the puzzle consisting of questions, answerers, and answers in terms of propositional logic, and show its adequacy by the truth values $(0, 1)$ semantics. Then it turns out that the two solutions in the different forms are logically equivalent under the semantics. Moreover, the hardest logic puzzle can be considered as a natural extension of the puzzles of knights and knaves, i.e., lying and truth-telling by Smullyan.

1 Introduction

To begin with, we quote the puzzle from Boolos [1]:

“The puzzle: Three gods, A, B, and C are called, in some order, True, False, and Random. True always speaks truly, False always speaks falsely, but whether Random speaks truly or falsely is a completely *random* matter. Your task is to determine the identities of A, B, C by asking three yes-no questions; each question must be put to exactly one god. The gods understand English, but will answer all questions in their own language, in which the words for “yes” and “no” are “da” and “ja,” in some order. *You do not know which word means which.*”

His solution in the style of if-and-only-if consists of the following steps.

1. Ask god A:
does da means yes iff, you are True iff B is Random?
2. Ask B or C¹:
does da mean yes iff Rome is in Italy?

*This work was supported by the Research Institute for Mathematical Science, a Joint Usage/Research Center located in Kyoto University. This work was partly supported by Grants-in-Aid for Scientific Research KAKENHI (C) 17K05343 and 20K03711.

¹If the answer is “da” then ask C, otherwise (i.e., “ja”) B.

3. Does da mean yes iff A is Random?

On the other hand, Roberts [4] and Rabern-Rabern [3] provided another solution in the style of embedded questions as follows.

1. Ask god A:

if I asked you if god B was Random, would you say da?

2. Ask B or C¹:

if I asked you if you always told the truth, would you say da?

3. If I asked you if god A was Random, would you say da?

Before formalizing the questions of the puzzle above, we call one of the most elementary puzzle of Knights and Knaves from the book Logical Labyrinths [2]. This puzzle is known as a good example for the study of logic or sociology of *lying* and *truth-telling*. There were the island of knights and knaves such that *knights* always tell the truth, and *knaves* always lie. Each inhabitant of the island is either a knight or a knave.

• Problem 1.3. [2]:

The island has two inhabitants, A and B. Now, A made the following statement:

“Both of us are knaves.”

What is A and what is B?

From the book, we review the simple solution to this puzzle. Let A, B be propositional variables which mean that A is a knight and B is a Knight, respectively. Then $\neg A$ means that A is *not* a knight (i.e., knave) from the definition of knights and knaves. Suppose A asserts a proposition which is expressed by a formula X . Then the definition of knights leads to the following fact.

• The inhabitant A is a knight *if and only if* X is true.

The fact can be formalized by the formula of bi-implication:

$$A \leftrightarrow X$$

The formalization of the well-formed relation between inhabitants and assertions can be justified by the truth table of bi-implication, following the case analysis on A of either a knight or a knave.

A	X	$A \leftrightarrow X$
t	t	t
t	f	f
f	t	f
f	f	t

Now recall problem 1.3, and then we obtain the formula $A \leftrightarrow \neg A \wedge \neg B$ from the statement of A. The solution of the puzzle is given by solving the satisfiability problem (SAT) of the formula as follows:

A	B	$\neg A$	$\neg B$	$\neg A \wedge \neg B$	$A \leftrightarrow \neg A \wedge \neg B$
t	t	f	f	f	f
t	f	f	t	f	f
$\textcircled{\text{f}}$	$\textcircled{\text{t}}$	t	f	f	$\textcircled{\text{t}}$
f	f	t	t	t	f

In other words, the puzzle and its solution form the relation of logical consequence:

$$A \leftrightarrow \neg A \wedge \neg B \models \neg A \wedge B$$

2 Formalization

This methodology for solving the puzzles of lying and truth-telling can be naturally extended to that for the hardest logic puzzle ever, so that the binary relation between inhabitants and assertions should be replaced with a ternary form of questions, answerers, and answers. Firstly, let X be a propositional variable for a question, which means either true or false, respectively represented by 1 or 0. Secondly, let A, B be propositional variables for answerers A, B, which mean either² True or False, respectively represented by 1 or 0. Lastly, let Y be a propositional variable for an answer. Here, an answer means either yes or no, respectively represented by 1 or 0. Instead, Y may be used for an answer da-ja, whose meaning is also either 1 or 0, but not fixed yet.

If we ask a question X of an answerer A and obtain an answer Y , then the situation is depicted by the following diagram.

$$X \longrightarrow \boxed{A} \longrightarrow Y$$

We formalize this relation of question-answerer-answer by the ternary form with the logical connective of bi-implication.

$$X \leftrightarrow A \leftrightarrow Y$$

Note trivial facts that \leftrightarrow is symmetric and associative and that a tautology is the unit. An adequacy of the formalization can be expounded by the truth values $(0, 1)$ semantics. Let **Prop** be the set of propositions (formulae), and $\{0, 1\}$ for the set of truth values. We write v for the assignment $v : \mathbf{Prop} \rightarrow \{0, 1\}$. Now the consistent relation of question-answerer-answer is stated as follows:

$$v(X \leftrightarrow A \leftrightarrow Y) = 1,$$

²In the puzzle of Boolos [1], one has Random in addition, but for the formalization here answerers are supposed to be either True or False. According to the solutions [1, 4, 3], Random can be handled by a certain *strategy* of asking questions which can be formalized here in terms of propositional logic.

under the assignment v such that $v(A := \text{“True”}) = v(Y := \text{“yes”}) = 1$. This statement can be justified by the case analysis on A , as follows.

- Case A of “True”: $v(X \leftrightarrow Y) = 1$

$$X \longrightarrow \boxed{\text{True}} \longrightarrow Y$$

- Case A of “False”: $v(X \leftrightarrow Y) = 0$

$$X \longrightarrow \boxed{\text{False}} \longrightarrow Y$$

Let us formalize the embedded question [4, 3]. Recall the first one from the solution:

- Ask god A Q_1 : if I asked you if god B was Random, would you say da?

$$\text{“}X \rightarrow \boxed{A} \rightarrow Y\text{”} \longrightarrow \boxed{A} \longrightarrow ?_1$$

where $X := \text{“B is Random”}$, $Y := \text{“da”}$.

Now suppose that A’s answer $?_1$ is “da”. Then this situation is formalized by the formula $(X \leftrightarrow A \leftrightarrow Y) \leftrightarrow A \leftrightarrow Y$, and hence for any assignment v we have the following equation

$$v((X \leftrightarrow A \leftrightarrow Y) \leftrightarrow A \leftrightarrow Y) = v(X),$$

since $A \leftrightarrow A$ and $Y \leftrightarrow Y$ are tautologies. This implies that one can identify the truth value of X from $?_1$, even if we know neither the semantics of A nor that of “da”. Moreover, this equation means that the following forms of question-answerer-answer are equivalent to each other under the semantics.

$$\begin{aligned} & \text{“}X \rightarrow \boxed{A} \rightarrow Y\text{”} \longrightarrow \boxed{A} \longrightarrow Y \\ \iff & X \longrightarrow \boxed{\text{True}} \longrightarrow \text{“yes”} \\ \iff & \text{“}X \text{ iff } A \text{ iff } Y\text{”} \longrightarrow \boxed{A} \longrightarrow Y \end{aligned}$$

Next, in order to analyze Boolos’ solution in the style of iff [1], we recall his solution:

- Ask god A Q_1 : does da means yes iff, you are True iff B is Random?

$$\text{“}(Y \text{ iff } Y_1) \text{ iff, } A \text{ iff } X\text{”} \longrightarrow \boxed{A} \longrightarrow ?$$

where $X := \text{“B is Random”}$, $Y := \text{“da”}$, $Y_1 := \text{“yes”}$.

Here, suppose A’s answer $?$ is “da”. Then we obtain the formula $((Y \leftrightarrow Y_1) \leftrightarrow (A \leftrightarrow X)) \leftrightarrow A \leftrightarrow Y$, so that for any assignment v the equation holds true

$$v(((Y \leftrightarrow Y_1) \leftrightarrow (A \leftrightarrow X)) \leftrightarrow A \leftrightarrow Y) = v(X \leftrightarrow Y_1).$$

Thanks to the semantics $v(X \leftrightarrow Y_1) = 1$, this form in the style of iff is equivalent to the definition of “True” as the embedded question is.

$$X \longrightarrow \boxed{\text{True}} \longrightarrow \text{“yes”}$$

Finally, we quote yet another puzzle from the film Labyrinth [5]. There are two doors with two guards, to say, A and B, either True or False. Your task is to determine whether which door leads to the castle by asking A or B one question. What kind of questions makes you reach the castle? In the film, Sarah asked A: Would he (B) tell me that this door leads to the castle? That is,

$$\text{“}X \rightarrow \boxed{B} \rightarrow Y\text{”} \longrightarrow \boxed{A} \longrightarrow ?$$

where $X := \text{“This door leads to the castle”}$, $Y := \text{“yes”}$. Now, suppose A’s answer ? is “no”, i.e., $\neg Y$. Then we obtain the formula $(X \leftrightarrow B \leftrightarrow Y) \leftrightarrow A \leftrightarrow \neg Y$, and hence for any assignment v the equation holds

$$v((X \leftrightarrow B \leftrightarrow Y) \leftrightarrow A \leftrightarrow \neg Y) = v(X),$$

where $B \leftrightarrow \neg A$. Note also that $A \leftrightarrow \neg A$ and $Y \leftrightarrow \neg Y$ are equivalent to the contradiction \perp where $v(\perp) = 0$ for any v , and of course, $\perp \leftrightarrow \perp$ is a tautology. Moreover, this methodology is still available for the setting of da-ja instead of yes-no.

3 Concluding remarks

We introduced a simple formalization of puzzles of questions-answerers-answer

$$X \longrightarrow \boxed{A} \longrightarrow Y$$

by using bi-implications $X \leftrightarrow A \leftrightarrow Y$, and the formalization can be justified under the truth values semantics. This method makes it possible to apply algebraic properties of the connective rather than making truth-tables. It turns out that Boolos’ solution in the style of if-and-only-if [1] and the solution in the style of embedded question [4, 3] are logically equivalent under this semantics. The hardest logic puzzle ever can be regarded as a generalization of Smullyan’s Knights-Knaves puzzles [2]. We show that this method is also applicable elegantly to the puzzle in the film Labyrinth [5]. Moreover, this method can formalize naturally n -times nesting of embedded question:

$$\text{“}X \rightarrow \boxed{A_1} \rightarrow Y_1\text{”} \rightarrow \boxed{A_2} \rightarrow Y_2 \rightarrow \cdots \rightarrow \boxed{A_n} \rightarrow Y_n$$

We make remarks on the definition of Random. According to [1], whether Random speaks truly or not should be thought of as depending on the flip of a coin hidden in his brain: if the coin comes down heads, he speaks truly; if tails,

falsely. Random will answer da or ja when asked any yes-no question, so that the definition can be depicted in the following.

$$X \rightarrow \boxed{R} \rightarrow \begin{cases} \text{“da”} & \begin{cases} \text{yes} & \text{for } X = 1 \\ \text{no} & \text{for } X = 0 \end{cases} & \text{if } R = T, \\ \text{“ja”} & \begin{cases} \text{yes} & \text{for } X = 0 \\ \text{no} & \text{for } X = 1 \end{cases} & \text{if } R = F. \end{cases}$$

That is, to put it simply, the diagram becomes the following one.

$$X \rightarrow \boxed{R} \rightarrow \begin{cases} \text{“da”} & \text{if heads,} \\ \text{“ja”} & \text{if tails.} \end{cases}$$

Here, we established a very good question (embedded question), so that one can verify whether X is 1 or 0 independent of the values of “da” and R . However, if the question X contains R as an answerer (i.e., embedded questions for R), when and how often does R flip a coin? If Random flips a coin everywhere for each question, for instance, in the diagram below:

$$\text{“}X \rightarrow \boxed{R} \rightarrow Y\text{”} \rightarrow \boxed{R} \rightarrow ?,$$

then one cannot enjoy the good property of the embedded questions. Fortunately, the solutions [1, 4, 3] of the puzzle can be provided without answering the question of when-and-how-often, by using an elegant strategy of asking three questions (Q_1, Q_2, Q_3) based on the case analysis such that $A = R \vee A \neq R$.

Finally, following our formalization we summarize the solution [4, 3] which consists of the questions Q_1, Q_2, Q_3 in this order, depending on the answer $?_1$:

1. Q_1 (Ask god A: if I asked you if god B was Random, would you say da?)

$$\text{“}B = R \rightarrow \boxed{A} \rightarrow \text{da”} \rightarrow \boxed{A} \rightarrow ?_1$$

2. $Q_2(Z := C)$ if $?_1 = \text{da}$ ($Q_2(Z := B)$ otherwise (i.e. $?_1 = \text{ja}$))

$$\text{“}Z = T \rightarrow \boxed{Z} \rightarrow \text{da”} \rightarrow \boxed{Z} \rightarrow ?_2$$

3. $Q_3(Z := C)$ (otherwise $Q_3(Z := B)$)

$$\text{“}A = R \rightarrow \boxed{Z} \rightarrow \text{da”} \rightarrow \boxed{Z} \rightarrow ?_3$$

As a solution we have $3!$ patterns consisting of R, T, or F for $\langle A, B, C \rangle$, and 2^3 patterns $\langle ?_1, ?_2, ?_3 \rangle$ for an answer to $\langle Q_1, Q_2, Q_3 \rangle$. Every candidate for $\langle A, B, C \rangle$ and $\langle Q_1, Q_2, Q_3 \rangle$, and the correlation are compacted in the simple table.

	A	B	C	Q_1	$Q_2(C)$	$Q_3(C)$?
1-1	R	T	F	da	ja	da	1
2-1	R	F	T	da	da	da	2
3	T	R	F	da	ja	ja	3
4	F	R	T	da	da	ja	4

	A	B	C	Q_1	$Q_2(B)$	$Q_3(B)$?
1-2	R	T	F	ja	da	da	5
2-2	R	F	T	ja	ja	da	6
5	T	F	R	ja	ja	ja	7
6	F	T	R	ja	da	ja	8

References

- [1] George Boolos: The Hardest Logic Puzzle Ever, *The Harvard Review of Philosophy* **6**, pp. 62–65, 1996.
- [2] Raymond Smullyan: *Logical Labyrinths*, A K Peters, Ltd 2009. (高橋昌一郎 監訳, 川辺治之 訳: スマリヤン 記号論理学, 丸善出版 2013)
- [3] Brian Rabern and Landon Rabern: A simple solution to the hardest logic puzzle ever, *ANALYSIS* **68.2**, pp 105–112, 2008.
- [4] Tim S. Roberts: Some Thoughts About The Hardest Logic Puzzle Ever, *Journal of Philosophical Logic* **30**, pp. 609–612, 2001.
- [5] Jim Henson with George Lucas: *Labyrinth*, Lucasfilm Ltd. 1986.