

Note on weighted q -Fock spaces and q -orthogonal polynomials

Nobuhiro ASAI (浅井暢宏) *
Department of Mathematics,
Aichi University of Education,
Kariya 448-8542, Japan.

Abstract

In this short note, we shall discuss weighted q -Fock spaces, field operators and their vacuum distributions, which have strong connections with q -orthogonal polynomials including discrete q -Hermite I polynomials. One can see that our general approach can treat not only known examples scattered in [1][5][8][9][10][13], but also can involve non-trivial and interesting examples, which were not referred in previous works [5][11]. This is a summary paper of our paper [4].

1 Weighted q -Deformation

Let \mathcal{H} be a complex Hilbert space equipped with the inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$, where the inner product is linear on the right and conjugate linear on the left. Let $\mathcal{F}_{\text{fin}}(\mathcal{H})$ denote the algebraic full Fock space over \mathcal{H} ,

$$\mathcal{F}_{\text{fin}}(\mathcal{H}) := \mathbb{C}\Omega \oplus \bigoplus_{n=1}^{\infty} \mathcal{H}^{\otimes n},$$

where Ω denotes the vacuum vector. We note that elements of $\mathcal{F}_{\text{fin}}(\mathcal{H})$ are expressed as finite linear combinations of the elementary vectors $f_1 \otimes \cdots \otimes f_n \in \mathcal{H}^{\otimes n}$. We equip $\mathcal{F}_{\text{fin}}(\mathcal{H})$ with the inner product

$$\langle f_1 \otimes \cdots \otimes f_m, g_1 \otimes \cdots \otimes g_n \rangle_0 := \delta_{m,n} \prod_{k=1}^n \langle f_k, g_k \rangle, \quad f_k, g_k \in \mathcal{H}.$$

For $q \in (-1, 1)$, define the q -symmetrization operator on $\mathcal{H}^{\otimes n}$ as

$$P_q^{(n)} = \sum_{\sigma \in \mathfrak{S}_n} q^{\ell(\sigma)} \sigma, \quad n \geq 1,$$
$$P_q^{(0)} = I_{\mathcal{H}^{\otimes 0}}, \quad P_0^{(n)} = I_{\mathcal{H}^{\otimes n}},$$

where we put $0^0 = 1$ and $\mathcal{H}^{\otimes 0} = \mathbb{C}\Omega$ by convention, \mathfrak{S}_n denotes the n -th symmetric group of permutations and $\ell(\sigma)$ means the number of inversion of a permutation $\sigma \in \mathfrak{S}_n$ defined by

$$\ell(\sigma) = \#\{(i, j) \mid 1 \leq i < j \leq n, \sigma(i) > \sigma(j)\}.$$

*Supported by JSPS KAKENHI Grant Numbers JP20K03652.

Definition 1.1 ([11]). Let $\{\tau_n\}_{n=1}^\infty$ be a sequence of strictly positive numbers and $[\tau_n]! := \prod_{i=1}^n \tau_i$. The τ -weighted q -symmetrization operators on $\mathcal{H}^{\otimes n}$ and $\mathcal{F}(\mathcal{H})$, respectively, are defined by

$$T_q^{(0)} = P_q^{(0)}, \quad T_q^{(n)} = [\tau_n]! P_q^{(n)}, \quad n \geq 1,$$

$$T_q = \bigoplus_{n=0}^{\infty} T_q^{(n)}.$$

Since $P_q^{(n)}$ and $\{\tau_n\}_{n=1}^\infty$ are a strictly positive operator and sequence, respectively, the τ -weighted q -inner product is defined by

$$\langle f_1 \otimes \cdots \otimes f_m, g_1 \otimes \cdots \otimes g_n \rangle_{q, \{\tau_n\}} := \delta_{m,n} \langle f_1 \otimes \cdots \otimes f_m, T_q^{(n)}(g_1 \otimes \cdots \otimes g_n) \rangle_0.$$

Let $\mathcal{F}_{q, \{\tau_n\}}(\mathcal{H})$ denote the τ -weighted (generalized) q -Fock space. In this paper, we do not take completion. The τ -weighted q -creation operator $b_{q, \{\tau_n\}}^\dagger(f)$ is defined as the usual left creation operator and $b_{q, \{\tau_n\}}(f)$ is its adjoint with respect to $\langle \cdot, \cdot \rangle_{q, \{\tau_n\}}$, that is, $b_{q, \{\tau_n\}} = (b_{q, \{\tau_n\}}^\dagger)^*$.

Proposition 1.2. (1) The τ -weighted q -annihilation operator $b_{q, \{\tau_n\}}$ acting on the elementary vectors is given as follows:

$$b_{q, \{\tau_n\}}(f)\Omega = 0, \quad b_{q, \{\tau_n\}}(f)f_1 = \tau_1 \langle f, f_1 \rangle \Omega, \quad f \in \mathcal{H},$$

$$b_{q, \{\tau_n\}}(f)(f_1 \otimes \cdots \otimes f_n) = \tau_n \sum_{k=1}^n q^{k-1} \langle f, f_k \rangle f_1 \otimes \cdots \otimes \overset{\vee}{f_k} \otimes \cdots \otimes f_n, \quad n \geq 2,$$

where $\overset{\vee}{f_k}$ means that f_k should be deleted from the tensor product.

(2) The τ -weighted q -creation and annihilation operators satisfy

$$b_{q, \{\tau_n\}}(f)b_{q, \{\tau_n\}}^\dagger(g) - q\beta_N b_{q, \{\tau_n\}}^\dagger(g)b_{q, \{\tau_n\}}(f) = \langle f, g \rangle \tau_{N+1}, \quad f, g \in \mathcal{H},$$

where $\{\beta_n := \tau_{n+1}/\tau_n\}_{n=1}^\infty$ and operators β_N and τ_N are defined as

$$\begin{cases} \varphi_N \Omega = \Omega, \quad \varphi_N(f_1 \otimes \cdots \otimes f_n) = \varphi_n(f_1 \otimes \cdots \otimes f_n), \quad n \geq 1, \\ \varphi \in \{\beta, \tau\}. \end{cases}$$

Corollary 1.3. Suppose $\tau_1 = 1$ and $\beta_n = Q > 0$ for $n \geq 1$. The following commutation relation holds:

$$b_{q, \{\tau_n\}}(f)b_{q, \{\tau_n\}}^\dagger(g) - qQ b_{q, \{\tau_n\}}^\dagger(g)b_{q, \{\tau_n\}}(f) = \langle f, g \rangle \tau_{N+1}, \quad f, g \in \mathcal{H}.$$

2 Examples

Let us begin with the following examples to proceed our discussion.

Example 2.1. Suppose $\tau_1 = 1$ and $q \in (-1, 1)$.

(1) $Q = 1$ implies $\tau_n = \tau_2 > 0$, $n \geq 2$. If we set $\tau_2 = t$, then one can get $T_q^{(n)} = t^{n-1} P_q^{(n)}$ and the $(q, t)_W$ -Fock space in the sense of Wojakowski [12]. If we take $q = 0$, one can derive the t -free deformation done by Bożejko-Wysoczański [9][10]. Moreover, if $\tau_n = 1$ for all $n \geq 1$, one can recover the well-known q -Fock space of Bożejko-Speicher [8] (See also [7]).

(2) If $Q = s^2$, $s \in (0, 1]$, then we have $\tau_n = s^{2(n-1)}$, $n \geq 1$. One can get $T_q^{(n)} = s^{n(n-1)} P_q^{(n)}$ and the $(q, s)_{BY}$ -Fock space by Bożejko-Yoshida [11]. The s -free deformation of Yoshida [13] can be

derived if $q = 0$. Moreover, one can see that a limiting case of $(q, s)_{BY}$ as $q \rightarrow 1$ coincides with the Q^N -deformation of the Boson Fock space [1].

(3) The Boolean Fock space can be derived as a limiting case of the $(0, t)_W$ -Fock space as $t \rightarrow 0$ and also $(0, s)_{BY}$ -Fock space as $s \rightarrow 0$.

One can derive a further deformation from (2) in Example 2.1. We shall show the relationship between Blitvić [5] construction and ours.

Remark 2.2. In [6], Bożejko-Ejmsmont-Hasebe constructed the (α, q) -Fock space, which is different from the (q, t) -Fock space by Blitvić [5]. In this note, the expression “ $\{q, t\}$ ” will be used to refer a symbol “ (q, t) ” to avoid confusions with the (α, q) -deformation.

In fact, if we replace s^2 by $t > 0$ in (2) of Example 2.1, then we have $\tau_n = t^{n-1}$ and $[\tau_n]! = t^{\binom{n}{2}}$ for $n \geq 1$. In addition, if one considers the (q/t) -symmetrization operator $P_{q/t}^{(n)}$, which is strictly positive for $|q| < t$, then one can consider the weighted (q/t) -symmetrization operator in forms of $T_{q/t}^{(0)} = P_{q/t}^{(0)}$ and $T_{q/t}^{(n)} = t^{\binom{n}{2}} P_{q/t}^{(n)}$, $n \geq 1$. From now on, we set

$$Q_{q,t}^{(0)} = T_{q/t}^{(0)}, \quad Q_{q,t}^{(n)} = T_{q/t}^{(n)}, \quad n \geq 1, \quad |q| < t,$$

$$Q_{q,t} = \bigoplus_{n=0}^{\infty} Q_{q,t}^{(n)},$$

which are called the $\{q, t\}$ -symmetrization operators on $\mathcal{H}^{\otimes n}$ and $\mathcal{F}(\mathcal{H})$, respectively. An inner product defined by

$$\langle f_1 \otimes \cdots \otimes f_m, g_1 \otimes \cdots \otimes g_n \rangle_{q,t} := \delta_{m,n} \langle f_1 \otimes \cdots \otimes f_m, Q_{q,t}^{(n)}(g_1 \otimes \cdots \otimes g_n) \rangle_0$$

is called the $\{q, t\}$ -inner product, which is the $(q/t, \sqrt{t})_{BY}$ -inner product. The free Fock space equipped with this $\{q, t\}$ -inner product is called the $\{q, t\}$ -Fock space denoted by $\mathcal{F}_{q,t}(\mathcal{H})$. Therefore, we have seen the following propositions:

Proposition 2.3. *Suppose $q \in (-1, 1)$, $t \in (0, 1]$ and $|q| < t$. The $(q/t, \sqrt{t})_{BY}$ -Fock space is equivalent to the $\{q, t\}$ -Fock space in the sense of [5].*

The $\{q, t\}$ -creation operator $a_{q,t}^\dagger(f)$ is defined as the usual left creation operator and $\{q, t\}$ -annihilation operator $a_{q,t}(f)$ as its adjoint with respect to $\langle \cdot, \cdot \rangle_{q,t}$. By replacing q by q/t and setting $Q = t$, $\tau_n = t^{n-1}$ in Proposition 1.2 and Corollary 1.3, then one can get the following proposition.

Proposition 2.4. (1) *The $\{q, t\}$ -annihilation operator $a_{q,t}$ acting on the elementary vectors is given as follows:*

$$a_{q,t}(f)\Omega = 0, \quad a_{q,t}(f)f_1 = \langle f, f_1 \rangle \Omega, \quad f \in \mathcal{H},$$

$$a_{q,t}(f)(f_1 \otimes \cdots \otimes f_n) = t^{n-1} \sum_{k=1}^n \left(\frac{q}{t}\right)^{k-1} \langle f, f_k \rangle f_1 \otimes \cdots \otimes \overset{\vee}{f_k} \otimes \cdots \otimes f_n \quad n \geq 2, \quad (2.1)$$

where $\overset{\vee}{f_k}$ means that f_k should be deleted from the tensor product.

(2) *The $\{q, t\}$ -creation and annihilation operators satisfy*

$$a_{q,t}(f)a_{q,t}^\dagger(g) - qa_{q,t}^\dagger(g)a_{q,t}(f) = \langle f, g \rangle t^N, \quad f, g \in \mathcal{H},$$

where the operator t^N is defined by

$$t^N \Omega = \Omega, \quad t^N(f_1 \otimes \cdots \otimes f_n) = t^n f_1 \otimes \cdots \otimes f_n, \quad n \geq 1.$$

We would like to consider the spectral measure (vacuum distribution) of the $\{q, t\}$ -Gaussian (field) operator $g_{q,t}(f)$ on $\mathcal{F}_{q,t}(\mathcal{H})$ defined by

$$g_{q,t}(f) := a_{q,t}^\dagger(f) + a_{q,t}(f), \quad f \in \mathcal{H},$$

with respect to the vacuum state $\langle \Omega, \cdot \rangle_{q,t}$. Orthogonal polynomials play important roles to compute a distribution of such a field operator with respect to the vacuum state. In [5], the $\{q, t\}$ -Hermite polynomials given by the recurrence relation,

$$\begin{aligned} H_0(x; q, t) &= 1, \quad H_1(x; q, t) = x, \\ xH_n(x; q, t) &= H_{n+1}(x; q, t) + [n]_{q,t}H_{n-1}(x; q, t), \quad n \geq 1, \end{aligned}$$

where $[n]_{q,t} := t^{n-1}[n]_{q/t}$ are mentioned. Note that $[n]_q := [n]_{q,1} = \sum_{k=0}^{n-1} q^k$ and $[n]_{q,q} = q^{n-1}n$. However, concrete densities of orthogonalizing measures are not mentioned except for a very restricted case, $0 = q < t$. We have been seeking examples for $q \neq 0$, which can be treated within the $\{q, t\}$ -deformation. In this paper, we shall present not only recognized examples, but also unrecognized ones in [5][11] as follows.

Example 2.5. Let us consider the $\{qs^2, s^2\}$ -deformation for $q \in (-1, 1), s \in (0, 1]$. This deformation is of interest and quite fruitful.

(I) The $\{qs^2, s^2\}$ -Gaussian (field) operator is equal to the $(q, s)_{BY}$ -Gaussian (field) operator. The $\{q, s^2\}$ -deformation is different from the $(q, s)_{BY}$ except for $q = 0$ or $s = 1$.

(II) In addition, the probability density for $(q, s)_{BY}$ case is known for the following three cases: (1) $s = 1, q \in (-1, 1)$ in [7][8], (2) $s \in (0, 1], q = 0$ in [5][13], and (3) $s = \sqrt{|q|}, |q| \in (0, 1)$.

The case (1) is obvious at this time and provides the (Roger's continuous) q -Hermite polynomials. Therefore, one can obtain the q -Gaussian operator ([7][8]).

In case (2), it is known that the $\{0, t\}$ -Hermite polynomials are the t -Chebyshev II polynomials ($q = 0 < t \leq 1$ and set $t = s^2$). The $\{0, 1\}$ -Gaussian measure is the semicircular measure. If $t \neq 1$, the $\{0, t\}$ -Gaussian measure is known to be a discrete probability measure with atoms at which are represented by the zeros of the t -Airy function (See [5] and references cited therein). The $\{0, t\}$ -Gaussian (field) operator is the same as the $(0, \sqrt{t})_{BY}$ -Gaussian (field) operator, which is nothing but the s -free Gaussian (field) operator [13]. Moreover, the limiting case $s \rightarrow 0$ implies the Boolean Gauss (field) operator, whose distribution is $\frac{1}{2}(\delta_1 + \delta_{-1})$. The case (3) is not referred as a particular example in [5][11]. One can see that the $\{q^2, |q|\}$ -Hermite polynomials are identified as a rescaled version of discrete $|q|$ -Hermite I polynomials¹. Let μ_q denote the orthogonalizing measure² for the discrete q -Hermite I polynomials. Correspondingly, the rescaled orthogonalizing measure of $\mu_{|q|}$ is given by $D_{1/\sqrt{1-|q|}}\mu_{|q|}, |q| \in (0, 1)$, where D_λ denotes the dilation of a probability measure μ by $D_\lambda\mu(\cdot) = \mu(\cdot/\lambda), \lambda \neq 0$. Moreover, the $\{q^2, |q|\}$ -Gaussian (field) operator coincides with the $(q, \sqrt{|q|})_{BY}$ -Gaussian (field) operator.

(III) Furthermore, since the $(q, s)_{BY}$ -Fock space as $q \rightarrow 1$ coincides with the Q^N -deformation of the Boson Fock space mentioned in (2) of Example 2.1, a limiting case of the $\{qs^2, s^2\}$ -Gaussian (field) operator as $q \rightarrow 1$ agrees with the Q^N -deformation of the classical Gaussian (field) operator [1]. It is our paper [4] which first points out this nontrivial relationship of interest.

Acknowledgment. This work was supported by the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University.

¹It is known that the discrete q -Hermite I polynomials are a symmetric case of Al-Salam-Carlitz I polynomials. In addition, the discrete q -Hermite I polynomials belong to the class IV of Brenke-Chihara polynomials. See [2][4] and references therein.

² μ_q is expressed as an infinite sum of atoms on $\{0, \pm q^k : k = 0, 1, \dots\}$.

References

- [1] Accardi, L. and Lu, Y.-G.: The qq -bit (III) : Symmetric q -Jordan-Wigner embeddings, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, **22**, (2019), 1850023 (19 pages).
- [2] Asai, N., Kubo, I. and Kuo, H.-H.: The Brenke type generating functions and explicit forms of MRM-triples by means of q -hypergeometric series, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, **16**, (2013), 1350010 (27 pages).
- [3] Asai, N. and Yoshida, H.: Poisson type operators on the Fock space of type B, *J. Math. Phys.*, **60**, (2019), 011702 (9 pages).
- [4] Asai, N. and Yoshida, H.: Deformed Gaussian operators on weighted q -Fock spaces, *J. Stoch. Anal.*, **1** (4), (2020), Article 6.
- [5] Blitvić, N.: The (q, t) -Gaussian process. *J. Funct. Anal.*, **263**, (2012), 3270–3305.
- [6] Bożejko, M., Ejsmont, W. and Hasebe, T.: Fock space associated with Coxeter groups of type B, *J. Funct. Anal.*, **269**, (2015), 1769–1795.
- [7] Bożejko, M., Kümmerer, B. and Speicher, R.: q -Gaussian processes: Non-Commutative and classical aspects, *Comm. Math. Phys.*, **185**, (1997), 129–154.
- [8] Bożejko, M. and Speicher, R.: An example of a generalized Brownian motion, *Comm. Math. Phys.*, **137**, (1991), 519–531.
- [9] Bożejko, M. and Wysoczański, J.: New examples of convolutions and non-commutative central limit theorems, *Banach Center Publ.*, **43**, (1998), 95–103.
- [10] Bożejko, M. and Wysoczański, J.: Remarks on t -transformations of measures and convolutions, *Ann. Inst. H. Poincaré Probab. Statist.*, **37**, (2001), 737–76.
- [11] Bożejko, M. and Yoshida, H.: Generalized q -deformed Gaussian random variables, *Banach Center Publ.*, **73**, (2006), 127–140.
- [12] Wojakowski, Ł.: Probability interpolating between free and boolean, Ph.D. dissertation, University of Wrocław, 2004.
- [13] Yoshida, H.: Remarks on the s -free convolution, in: *Non-commutativity, Infinite Dimensionality, and Probability at the Crossroads*, N. Obata et al. (eds.), World scientific, Singapore, 2002, 412–433.