

SELECTIVE INDEPENDENCE AND h -PERFECT TREE FORCING NOTIONS

COREY BACAL SWITZER

ABSTRACT. Generalizing the proof for Sacks forcing, we show that the h -perfect tree forcing notions introduced by Goldstern, Judah and Shelah preserve selective independent families even when iterated. As a result we obtain new proofs of the consistency of $\mathfrak{i} = \mathfrak{u} < \text{non}(\mathcal{N}) = \text{cof}(\mathcal{N})$ and $\mathfrak{i} < \mathfrak{u} = \text{non}(\mathcal{N}) = \text{cof}(\mathcal{N})$ as well as some related results.

1. INTRODUCTION

A family $\mathcal{I} \subseteq [\omega]^\omega$ is said to be *independent* if for all finite subsets $X_0, \dots, X_{n-1} \in \mathcal{I}$ and all $g : n \rightarrow 2$ we have that $X_0^{g(0)} \cap \dots \cap X_{n-1}^{g(n-1)}$ is infinite where $X_i^{g(i)}$ means X_i if $g(i) = 1$ and $\omega \setminus X_i$ if $g(i) = 0$. Such a set is said to be *maximal* if it is not properly contained in any other independent set. The *independence number* \mathfrak{i} is the least size of a maximal independent family. Despite being one of the classical cardinal characteristics, \mathfrak{i} is notoriously difficult to manipulate. Indeed many relatively simple open questions remain surrounding \mathfrak{i} , most notably the consistency of $\mathfrak{i} < \mathfrak{a}$ where \mathfrak{a} is the almost disjointness number¹. Part of the issue is that \mathfrak{i} has no known upper bound, besides the trivial 2^{\aleph_0} while it has several lower bounds thus preserving \mathfrak{i} small requires preserving the smallness of several other cardinal characteristics simultaneously.

One of the first breakthroughs in studying \mathfrak{i} came in [14] where the consistency of $\mathfrak{i} < \mathfrak{u}$ was established². There, a special independent family, now known as a *selective* independent family was constructed under CH and it was shown that under somewhat delicate conditions such a family's maximality could be preserved over a countable support iteration of length ω_2 of certain proper forcing notions. Since then selective independence has become one of the main tools in providing models of $\mathfrak{i} < 2^{\aleph_0}$ with interesting properties. See e.g [3, 5, 6]. In particular selective independent families are Sacks indestructible.

All the published examples in the literature³ of countable support iterations of proper forcing notions which are shown to preserve selective independent families

2000 *Mathematics Subject Classification.* 03E17, 03E35, 03E50.

Acknowledgements: The author would like to thank the Austrian Science Fund (FWF) for the generous support through grant number Y1012-N35. This paper was written for the RIMS Kôkyûroku volume on RIMS set theory workshop 2021. The author thanks the organizers of that conference.

¹See the appendix of [6] for a discussion of this problem.

²The cardinal \mathfrak{u} , the *ultrafilter number* is the least size of an ultrafilter base on ω .

³At least all examples the author is aware of.

are such that the iterands all have the Sacks property. The Sacks property is used throughout these proofs⁴. However the Sacks property is not needed and is overkill. In this article we show that the h -perfect tree forcing notions⁵ introduced by Goldstern, Judah and Shelah in [7] also preserve selective independence, though they may, and often do depending on h , fail to have the Sacks property.

Main Theorem 1.1 (CH). (1) *Let $h : \omega \rightarrow \omega$ be any function so that for all $n < \omega$ we have $1 < h(n) < \omega$ then the h -perfect tree forcing, \mathbb{P}_h preserves any ground model selective independent family.*

(2) *Let δ be an ordinal and $\langle \mathbb{P}_\alpha, \dot{Q}_\alpha \mid \alpha < \delta \rangle$ be a countable support iteration so that for all $\alpha < \delta$ we have*

\Vdash_α “ \dot{Q}_α is the h -perfect tree forcing for some $h \in \omega^\omega$ with $1 < h(n) < \omega$ for all $n < \omega$ ”.

Then \mathbb{P}_δ preserves all ground model selective independent families.

This allows us to show that in the models obtained by iterating such forcing notions there is a selective independent family of size \aleph_1 and, in particular, $\mathfrak{i} = \aleph_1$. As a result we obtain the following consistency results.

Main Theorem 1.2. *The following are consistent.*

- (1) $\mathfrak{i} = \mathfrak{u} < \text{non}(\mathcal{N}) = \text{cof}(\mathcal{N}) = 2^{\aleph_0}$
- (2) $\mathfrak{i} < \mathfrak{u} = \text{non}(\mathcal{N}) = \text{cof}(\mathcal{N}) = 2^{\aleph_0}$

Finally riffing off work of Brendle, Fischer and Khomskii [3], Schilhan [13] and Bergfalk, Fischer and the author [9] we can obtain that the cardinal characteristic inequalities above are consistent with Π_1^1 -definable witnesses and a Δ_3^1 -definable well order of the reals.

Main Theorem 1.3. *The cardinal characteristic inequalities featured in Main Theorem 1.2 are consistent with a Δ_3^1 well-order of the reals, a Π_1^1 witness to $\mathfrak{i} = \aleph_1$ and, in the case of the first inequality, a Π_1^1 witness to $\mathfrak{u} = \aleph_1$.*

The rest of this paper is organized as follows. In the next section we review the basics of maximal independent families and selective independence. In the following section we introduce the h -perfect tree forcing of [7] and prove Main Theorem 1.1. In Section 4 we move on to applications and prove in particular Main Theorems 1.2 and 1.3. We also discuss the relation between independent families and strong measure zero sets. Section 5 concludes with a discussion and some open questions. Throughout most of our terminology is standard and conforms e.g. to that of [11] and [10]. For combinatorial cardinal characteristics of the continuum we follow [2].

Let us finally stress that most of the results in this paper, in particular Main Theorem 1.1 are probably not new and indeed were suggested to the author by

⁴Though note that the Sacks property is not enough to ensure that selective independence is preserved: Silver forcing has the Sacks property but will kill the maximality of any ground model independent family and, if iterated ω_2 many times will result in a model of $\mathfrak{i} = 2^{\aleph_0} > \aleph_1$.

⁵See Definition 3.1 for the definition of these posets.

both Jörg Brendle and Vera Fischer⁶. However, they do not seem to ever have been written down, at least not explicitly and this seemed worth while to do. In particular, while the consistency of $\mathfrak{i} < \text{non}(\mathcal{N})$ was shown in [4, Theorem 3.8], the proof uses a short finite support iteration of ccc forcing notions over a model of MA (a “dual iteration”) and hence is very different than the model constructed here. Moreover in the model in [4] we do not know the value of \mathfrak{u} .

Acknowledgments. The author thanks Jörg Brendle for pointing out [7] to him and suggesting that h -perfect forcing may preserve selective independent families. The author thanks Vera Fischer for many very helpful conversations on this material and sharing her wealth of knowledge on selective independence and the cardinal \mathfrak{i} .

2. SELECTIVE INDEPENDENCE

In this section we introduce the notion of a *selective independent family*. The reader familiar with this idea, for example as presented in [3] or [5], can comfortably skip this section as nothing is new. Selective independent families were introduced implicitly in Shelah’s proof of the consistency of $\mathfrak{i} < \mathfrak{u}$ in [14]. To facilitate the discussion we utilize the following notation.

Notation 1. For $\mathcal{I} \subseteq [\omega]^\omega$,

- let $\text{FF}(\mathcal{I})$ denote the set of finite partial functions g from \mathcal{I} to $\{0, 1\}$, and
- for $g \in \text{FF}(\mathcal{I})$ write \mathcal{I}^g for

$$\bigcap \{A \mid A \in \text{dom}(g) \text{ and } g(A) = 1\} \cap \bigcap \{\omega \setminus A \mid A \in \text{dom}(g) \text{ and } g(A) = 0\}.$$

In this notation, a family $\mathcal{I} \subseteq [\omega]^\omega$ is *independent* if \mathcal{I}^g is infinite for all $g \in \text{FF}(\mathcal{I})$. An independent family \mathcal{I} is *maximal* if

$$\forall X \in [\omega]^\omega \exists g \in \text{FF}(\mathcal{I}) \text{ such that } \mathcal{I}^g \cap X \text{ or } \mathcal{I}^g \setminus X \text{ is finite,}$$

We will need a slight strengthening of maximality.

Definition 2.1. An independent family \mathcal{I} is *densely maximal* if

$$\forall X \in [\omega]^\omega \text{ and } g' \in \text{FF}(\mathcal{I}) \exists g \supseteq g' \text{ in } \text{FF}(\mathcal{I}) \text{ such that } \mathcal{I}^g \cap X \text{ or } \mathcal{I}^g \setminus X \text{ is finite.}$$

In other words, an independent family \mathcal{I} is *densely maximal* if for each $X \in [\omega]^\omega$ the collection of g ’s witnessing that $\mathcal{I} \cup \{X\}$ is not a larger independent family is dense in $(\text{FF}(\mathcal{I}), \supseteq)$.

Definition 2.2. Let \mathcal{I} be an independent family. The *density ideal* of \mathcal{I} , denoted $\text{id}(\mathcal{I})$, is

$$\{X \subseteq \omega \mid \forall g' \in \text{FF}(\mathcal{I}) \exists g \supseteq g' \text{ in } \text{FF}(\mathcal{I}) \text{ such that } \mathcal{I}^g \cap X \text{ is finite}\}.$$

Dual to the density ideal of \mathcal{I} is the *density filter* of \mathcal{I} , denoted $\text{fil}(\mathcal{I})$ and defined as

$$\{X \subseteq \omega \mid \forall g' \in \text{FF}(\mathcal{I}) \exists g \supseteq g' \text{ in } \text{FF}(\mathcal{I}) \text{ such that } \mathcal{I}^g \setminus X \text{ is finite}\}.$$

⁶Private communication.

Observe also that for an infinite independent family \mathcal{I} , none of the above definitions' meanings change if we replace the word "finite" with "empty". We have as well the following from [9, Lemma 5.4].

Lemma 2.3. *A family $\mathcal{I} \subseteq [\omega]^\omega$ is densely maximal if and only if*

$$P(\omega) = \text{fil}(\mathcal{I}) \cup \langle \omega \setminus \mathcal{I}^g \mid g \in \text{FF}(\mathcal{I}) \rangle_{\text{dn}}.$$

Where for a set $\mathcal{X} \subseteq [\omega]^\omega$ the set $\langle \mathcal{X} \rangle_{\text{dn}}$ denotes the downward closure of \mathcal{X} under \subseteq^* i.e. $A \in \langle \mathcal{X} \rangle_{\text{dn}}$ if and only if there is an $X \in \mathcal{X}$ with $A \subseteq^* X$. Later on we will similarly denote $\langle \mathcal{X} \rangle_{\text{up}}$ for the upward closure of \mathcal{X} under \subseteq^* .

The following are easily verified, see [9, Lemma 5.5].

Lemma 2.4. (1) *If \mathcal{I}' is an independent family and $\mathcal{I} \subseteq \mathcal{I}'$ then $\text{fil}(\mathcal{I}) \subseteq \text{fil}(\mathcal{I}')$;*
 (2) *if κ is a regular uncountable cardinal and $\langle \mathcal{I}_\alpha \mid \alpha < \kappa \rangle$ is a continuous increasing chain of independent families then $\text{fil}(\bigcup_{\alpha < \kappa} \mathcal{I}_\alpha) = \bigcup_{\alpha < \kappa} \text{fil}(\mathcal{I}_\alpha)$;*
 (3) *If \mathcal{I} is an independent family then $\text{fil}(\mathcal{I}) = \bigcup \{ \text{fil}(\mathcal{J}) \mid \mathcal{J} \in [\mathcal{I}]^{\leq \omega} \}$.*

Recall that given a family \mathcal{F} of subsets of ω we say that

- (1) \mathcal{F} is a *P-set* if every countable family $\{A_n \mid n < \omega\} \subseteq \mathcal{F}$ has a pseudointersection $B \in \mathcal{F}$,
- (2) \mathcal{F} is a *Q-set* if given every partition of ω into finite sets $\{I_n \mid n < \omega\}$ there is a *semiselect* $A \in \mathcal{F}$ i.e. $|A \cap I_n| \leq 1$ for all $n < \omega$,
- (3) \mathcal{F} is *Ramsey* if it is both a *P-set* and a *Q-set*.

If \mathcal{F} is a filter and a *P-set* (respectively a *Q-set*, *Ramsey set*) we call \mathcal{F} a *P-filter* (respectively a *Q-filter*, *Ramsey filter*).

Definition 2.5. An independent family \mathcal{I} is *selective* if it is densely maximal and $\text{fil}(\mathcal{I})$ is *Ramsey*.

Fact 2.6 (Shelah, see [14]). *CH implies the existence of a selective independent family.*

Under certain circumstances countable support iterations of proper forcing notions over a model of CH will preserve that a given ground model selective independent family is maximal. Towards clarifying the meaning of "certain conditions" recall the following preservation result, due to Shelah, see [14, Lemma 3.2].

Theorem 2.7. *Assume CH. Let δ be a limit ordinal and let $\langle \mathbb{P}_\alpha, \dot{Q}_\alpha \mid \alpha < \delta \rangle$ be a countable support iteration of ${}^\omega\omega$ -bounding proper posets. Let $\mathcal{F} \subseteq P(\omega)$ be a Ramsey set and let \mathcal{H} be a subset of $P(\omega) \setminus \langle \mathcal{F} \rangle_{\text{up}}$. If $V^{\mathbb{P}_\alpha} \models P(\omega) = \langle \mathcal{F} \rangle_{\text{up}} \cup \langle \mathcal{H} \rangle_{\text{dn}}$ for all $\alpha < \delta$ then $V^{\mathbb{P}_\delta} \models P(\omega) = \langle \mathcal{F} \rangle_{\text{up}} \cup \langle \mathcal{H} \rangle_{\text{dn}}$ as well.*

We will also need the notion of *Cohen preserving*.

Definition 2.8. Let \mathbb{P} be a forcing notion. We say that \mathbb{P} is *Cohen preserving* if every every new dense open subset of $2^{<\omega}$ (or, equivalently $\omega^{<\omega}$, ...) contains an old dense subset. More formally, \mathbb{P} is *Cohen preserving* if for all $p \in \mathbb{P}$ and all \mathbb{P} -names \dot{D} so that $p \Vdash \dot{D} \subseteq 2^{<\omega}$ is dense open" there is a dense $E \subseteq 2^{<\omega}$ in the ground model and a $q \leq_{\mathbb{P}} p$ so that $q \Vdash \dot{E} \subseteq \dot{D}$.

Being Cohen preserving is preserved by countable support iterations of proper forcing notions.

Theorem 2.9 (Shelah, See Conclusion 2.15D, pg. 305 of [15], see also [5], Theorem 27). *If δ is an ordinal and $\langle \mathbb{Q}_\alpha, \dot{\mathbb{R}}_\alpha \mid \alpha < \delta \rangle$ is a countable support iteration of forcing notions so that for each $\alpha < \delta$ we have \Vdash_α “ $\dot{\mathbb{R}}_\alpha$ is proper and Cohen preserving” then \mathbb{Q}_δ is proper and Cohen preserving.*

3. h -PERFECT TREES PRESERVE SELECTIVE INDEPENDENT FAMILIES

In this section we prove Main Theorem 1.1. First, recall the definition of h -perfect tree forcing from [7].

Definition 3.1. Given a function $h : \omega \rightarrow \omega$ with $1 < h(n) < \omega$ for all $n < \omega$, the h -perfect tree forcing, denoted \mathbb{PT}_h , is the forcing notion consisting of trees $p \subseteq \omega^{<\omega}$ so that the following hold:

- (1) For all $t \in p$ and all $l \in \text{dom}(t)$ we have $t(l) < h(l)$.
- (2) Every $t \in p$ has either one or $h(l(t))$ -many immediate successors in T .
- (3) For every $t \in p$ there is a $t' \supseteq t$ with $t' \in p$ and there are $h(l(t'))$ many immediate successors of t' in p .

The order is inclusion.

Note that the case where $h(n) = 2$ for all $n < \omega$ is simply Sacks forcing, while for a fast growing $h : \omega \rightarrow \omega$ this forcing will make the ground model reals measure zero so which h we choose can affect the properties of the forcing significantly. This forcing notion was first considered in [7]. In [7] the following is shown.

Fact 3.2 ([7]). *For any $h : \omega \rightarrow \omega$ with $1 < h(n) < \omega$ for all $n < \omega$ the following hold.*

- (1) \mathbb{PT}_h is proper, and in fact satisfies Axiom A.
- (2) \mathbb{PT}_h is ω^ω -bounding.
- (3) \mathbb{PT}_h preserves P -points.

For the rest of this section fix an arbitrary function $h : \omega \rightarrow \omega$ so that $1 < h(n) < \omega$ for all $n < \omega$. We will prove first that forcing with \mathbb{PT}_h preserves a ground model selective independent family. Then we will show that under CH arbitrary countable support iterations of h -perfect tree forcings (where the h can change and need not even be in the ground model) will preserve selective independent families. First we introduce some arboreal terminology. If $p \in \mathbb{PT}_h$ and $n < \omega$ then a node $t \in p$ is an n^{th} -splitting node if it has $h(l(t))$ many immediate successors and it has the $n - 1$ predecessors with this property. Denote by $\text{Split}_n(p)$ the set of n -splitting nodes. We say that for two h -perfect trees $p, q \in \mathbb{PT}_h$ that $q \leq_n p$ if $q \leq p$ and for all $i < n + 1$ $\text{Split}_i(p) = \text{Split}_i(q)$. Given any $p \in \mathbb{PT}_h$ and any node $t \in p$ we let p_t denote the tree $\{s \in p \mid s \subseteq t \text{ or } t \subseteq s\}$. Note that $p_t \in \mathbb{PT}_h$ and $p_t \leq p$ for any $t \in p$.

Lemma 3.3. *Let $p \in \mathbb{P}\mathbb{T}_h$ and let \dot{X} be a $\mathbb{P}\mathbb{T}_h$ -name for an infinite subset of ω . There is a $q \leq p$ so that for all $n < \omega$ and any n -splitting node $t \in q$ we have that q_t decides $\dot{X} \cap \check{n}$.*

Such a q is called *preprocessed* for \dot{X} .

Proof. This is a standard fusion argument but we sketch it for completeness. Fix p and \dot{X} as in the lemma. Inductively define a fusion sequence $\dots \leq_n p_n \leq_{n-1} p_{n-1} \leq_{n-2} \dots \leq_1 p_1 \leq_0 p_0 = p$ as follows. Given $k < \omega$ and p_k , for each k^{th} splitting node $t \in \text{Split}_k(p_k)$ let $p'_t \leq (p_k)_t$ decide $\dot{X} \cap \check{k}$. Let $p_{k+1} = \bigcup_{t \in \text{Split}_k(p_k)} p'_t$. Clearly $p_{k+1} \leq_k p_k$ and the fusion $q := \bigcap_{k < \omega} p_k$ is as needed. \square

We now move to the first substantial lemma.

Lemma 3.4. *$\mathbb{P}\mathbb{T}_h$ is Cohen preserving.*

Proof. Fix $p \in \mathbb{P}\mathbb{T}_h$ and suppose that \dot{D} is a $\mathbb{P}\mathbb{T}_h$ -name for a dense open subset of $2^{<\omega}$. Enumerate $2^{<\omega}$ as $\{s_n \mid n < \omega\}$. We will inductively construct sequences $\{p_n \mid n < \omega\}$ and $\{t_n \mid n < \omega\}$ so that the following hold.

- (1) $p_0 = p$.
- (2) $p_{n+1} \leq_n p_n$ for all $n < \omega$.
- (3) $t_n \supseteq s_n$ for all $n < \omega$.
- (4) $p_{n+1} \Vdash \check{t}_n \in \dot{D}$.

Given such sequences, let $q = \bigcap_{n < \omega} p_n$ and $E = \{t_n \mid n < \omega\}$. Clearly $q \Vdash \check{E} \subseteq \dot{D}$ and E is dense so, assuming we can construct such sequences we will be done. This is done by induction. Given p_k , enumerate $\text{Split}_k(p_k)$ as $\{u_i \mid i < l\}$. Note that $l \in \omega$ depends not just on h but also on p_k but what matters here is that it is finite (which it is). Now let $p'_{k,0} \leq (p_k)_{u_0}$ decide some $t_k^0 \supseteq s_k$ to be in \dot{D} (since \dot{D} is forced to be dense this is possible). Next, let $p'_{k,1} \leq (p_k)_{u_1}$ decide some $t_k^1 \supseteq t_k^0$ to be in \dot{D} . Continuing this way, inductively, let for all $0 < i < l$ let $p'_{k,i} \leq (p_k)_{u_i}$ decide some $t_k^i \supseteq t_k^{i-1}$ to be in \dot{D} . Let $p_{k+1} = \bigcup_{i < l} p'_{k,i}$ and t_{k+1} be t_k^{l-1} . Since \dot{D} is forced to be open we have that $p_{k+1} \Vdash \check{t}_{k+1} \in \dot{D}$ so we're done. \square

Fix a selective independent family \mathcal{I} in the ground model.

Lemma 3.5. *If $G \subseteq \mathbb{P}\mathbb{T}_h$ is generic over V then in $V[G]$ the ideal $\text{id}(\mathcal{I})$ is generated by $\text{id}(\mathcal{I}) \cap V$.*

Note that dually this lemma implies that in $V[G]$ the filter $\text{fil}(\mathcal{I})$ is generated by $\text{fil}(\mathcal{I}) \cap V$.

Proof. Let $p \in \mathbb{P}\mathbb{T}_h$, let \dot{X} be a $\mathbb{P}\mathbb{T}_h$ -name for an infinite subset of ω and suppose that $p \Vdash \dot{X} \in \text{id}(\mathcal{I})$. We need to find a ground model $Y \in [\omega]^\omega$ and an $r \leq p$ so that $r \Vdash \dot{X} \subseteq \check{Y}$. Towards this, via a fusion argument, or just using the properness of $\mathbb{P}\mathbb{T}_h$, find a $q \leq p$ and a countable $\mathcal{J} \subseteq \mathcal{I}$ so that $q \Vdash \dot{X} \in \text{id}(\mathcal{J})$. Since \mathcal{J} is countable, we can associate $\text{FF}(\mathcal{J})$ with $2^{<\omega}$. Let \dot{D} be the $\mathbb{P}\mathbb{T}_h$ -name for the dense

open subset of $2^{<\omega}$ defined by $q \Vdash \check{g} \in \dot{D}$ if and only if $\mathcal{J}^g \cap \dot{X} = \emptyset$. Since $\mathbb{P}\mathbb{T}_h$ is Cohen preserving there is an $r \leq q$ and a dense $E \subseteq 2^{<\omega}$ so that $r \Vdash \check{E} \subseteq \dot{D}$. Let $Y = \bigcap_{h \in E} (\omega \setminus \mathcal{J}^h)$. Observe that $Y \in \text{id}(\mathcal{J})$ and hence $Y \in \text{id}(\mathcal{I})$. To see this, let $g' \in \text{FF}(\mathcal{J})$ be arbitrary and let $g \supseteq g'$ be in E . We have that $Y \subseteq \omega \setminus \mathcal{J}^g$ and hence $Y \cap \mathcal{J}^g = \emptyset$ which by definition means that Y is in the ideal. The following claim now completes the proof.

Claim 3.6. $r \Vdash \dot{X} \subseteq \check{Y}$.

Proof. Let $r \in G$ be $\mathbb{P}\mathbb{T}_h$ -generic over V . Note that by the way \dot{D}^G is defined we have that $\dot{X} = \bigcap_{g \in \dot{D}^G} (\omega \setminus \mathcal{B}^g)$ and since $E \subseteq \dot{D}^G$ we're done. \square

\square

Theorem 3.7 (CH). *If \mathcal{I} is a selective independent family and $G \subseteq \mathbb{P}\mathbb{T}_h$ is generic over V then $V[G] \models \text{“}\mathcal{I} \text{ is a selective independent family”}$.*

Proof. There are three things to check. We need to show 1) that $\text{fil}(\mathcal{I})$ is a P -filter, 2) that $\text{fil}(\mathcal{I})$ is a Q -filter and 3) that \mathcal{I} is densely maximal. We take these one at a time. First though, since we assume CH in V , note that the fact that $\text{fil}(\mathcal{I})$ is a P -set implies that we can assume that $\text{fil}(\mathcal{I})$ is generated by an ω_1 length \subseteq^* -descending sequence $\{B_\alpha \mid \alpha < \omega_1\}$. In other words for all $\alpha < \beta < \omega_1$ we have $B_\beta \subseteq^* B_\alpha$ and every $A \in \text{fil}(\mathcal{I})$ is almost contained in some (equivalently a tail of) B_γ . Fix such a sequence $\{B_\alpha \mid \alpha < \omega_1\}$.

To see that $\text{fil}(\mathcal{I})$ is a P -filter in $V[G]$ then, note that if $\{A_n \mid n < \omega\} \subseteq \text{fil}(\mathcal{I})$ in $V[G]$ by Lemma 3.5 there are countable ordinals $\{\gamma_n \mid n < \omega\}$ so that for all $n < \omega$ we have $B_{\gamma_n} \subseteq^* A_n$. Let $\gamma \geq \sup_{n < \omega} \gamma_n$. We have $B_\gamma \subseteq^* A_n$ for all $n < \omega$ so $\text{fil}(\mathcal{I})$ is a P -filter in $V[G]$.

The fact that $\text{fil}(\mathcal{I})$ is a Q -filter still in $V[G]$ follows immediately from the fact that $\mathbb{P}\mathbb{T}_h$ is ω^ω -bounding.

Thus it remains to see that \mathcal{I} remains densely maximal in $V[G]$. Suppose not and let \dot{X} be a $\mathbb{P}\mathbb{T}_h$ -name for an infinite subset of ω so that in $V[G]$ \dot{X}^G is not in $\text{fil}(\mathcal{I})$ and for all $g \in \text{FF}(\mathcal{I})$ we have $\dot{X} \not\subseteq \omega \setminus \mathcal{I}^g$. Let $p \in G$ force this. Without loss of generality we may assume that p is preprocessed for \dot{X} i.e. for each $n < \omega$ and every $t \in \text{Split}_n(p)$ we have that p_t decides $\dot{X} \cap \check{n}$. For each split node t of p let $Y_t = \{n \mid p_t \Vdash \check{n} \notin \dot{X}\}$. Note that for all split nodes t we have that $p_t \Vdash \dot{X} \subseteq \check{Y}_t$.

Claim 3.8. *For all split nodes t of p we have $Y_t \in \text{fil}(\mathcal{I})$.*

Proof. Otherwise there is a $t \in \text{Split}(p)$ so that $Y_t \subseteq \omega \setminus \mathcal{I}^g$ for some $g \in \text{FF}(\mathcal{I})$ but since $p_t \Vdash \dot{X} \subseteq \check{Y}_t$ we have that $p_t \Vdash \dot{X} \subseteq \omega \setminus \mathcal{I}^g$. However this contradicts the choice of p . \square

Since $\text{fil}(\mathcal{I})$ is a P -filter generated by ground model elements there is a $C \in \text{fil}(\mathcal{I}) \cap V$ so that $C \subseteq^* Y_t$ for all $t \in \text{Split}(p)$. Let $f \in \omega^\omega$ be a strictly increasing function such that for all $n < \omega$ we have $C \setminus f(n) \subseteq \bigcap \{Y_t \mid t \in \text{Split}_j(p), j \leq n+2\}$. The following is proved in [14], as well as [6, Lemma 3.15] but we include it for completeness.

Claim 3.9. *There is a $C^* \subseteq C$ so that $C^* \in \text{fil}(\mathcal{I}) \cap V$ and, letting $\{k_n \mid n < \omega\}$ be a strictly increasing enumerate of C^* , we have that $f(k_n) < k_{n+1}$ for all $n < \omega$ and $f(1) < k_1$.*

Proof. This follows from the fact that $\text{fil}(\mathcal{I})$ is a Q -filter⁷. Inductively find a sequence $\{n_l\}_{l \in \omega}$ so that $n_0 = 0$ and

$$n_{l+1} = \min\{n \mid n_l < n \text{ and for all } m < n_l, f(m) < n\}$$

. Consider the interval partition $\mathcal{E}_0 = \{\{n_{3l}, n_{3l+3}\}\}_{l \in \omega}$. Since $\text{fil}(\mathcal{I})$ is a Q -set and a filter there is a $C_1 \subseteq C$ so that for all $l < \omega$ we have $|C_1 \cap [n_{3l}, n_{3l+3}]| \leq 1$. Now define an equivalence relation \mathcal{E}_1 on ω by

$$m \equiv_{\mathcal{E}_1} k \text{ iff } m = k \vee m, k \in C_1 \wedge (m < k \leq f(m) \vee k < m \leq f(k)).$$

In words, this says that every element of $\omega \setminus C_1$ is in their own equivalence class and distinct elements $m, k \in C_1$ are \mathcal{E}_1 -equivalent just in case applying f to the smaller one is greater or equal to the bigger one. Every \mathcal{E}_1 equivalence class has at most two members. To see this, suppose $m_1 < m_2 < m_3 \in C_1$ were all in the same equivalence class. By definition of \mathcal{E}_1 we have $m_1 < m_2 < m_3 \leq f(m_1)$. However, since C_1 is a semiselector for the interval partition \mathcal{E}_0 there are distinct $l_1 < l_2 < l_3$ so that for all $i \in \{1, 2, 3\}$ we have $m_i \in [n_{3l_i}, n_{3l_i+1})$. Thus we get $m_1 < n_{3l_2} \leq m_2 < n_{3l_3} \leq m_3 \leq f(m_1)$ but by the definition of the n_l sequence we also have $f(m_1) \leq n_{3l_2+1} < n_{3l_3}$ which is a contradiction.

Now let $C_2 \subseteq C_1$ be a semiselector for \mathcal{E}_1 in $\text{fil}(\mathcal{I})$. Without loss of generality $0 \in C_2$. Let $\{k_n\}$ be an increasing enumeration of C_2 . For all $n < n'$ we have that n and n' are not in the same \mathcal{E}_1 equivalence class and therefore $f(n) < n'$. As such $C_2 = C^*$ is as needed.

For the final point note that $\text{fil}(\mathcal{I})$ is closed under finite changes to elements so we can augment C^* to get $f(1) < k_1$ as needed. \square

We will find a $q \leq p$ forcing that $C^* \subseteq \dot{X}$ which contradicts the choice of X . Obviously this will be a fusion argument. Let $t^* = \text{Stem}(p)$ and let $p_0 = p = p_{t^*}$. Now for each $i \in h(l(t^*))$ let $w(t^*, i) \in p_0$ be a 1-splitting node extending $(t^*) \frown i$. Since $k_1 > f(1)$ we have that $k_1 \in \bigcap \{Y_{w(t^*, i)} \mid i \in h(l(t^*))\}$. This means that for each $i \in h(l(t^*))$ there is a $w'(t^*, i) \in \text{Split}_{k_1+1}(p_0)$ extending $w(t^*, i)$ which forces that $k_1 \in \dot{X}$ since p_0 is preprocessed. Let $p_1 = \bigcup_{i \in h(l(t^*))} p_{w'(t^*, i)}$. Note that $p_1 \leq_0 p_0$ and forces that $k_1 \in \dot{X}$. Also note that $\text{Split}_1(p_1) \subseteq \text{Split}_{k_1+1}(p_0)$ and, by construction, for each m we have that $\text{Split}_m(p_1) \subseteq \text{Split}_{k_1+m}(p_0)$.

Now proceed inductively defining p_{n+1} as follows. Assume p_n has been defined and that for all $m < \omega$ we have $\text{Split}_{n+m}(p_n) \subseteq \text{Split}_{k_n+m}(p_0)$. Observe that we have $k_{n+1} \in \bigcap \{Y_t \mid t \in \text{Split}_n(p_n)\}$ since $k_{n+1} > f(k_n)$ and hence we can find for each $t \in \text{Split}_n(p_n)$ and each $i \in h(l(t))$ a $w(t, i) \in \text{Split}_{k_{n+1}+1}(p_0)$ in p_n which contains

⁷In fact a family $\mathcal{F} \subseteq [\omega]^\omega$ is a Q -filter if and only if for each increasing $f \in \omega^\omega$ there is a $C^* = \{k_n \mid n < \omega\} \in \mathcal{F}$ such that $f(k_n) < k_{n+1}$, see [6, Lemma 3.15].

$t \frown i$ so that $(p_n)_{w(t,i)} \Vdash \check{k}_{n+1} \in \dot{X}$. Let $p_{n+1} = \bigcup_{t \in \text{Split}_n(p_n)} \bigcup_{i \in h(l(t))} (p_n)_{w(t,i)}$. Clearly this is as needed.

Let q be the fusion of the p_n 's. We have that $q \Vdash \check{C}^* \subseteq \dot{X}$ contradicting the fact that \dot{X} is forced not to be in $\text{fil}(\mathcal{I})$ so we're done. \square

Now we show how to lift the above proof to show that iterations of h -perfect tree forcing notions preserve selective independent families.

Theorem 3.10 (CH). *Let δ be an ordinal and \mathcal{I} be a selective independent family. Let $\langle \mathbb{P}_\alpha, \dot{\mathbb{Q}}_\alpha \mid \alpha < \delta \rangle$ be a countable support iteration of posets so that for all $\alpha < \delta$ we have $\Vdash_\alpha \dot{\mathbb{Q}}_\alpha$ is $\mathbb{P}\mathbb{T}_h$ for some $h \in \omega^\omega$ with $1 < h(n) < \omega$ for all $n < \omega$. If $G \subseteq \mathbb{P}_\delta$ is generic over V then $V[G] \models \text{“}\mathcal{I} \text{ is a selective independent family”}$.*

Proof. The proof is by induction on δ . Note first that by Theorems 2.7 and 2.9 and Lemmas 3.4 and 3.5 we have that for each $\alpha \leq \delta$ that \mathbb{P}_α is Cohen preserving and forces that $\text{fil}(\mathcal{I})$ is generated by ground model sets. This guarantees that \mathbb{P}_α forces that $\text{fil}(\mathcal{I})$ is a P -filter. Moreover being ω^ω -bounding ensures that $\text{fil}(\mathcal{I})$ is a Q -filter hence $\text{fil}(\mathcal{I})$ is forced to be Ramsey by every \mathbb{P}_α for $\alpha \leq \delta$. Therefore we just need to ensure that $\Vdash_\delta \text{“}\mathcal{I} \text{ is densely maximal”}$ under the assumption that for all $\alpha < \delta$ we have that $\Vdash_\alpha \text{“}\mathcal{I} \text{ is densely maximal”}$. To show this we will use the characterization of dense maximality given by Lemma 2.3. We now consider two separate cases:

Case 1: $\delta = \beta + 1$ for some β . The proof of this case is almost verbatim the same as the proof of theorem 3.7 noting that, by the above, we can assume that $\text{fil}(\mathcal{I})^{V^{\mathbb{P}_\beta}}$ is generated by $\text{fil}(\mathcal{I}) \cap V$ and the f and C^* found in that proof can be assumed to come from the ground model by ω^ω -boundedness.

Case 2: δ is a limit ordinal. Inductively we have that if $\beta < \delta$ and $G_\beta \subseteq \mathbb{P}_\beta$ is generic over V then

$$V[G_\beta] \models P(\omega) = \langle \text{fil}(\mathcal{I}) \cap V \rangle_{\text{up}} \cup \langle \omega \setminus \mathcal{I}^g \mid g \in \text{FF}(\mathcal{I}) \rangle_{\text{dn}}.$$

But then by Theorem 2.7 plus the fact that $\text{fil}(\mathcal{I})$ is a Ramsey filter in $V^{\mathbb{P}_\delta}$ we get

$$\Vdash_\delta P(\omega) = \langle \text{fil}(\mathcal{I}) \cap V \rangle_{\text{up}} \cup \langle \omega \setminus \mathcal{I}^g \mid g \in \text{FF}(\mathcal{I}) \rangle_{\text{dn}}.$$

Which, by Lemma 2.3 is exactly what we needed to show. \square

4. APPLICATIONS

We now turn to applications of the results from the previous section. The most obvious of these is that there is a small independent family in any model obtained by iteratively forcing with h -perfect tree partial orders. In particular we get the following as a corollary to Theorem 3.10.

Corollary 4.1. *It is consistent that $\mathfrak{i} = \aleph_1 < \text{non}(\mathcal{N}) = \aleph_2$.*

As mentioned in the introduction this consistent inequality was first shown in [4, Theorem 3.8], though by a very different construction.

In the models constructed in [7] many interesting properties hold with regards to the structure of the strong measure zero sets, for example the consistency of “the

additivity of the strong measure zero ideal is $\aleph_2 = 2^{\aleph_0}$. As a consequence all of these are also consistent with $\mathfrak{i} = \aleph_1$. One thing to note as a consequence of this is the following.

Corollary 4.2. *It is independent of ZFC whether there is a maximal independent family of strong measure zero.*

Proof. In the Laver model, [1, Model 7.6.13] every strong measure zero set is countable so no maximal independent family has strong measure zero. By contrast if the additivity of the strong measure zero ideal is \aleph_2 then in particular any set of reals of size \aleph_1 will be strong measure zero, in particular any selective independent family from the ground model. □

We can also now iteration mixings of h -perfect posets with other proper partial orders which iteratively preserve small selective independent families.

Theorem 4.3. *The following are consistent.*

- (1) $\mathfrak{i} = \mathfrak{u} < \text{non}(\mathcal{N})$
- (2) $\mathfrak{i} < \mathfrak{u} = \text{non}(\mathcal{N})$

Proof. For the first inequality, as noted above in Fact 3.2, for any h we have that $\mathbb{P}\mathbb{T}_h$ preserves P -points hence in any model constructed by iterating h -perfect tree forcings with countable support over a model of CH there will be a P -point base of size \aleph_1 . For the second inequality alternating between $\mathbb{P}\mathbb{T}_h$ for e.g. $h(n) = 2^n$ and the forcing notions \mathbb{Q}_T of [14] (alongside some bookkeeping device) will increase \mathfrak{u} but preserves selective independent families. □

Finally we note some applications to definability.

Theorem 4.4. *Both inequalities featured in Theorem 4.3 are consistent with a Π_1^1 independent family of size \aleph_1 , a Δ_3^1 well order of the reals and, in the case of the first inequality, a Π_1^1 ultrafilter base for a P -point of size \aleph_1 .*

Proof. Schilhan [13] has shown that in L there is a Π_1^1 ultrafilter base for a P -point and Brendle, Fischer and Khomskii [3] have shown that in L there is a Σ_2^1 selective independent family and that if there is a Σ_2^1 maximal independent family then there is a Π_1^1 maximal independent family. It follows that all of these objects can be preserved by the iterations described in the proof of Theorem 4.3 assuming the ground model is L . Finally for the Δ_3^1 -well order of the reals we apply the forcing from [9] noting that the main theorem of that paper is precisely that such objects can be preserved by this forcing, even when other forcing notions, such as $\mathbb{P}\mathbb{T}_h$ are added to the iteration. □

5. CONCLUSION AND OPEN QUESTIONS

The proof of Main Theorem 1.1 are almost verbatim the same as those for Sacks forcing [5], Shelah's forcing for killing a maximal ideal used in [14] and very similar

to the proof for the coding with perfect tree forcing from [9]. In particular really only structural properties of the forcing are used. This suggests there should be a general property of proper, ω^ω -bounding forcing notions which imply that small selective independent families are preserved. The following seems like the first place to go to isolate such a property.

Question 1. Suppose δ is an ordinal and $\langle \mathbb{P}_\alpha, \dot{\mathbb{Q}}_\alpha \mid \alpha < \delta \rangle$ is a countable support iteration of proper, ω^ω -bounding, Cohen preserving forcing notions. Let \mathcal{I} is a selective independent family in $V \models \text{CH}$. If for all $\alpha < \delta \Vdash_\alpha$ “ $\dot{\mathbb{Q}}_\alpha$ forces “ $\text{fil}(\mathcal{I})$ is generated by ground model sets” then does \mathbb{P}_δ preserve the maximality of \mathcal{I} ?

REFERENCES

- [1] Tomek Bartoszyński and Haim Judah. *Set Theory: On the Structure of the Real Line*. A.K. Peters, Wellsley, MA, 1995.
- [2] Andreas Blass. Combinatorial cardinal characteristics of the continuum. In Matthew Foreman and Akihiro Kanamori, editors, *Handbook of Set Theory*, pages 395–489. Springer, Dordrecht, 2010.
- [3] Jörg Brendle, Vera Fischer and Yuri Khomskii. Definable Maximal Independent Families *Proceedings of the American Mathematical Society*, 147: 3547-3557, 2019.
- [4] B. Balcar, F. Hernández-Hernández and M. Hrušák. Combinatorics of Dense Subsets of the Rationals *Fund. Math.*, 183: 59-80, 2004.
- [5] Vera Fischer and Diana C. Montoya. Ideals of Independence *Arch. for Math. Logic*, 58(5-6): 767-785, 2019.
- [6] Jorge A. Cruz-Chapital, Vera Fischer, Osvaldo Guzmán and Jaroslav Šupina. Filter Games, Compact Partitions and MAD Families *Submitted* 2021.
- [7] Martin Goldstern, Haim Judah and Saharon Shelah. Strong Measure Zero Sets without Cohen Reals *J. Symbolic Logic*, 58(4): 1323-1341, 1993.
- [8] Vera Fischer and Corey Bacal Switzer. Tight Eventually Different Families *Submitted*, 2021.
- [9] Jeffrey Bergfalk, Vera Fischer and Corey Bacal Switzer. Projective Well Orders and Coanalytic Witnesses *Submitted*, 2021.
- [10] Lorenze J. Halbeisen. *Combinatorial Set Theory with a Gentle Introduction to Forcing*. Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2017.
- [11] Thomas Jech. *Set Theory*. Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2003.
- [12] Arnold W. Miller. Infinite combinatorics and definability. *Ann. Pure Appl. Logic*, 41(2):179–203, 1989.
- [13] Johnathan Schilhan. Coanalytic ultrafilter bases. *Submitted*, 2019.
- [14] Saharon Shelah. $\text{CON}(u > i)$. *Arch. Math. Logic*, 31(6): 433-443, 1992.
- [15] Saharon Shelah. *Proper and Improper Forcing*. Perspectives in Logic, Cambridge University Press, Cambridge, 3rd Edition, 2016.

(C. B. Switzer) INSTITUT FÜR MATHEMATIK, KURT GÖDEL RESEARCH CENTER, UNIVERSITÄT WIEN, KOLINGASSE 14-16, 1090 WIEN, AUSTRIA
Email address: corey.bacal.switzer@univie.ac.at