

SPHERICAL HARMONICS AND HARDY'S INEQUALITIES

SHUJI MACHIHARA

ABSTRACT. We consider the derivative operators for radial direction and spherical direction. We also investigate the operator which takes the spherical average for functions. We reconfirm those properties with particular attention to orthogonality. As an application, the Hardy type inequality is presented with spherical derivatives in the framework of equalities. This clarifies the difference between contribution by radial and spherical derivatives in the improved Hardy inequality as well as nonexistence of nontrivial extremizers without compactness arguments.

This paper is based on the joint work with Neal Bez and Tohru Ozawa.

1. PRELIMINARIES PART 1

In this section we define and investigate some derivative operators in Euclian space. Although the facts are well-known, we here check the all lines for calclations and the orthogonality of those. We define the radial derivative and the spherical derivative

$$\begin{aligned}\partial_r &= \frac{x}{|x|} \cdot \nabla = \sum_{j=1}^n \frac{x_j}{|x|} \partial_j, \\ L &= \nabla - \frac{x}{|x|} \partial_r = \left(\partial_1 - \frac{x_1}{|x|} \partial_r, \dots, \partial_n - \frac{x_n}{|x|} \partial_r \right).\end{aligned}$$

where $\partial_j = \partial/\partial x_j, j = 1, \dots, n$. We define

$$L_j = \partial_j - \frac{x_j}{|x|} \partial_r = \partial_j - \sum_{k=1}^n \frac{x_j x_k}{|x|^2} \partial_k, \quad j = 1, \dots, n,$$

to have $Lf = (L_1 f, \dots, L_n f)$. We use the polar coordinate $x = r\omega, r = |x|, \omega = x/|x|$ for $x \in \mathbb{R}^n \setminus \{0\}$. We may write $f(x) = f(r, \omega)$. If $f(r, \omega)$ depends on r only, that is $f(x) = f(r)$, we have

$$\begin{aligned}\partial_r f(x) &= \sum_{j=1}^n \frac{x_j}{|x|} \partial_j f(|x|) = \sum_{j=1}^n \frac{x_j}{|x|} f'(r) \partial_j |x| = f'(r) \sum_{j=1}^n \frac{x_j^2}{|x|^2} = f'(r), \\ L_j f(x) &= \left(\partial_j - \frac{x_j}{|x|} \partial_r \right) f(|x|) = f'(r) \frac{x_j}{|x|} - \frac{x_j}{|x|} f'(r) = 0.\end{aligned}$$

2010 *Mathematics Subject Classification.* Primary 26D10, Secondary 35A23.

Key words and phrases. Spherical harmonic decomposition, Improved Hardy inequalities, radial and spherical derivatives.

If $f(r, \omega)$ depends on ω only, that is $f(x) = f(\omega) = f\left(\frac{x}{|x|}\right)$, we have

$$\begin{aligned}\partial_r f(x) &= \sum_{j=1}^n \frac{x_j}{|x|} \partial_j \left(f\left(\frac{x}{|x|}\right) \right) = \sum_{j=1}^n \frac{x_j}{|x|} \sum_{k=1}^n (\partial_k f)\left(\frac{x}{|x|}\right) \partial_j \frac{x_k}{|x|} \\ &= \sum_{j=1}^n \frac{x_j}{|x|} \sum_{k=1}^n (\partial_k f)(\omega) \left(\frac{\delta_{j,k}}{|x|} - \frac{x_j x_k}{|x|^3} \right) \\ &= \sum_{j=1}^n \frac{x_j}{|x|} (\partial_j f)(\omega) \frac{1}{|x|} - \sum_{j=1}^n \sum_{k=1}^n (\partial_k f)(\omega) \frac{x_j^2 x_k}{|x|^4} = 0, \\ L_j f(x) &= \left(\partial_j - \frac{x_j}{|x|} \partial_r \right) \left(f\left(\frac{x}{|x|}\right) \right) = \partial_j \left(f\left(\frac{x}{|x|}\right) \right) = \partial_j f(x),\end{aligned}$$

if we continue to calculate

$$\begin{aligned}\partial_j \left(f\left(\frac{x}{|x|}\right) \right) &= \frac{1}{|x|} (\partial_j f)(\omega) - \sum_{k=1}^n (\partial_k f)(\omega) \frac{x_j x_k}{|x|^3} \\ &= \frac{1}{|x|} (\partial_j f)(\omega) - \frac{x_j}{|x|^2} (\partial_r f)(\omega) \\ &= \frac{1}{|x|} (L_j f)(\omega) = \left(\frac{1}{|x|} L_j f \right)(\omega) = (L_j f)(\omega).\end{aligned}$$

Since we have Leibniz rule

$$\partial_r(fg) = f\partial_r g + g\partial_r f, \quad L_j(fg) = fL_j g + gL_j f,$$

we estimate for $f = f(r)$, $g = g(\omega)$ and general $h = h(r, \omega)$ as

$$(1.1) \quad \partial_r(gh) = g\partial_r h, \quad L_j(fh) = fL_j h.$$

We have the pointwise orthogonality between Lf for any f and x as

$$Lf \cdot x = \sum_{j=1}^n \left(\partial_j f - \frac{x_j}{|x|} \partial_r f \right) x_j = x \cdot \nabla f - |x| \partial_r f = 0.$$

Therefore we have the orthogonality pointwisely between Lf and $\frac{x}{|x|} \partial_r f$ for any f ,

$$Lf \cdot \frac{x}{|x|} \partial_r f = \frac{\partial_r f}{|x|} (Lf \cdot x) = 0.$$

We use the inner product and the norm for $L^2(\mathbb{R}^n)$,

$$(f|g) = \int_{\mathbb{R}^n} f(x) \overline{g(x)} dx, \quad \|f\|_2 = \|f\|_{L^2} = \sqrt{(f|f)}.$$

We define

$$\|Lf\|_2^2 = \sum_{j=1}^n \|L_j f\|_2^2.$$

We have the relation $\nabla = \frac{x}{|x|} \partial_r + L$ and, as a corollary, the following orthogonality in the Hilbert space $L^2(\mathbb{R}^n)$.

Proposition 1.

$$(1.2) \quad \|\nabla f\|_2^2 = \|\partial_r f\|_2^2 + \|L_f\|_2^2.$$

Proof. We estimate

$$\begin{aligned} \|\nabla f\|_2^2 &= \sum_{j=1}^n \left\| \frac{x_j}{|x|} \partial_r f + L_j f \right\|_2^2 \\ &= \sum_{j=1}^n \left\| \frac{x_j}{|x|} \partial_r f \right\|_2^2 + 2\operatorname{Re} \int_{\mathbb{R}^n} \frac{\partial_r f}{|x|} x \cdot \overline{L_f} dx + \sum_{j=1}^n \|L_j f\|_2^2 = \|\partial_r f\|_2^2 + \|L_f\|_2^2. \end{aligned}$$

□

We give the following two results by parts for L_j .

$$\begin{aligned} (L_j u | v) &= -(u | \partial_j v) + \sum_{k=1}^n \left(u \left| \partial_k \left(\frac{x_j x_k}{|x|^2} v \right) \right. \right) \\ &= -(u | L_j v) + (n-1) \left(u \left| \frac{x_j}{|x|^2} v \right. \right), \end{aligned}$$

and so

$$\sum_{j=1}^n (L_j u | L_j v) = - \sum_{j=1}^n (u | L_j^2 v) + (n-1) \left(u \left| \frac{1}{|x|^2} x \cdot Lv \right. \right) = - \sum_{j=1}^n (u | L_j^2 v).$$

We consider second order derivative. We introduce the Laplacian on the unit sphere, that is, the Laplace-Beltrami operator.

$$\Delta_{S^{n-1}} := \sum_{1 \leq j < k \leq n} (x_j \partial_k - x_k \partial_j)^2.$$

We have the following relation between the second order of the spherical derivative L and the Laplace-Beltrami operator.

$$\begin{aligned} \sum_{j=1}^n (|x| L_j)^2 &= |x|^2 \sum_{j=1}^n L_j^2 = |x|^2 \sum_{j=1}^n \left(\partial_j^2 - 2 \frac{x_j}{|x|} \partial_j \partial_r + \frac{x_j^2}{|x|^2} \partial_r^2 \right) \\ &= (x_1^2 + \cdots + x_n^2) (\partial_1^2 + \cdots + \partial_n^2) - (x_1^2 \partial_1^2 + \cdots + x_n^2 \partial_n^2) - 2 \sum_{1 \leq j < k \leq n} x_j \partial_j x_k \partial_k \\ &= \sum_{1 \leq j < k \leq n} (x_j^2 \partial_k^2 + x_k^2 \partial_j^2) - 2 \sum_{1 \leq j < k \leq n} x_j \partial_j x_k \partial_k = \Delta_{S^{n-1}}, \end{aligned}$$

where we use (1.1) at the first equality. The following is also well-known

$$\Delta = \partial_r^2 + \frac{n-1}{|x|} \partial_r + \frac{1}{|x|^2} \Delta_{S^{n-1}}.$$

2. PRELIMINARIES PART 2

We denote by $L_{\text{rad}}^2(\mathbb{R}^n)$ and $H_{\text{rad}}^1(\mathbb{R}^n)$ the closed subspaces $L^2(\mathbb{R}^n)$ and $H^1(\mathbb{R}^n)$, respectively, of radial functions:

$$\begin{aligned} L_{\text{rad}}^2(\mathbb{R}^n) &:= \{f \in L^2(\mathbb{R}^n); \text{ There exists } u \in L^2(0, \infty) \text{ such that} \\ &\quad f(x) = u(|x|)|x|^{\frac{1-n}{2}} \text{ for almost all } x \in \mathbb{R}^n \setminus \{0\}\}, \\ H_{\text{rad}}^1(\mathbb{R}^n) &:= \{f \in (H^1 \cap L_{\text{rad}}^2)(\mathbb{R}^n); \partial_r f \in L_{\text{rad}}^2(\mathbb{R}^n)\}. \end{aligned}$$

For any $f \in L^2(\mathbb{R}^n)$, we denote by Pf its radial average over the unit sphere

$$(Pf)(x) := \frac{1}{\sigma_{n-1}} \int_{S^{n-1}} f(|x|\omega) d\sigma(\omega), \quad x \in \mathbb{R}^n.$$

Then $P : f \mapsto Pf$ induces the orthogonal projection from $L^2(\mathbb{R}^n)$ onto $L_{\text{rad}}^2(\mathbb{R}^n)$ as well as from $H^1(\mathbb{R}^n)$ onto $H_{\text{rad}}^1(\mathbb{R}^n)$ (see Proposition 3 below).

Lemma 2.

$$P^2 = P, \quad P^* = P.$$

Proof. For any radial function $f \in L^2(\mathbb{R}^n)$, and we write $f(x) = f(|x|)$, we have

$$(Pf)(x) = \frac{1}{\sigma_{n-1}} \int_{S^{n-1}} f(|x|) d\sigma(\omega) = \frac{f(|x|)}{\sigma_{n-1}} \int_{S^{n-1}} d\sigma(\omega) = f(x).$$

Therefore for any f we have the radial Pf and so we have $P(Pf) = Pf$. For any u, v , we calculate

$$\begin{aligned} (Pu|v) &= \int_{\mathbb{R}^n} \frac{1}{\sigma_{n-1}} \int_{S^{n-1}} u(|x|\omega) d\sigma(\omega) \bar{v}(x) dx \\ &= \int_{S^{n-1}} \int_0^\infty \left(\frac{1}{\sigma_{n-1}} \int_{S^{n-1}} u(r\omega) d\sigma(\omega) \right) \bar{v}(r\theta) r^{n-1} dr d\sigma(\theta) \\ &= \int_{S^{n-1}} \int_0^\infty \left(\frac{1}{\sigma_{n-1}} \int_{S^{n-1}} \bar{v}(r\theta) d\sigma(\theta) \right) u(r\omega) r^{n-1} dr d\sigma(\omega) = (u|Pv). \end{aligned}$$

□

We define the orthogonal projection onto the orthogonal complement of these spaces

$$P^\perp = I - P.$$

We have $f = Pf + P^\perp f$.

Proposition 3. *The following relations hold:*

- (1) $(P^\perp)^2 = P^\perp, \quad (P^\perp)^* = P^\perp.$
- (2) $PP^\perp = P^\perp P = 0.$
- (3) $(Pu|P^\perp v) = 0, \quad \|u\|_2^2 = \|Pu\|_2^2 + \|P^\perp u\|_2^2, \quad u, v \in L^2(\mathbb{R}^n).$
- (4) $(\nabla Pu|\nabla P^\perp v) = 0, \quad \|\nabla u\|_2^2 = \|\nabla Pu\|_2^2 + \|\nabla P^\perp u\|_2^2, \quad u, v \in H^1(\mathbb{R}^n).$

Proof. The all orthogonal projections satisfy the properties (1), (2) and (3).

(1) From Lemma 2,

$$\begin{aligned} (P^\perp)^2 &= I - 2P + P^2 = I - 2P + P = P^\perp, \\ (P^\perp u|v) &= (u|v) - (Pu|v) = (u|v) - (u|Pv) = (u|P^\perp v). \end{aligned}$$

(2) From Lemma 2,

$$PP^\perp = P^\perp P = P - P^2 = P - P = 0.$$

(3) From (2),

$$(Pu|P^\perp v) = (u|PP^\perp v) = (u|0) = 0.$$

$$\|u\|^2 = \|Pu + P^\perp u\|^2 = \|Pu\|^2 + 2\operatorname{Re}(Pu|P^\perp v) + \|P^\perp v\|^2 = \|Pu\|^2 + \|P^\perp v\|^2.$$

(4) We have

$$\begin{aligned} \partial_r Pu &= \frac{1}{\sigma_{n-1}|x|} \sum_{j=1}^n x_j \int_{S^{n-1}} \partial_j(u(|x|\omega)) d\sigma(\omega) \\ &= \frac{1}{\sigma_{n-1}|x|} \int_{S^{n-1}} \sum_{j=1}^n x_j \frac{x_j}{|x|} (\nabla u)(|x|\omega) \cdot \omega d\sigma(\omega) = P(\nabla u \cdot \omega) = P\partial_r u. \end{aligned}$$

We have seen $L_j v = 0$ for any radial function v , and so $Lv = 0$ and $LP = 0$. □

By using the orthogonal projection P and P^\perp , we have the following orthogonal decompositions:

$$\begin{aligned} L^2(\mathbb{R}^n) &= L_{\text{rad}}^2(\mathbb{R}^n) \oplus (L_{\text{rad}}^2(\mathbb{R}^n))^\perp, \\ H^1(\mathbb{R}^n) &= H_{\text{rad}}^1(\mathbb{R}^n) \oplus (H_{\text{rad}}^1(\mathbb{R}^n))^\perp. \end{aligned}$$

We also use the complete orthogonal decomposition

$$(2.1) \quad L^2(\mathbb{R}^n) = \bigoplus_{k \geq 0} \mathcal{H}^k(\mathbb{R}^n),$$

where $\mathcal{H}^k(\mathbb{R}^n)$ is a closed subspace spanned by spherical harmonics of order k multiplied by radial functions. We denote by P_k the associated orthogonal projection. We refer the reader to [4, 5, 6, 24, 26] for details on the decomposition (2.1). Here we notice that

$$\begin{aligned} P &= P_0, \\ L_{\text{rad}}^2(\mathbb{R}^n) &= \mathcal{H}^0(\mathbb{R}^n). \end{aligned}$$

We introduce the eigenvalues of the spherical harmonics

$$\Delta_{S^{n-1}} P_k = -k(k+n-2)P_k, \quad k \geq 0.$$

3. HARDY'S INEQUALITIES AND EQUALITIES, KNOWN RESULTS.

In this talk, we study the classical Hardy inequality of the form

$$(3.1) \quad \left(\frac{n-2}{2}\right)^2 \left\| \frac{f}{|x|} \right\|_2^2 \leq \|\nabla f\|_2^2$$

for all $f \in H^1(\mathbb{R}^n)$ with $n \geq 3$. We remark the constant $(n-2/2)^2$ is optimal and is not attained. The following also holds

$$\left(\frac{n-2}{2}\right)^2 \left\| \frac{f}{|x|} \right\|_2^2 \leq \|\partial_r f\|_2^2$$

for all $f \in H^1(\mathbb{R}^n)$ with $n \geq 3$.

We call a kinds of the followings Hardy's equalities. Dolbeault-Volzone (2012) and Bogdan-Dyda-Kim (2016) showed

$$\left(\frac{n-2}{2}\right)^2 \left\| \frac{f}{|x|} \right\|_2^2 = \|\nabla f\|_2^2 - \left\| \nabla f + \frac{n-2}{2} \frac{x}{|x|^2} f \right\|_2^2.$$

M-Ozawa-Wadade (2016) showed

$$\left(\frac{n-2}{2}\right)^2 \left\| \frac{f}{|x|} \right\|_2^2 = \|\partial_r f\|_2^2 - \left\| \partial_r f + \frac{n-2}{2|x|} f \right\|_2^2.$$

Meanwhile, Ekholm-Frank (2006) introduce the following as an improved Hardy inequality

$$(3.2) \quad \frac{n^2}{4} \left\| \frac{f}{|x|} \right\|_2^2 \leq \|\nabla f\|_2^2$$

for all $f \in C_0^\infty(\mathbb{R}^n)$ with $n \geq 2$ satisfying

$$(3.3) \quad \int_{S^{n-1}} f(r\omega) d\sigma(\omega) = 0$$

for all $r \geq 0$, where σ is the Lebesgue measure on the unit sphere $S^{n-1} = \{\omega \in \mathbb{R}^n; |\omega| = 1\}$, [7, 12]. In [12], the inequality (3.2) is referred to as an improved Hardy inequality on the basis of the improvement in the coefficient $\frac{n^2}{4}$ on the left hand side of (3.2), which is larger than the corresponding coefficient $\left(\frac{n-2}{4}\right)^2$ on the left hand side of (3.1), as well as of the applicable range of dimensions, in particular, $n = 2$ is now admissible.

4. HARDY'S INEQUALITIES AND EQUALITIES, THEOREMS.

We now state the main results in this paper.

Theorem 4. *Let $n \geq 2$. Then, the following equality*

$$(4.1) \quad (n-1) \left\| \frac{P^\perp f}{|x|} \right\|_2^2 = \|LP^\perp f\|_2^2 - \sum_{k=2}^{\infty} (k-1)(k+n-1) \left\| \frac{P_k f}{|x|} \right\|_2^2$$

holds for all $f \in H^1(\mathbb{R}^n)$.

Corollary 5. *Let $n \geq 2$. Then the following inequality*

$$(4.2) \quad (n-1) \left\| \frac{P^\perp f}{|x|} \right\|_2^2 \leq \|LP^\perp f\|_2^2$$

holds for all $f \in H^1(\mathbb{R}^n)$. Equality holds in (4.2) if and only if there exist $\alpha \in \mathbb{C}^n$ and $g, h \in H_{\text{rad}}^1(\mathbb{R}^n)$ such that

$$(4.3) \quad f(x) = (\alpha \cdot x)g(x) + h(x)$$

for almost all $x \in \mathbb{R}^n \setminus \{0\}$, where $\alpha \cdot y = \sum_{j=1}^n \alpha_j y_j$ for $y \in \mathbb{R}^n$. In this case, both sides of (4.2) are given by

$$(4.4) \quad (n-1) \left\| \frac{P^\perp f}{|x|} \right\|_2^2 = \|LP^\perp f\|_2^2 = \frac{n-1}{n} \sigma_{n-1} |\alpha|^2 \int_0^\infty |u(r)|^2 dr,$$

where $u \in L^2(0, \infty)$ satisfies $g(x) = u(|x|)|x|^{-\frac{n-1}{2}}$ for almost all $x \in \mathbb{R}^n \setminus \{0\}$ and $|\alpha|^2 = \sum_{j=1}^n \alpha_j \bar{\alpha}_j$.

Theorem 6. *Let $n \geq 2$. Then, the following equalities*

$$\begin{aligned}
(4.5) \quad & \left(\frac{n-2}{2}\right)^2 \left\| \frac{f}{|x|} \right\|_2^2 + (n-1) \left\| \frac{P^\perp f}{|x|} \right\|_2^2 \\
&= \left(\frac{n-2}{2}\right)^2 \left\| \frac{Pf}{|x|} \right\|_2^2 + \frac{n^2}{4} \left\| \frac{P^\perp f}{|x|} \right\|_2^2 \\
&= \|\nabla f\|_2^2 - \left\| \left(\partial_r + \frac{n-2}{2|x|} \right) f \right\|_2^2 - \sum_{k=2}^{\infty} (k-1)(k+n-1) \left\| \frac{P_k f}{|x|} \right\|_2^2
\end{aligned}$$

hold for all $f \in H^1(\mathbb{R}^n)$.

Corollary 7. *Let $n \geq 2$. Then, the following inequality*

$$(4.6) \quad \left(\frac{n-2}{2}\right)^2 \left\| \frac{f}{|x|} \right\|_2^2 + (n-1) \left\| \frac{P^\perp f}{|x|} \right\|_2^2 \leq \|\nabla f\|_2^2.$$

holds for all $f \in H^1(\mathbb{R}^n)$. Equality holds in (4.6) if and only if $f = 0$.

Theorem 8. *Let $n \geq 2$. Then, the following equality*

$$\begin{aligned}
(4.7) \quad & \frac{n^2}{4} \left\| \frac{P^\perp f}{|x|} \right\|_2^2 = \|\nabla P^\perp f\|_2^2 - \left\| \left(\partial_r + \frac{n-2}{2|x|} \right) P^\perp f \right\|_2^2 \\
& \quad - \sum_{k=2}^{\infty} (k-1)(k+n-1) \left\| \frac{P_k f}{|x|} \right\|_2^2
\end{aligned}$$

holds for all $f \in H^1(\mathbb{R}^n)$.

Corollary 9 (Improved Hardy inequality [7, 12]). *Let $n \geq 2$. Then, the following inequality*

$$(4.8) \quad \frac{n^2}{4} \left\| \frac{P^\perp f}{|x|} \right\|_2^2 \leq \|\nabla P^\perp f\|_2^2$$

holds for all $f \in H^1(\mathbb{R}^n)$. Equality holds in (4.8) if and only if $f \in H_{\text{rad}}^1(\mathbb{R}^n)$. In this case, both sides of (4.8) vanish.

The inequality (4.6) improved the Hardy inequality (3.1) in the sense that (4.6) reveals a novel term $(n-1) \left\| \frac{P^\perp f}{|x|} \right\|_2^2$ on the left hand side and that the improvement in (3.2) in two space dimensions arises as a result of the existence of the novel term $\left\| \frac{P^\perp f}{|x|} \right\|_2^2$ when $n = 2$. Moreover, (4.6) clarifies the contribution by the orthogonal component to $H_{\text{rad}}^1(\mathbb{R}^n)$ with coefficient $n-1$, which together with the standard coefficient $\left(\frac{n-2}{2}\right)^2$ yields the improved coefficient $\frac{n^2}{4}$ in (3.2) on the basis of the simple identity $\left(\frac{n-2}{2}\right)^2 + (n-1) = \frac{n^2}{4}$.

In Section 2, we prove a density lemma, which enables us to prove the main theorems for functions in $C_0^\infty(\mathbb{R}^n \setminus \{0\})$. In Section 3, we prove the main results stated above. Furthermore, we also include a justification of the observation that the constant $\frac{n^2}{4}$ in the improved Hardy inequality (4.8) is best possible and thus we establish the nonexistence of nontrivial extremizers for the improved Hardy inequality.

5. PROOFS OF THE MAIN THEOREMS

The following proposition is important in the proofs of the main theorems:

Proposition 10. $P^\perp(C_0^\infty(\mathbb{R}^n \setminus \{0\}))$ is dense in $P^\perp(H^1(\mathbb{R}^n)) = (H_{\text{rad}}^1(\mathbb{R}^n))^\perp$ if $n \geq 2$.

In this section, we prove Theorems 4, 6, 8 and their corollaries. By a density argument based on Proposition 10, it suffices to prove the theorems for functions in $C_0^\infty(\mathbb{R}^n \setminus \{0\})$. In the proofs below, all functions are supposedly elements of $C_0^\infty(\mathbb{R}^n \setminus \{0\})$.

Proof of Theorem 4. Let $f \in C_0^\infty(\mathbb{R}^n \setminus \{0\})$. By Propositions 3 and ??, the first term on the right hand side of (4.1) is represented as

$$\begin{aligned}
 \|LP^\perp f\|_2^2 &= \sum_{j=1}^n \|L_j P^\perp f\|_2^2 \\
 &= - \sum_{j=1}^n (L_j^2 P^\perp f | P^\perp f) \\
 &= - \sum_{j=1}^n \sum_{k=0}^{\infty} (L_j^2 P^\perp P_k f | P^\perp f) \\
 &= - \sum_{j=1}^n \sum_{k=1}^{\infty} (L_j^2 P^\perp P_k f | P^\perp f) \\
 &= - \sum_{k=1}^{\infty} (|x|^{-2} \Delta_{S^{n-1}} P_k P^\perp f | P^\perp f)
 \end{aligned}$$

and hence

$$\begin{aligned}
 \|LP^\perp f\|_2^2 &= - \sum_{k=1}^{\infty} \int_0^\infty (\Delta_{S^{n-1}} P_k P^\perp f)(r(\cdot) | P^\perp f(r(\cdot)))_{L^2(S^{n-1})} r^{n-3} dr \\
 &= \sum_{k=1}^{\infty} k(k+n-2) \int_0^\infty (P_k P^\perp f)(r(\cdot) | P^\perp f(r(\cdot)))_{L^2(S^{n-1})} r^{n-3} dr \\
 &= \sum_{k=1}^{\infty} k(k+n-2) \left\| \frac{1}{|x|} P_k P^\perp f \right\|_2^2 \\
 &= (n-1) \sum_{k=1}^{\infty} \left\| \frac{1}{|x|} P_k P^\perp f \right\|_2^2 + \sum_{k=1}^{\infty} (k(k+n-2) - (n-1)) \left\| \frac{1}{|x|} P_k P^\perp f \right\|_2^2 \\
 &= (n-1) \sum_{k=0}^{\infty} \left\| P_k \left(\frac{P^\perp f}{|x|} \right) \right\|_2^2 + \sum_{k=2}^{\infty} (k-1)(k+n-1) \left\| \frac{1}{|x|} P_k P^\perp f \right\|_2^2 \\
 &= (n-1) \left\| \frac{P^\perp f}{|x|} \right\|_2^2 + \sum_{k=2}^{\infty} (k-1)(k+n-1) \left\| \frac{P_k f}{|x|} \right\|_2^2,
 \end{aligned}$$

where we have used relations

$$\begin{aligned} P_0 P^\perp &= P^\perp P_0 = 0, \\ P_k P^\perp &= P_k(I - P_0) = P_k \quad \text{for } k \geq 1, \\ P_k \left(\frac{g}{|x|} \right) &= \frac{1}{|x|} P_k g \quad \text{for } k \geq 0. \end{aligned}$$

This completes the proof. \square

Proof of Corollary 5. The inequality (4.2) is a direct consequence of (4.1). The equality in (4.2) holds if and only if $\frac{P_k f}{|x|} = 0$ for all nonnegative integers k with $k \geq 2$, namely, $\frac{f}{|x|} \in \mathcal{H}_0 \oplus \mathcal{H}_1$. This proves (4.3). Then we take $g, h \in H_{\text{rad}}^1(\mathbb{R}^n)$ as in (4.3). In this case, we have

$$\begin{aligned} (P^\perp f)(x) &= (\alpha \cdot x)g(x), \\ (LP^\perp f)(x) &= \left(\alpha - \frac{x}{|x|} \left(\alpha \cdot \frac{x}{|x|} \right) \right) g(x) \end{aligned}$$

for almost all $x \in \mathbb{R}^n \setminus \{0\}$. Since g is radial and $\frac{P^\perp f}{|x|} \in \mathcal{H}_1$, a new function $u \in L^2(0, \infty)$ is defined to satisfy $g(x) = u(|x|)|x|^{-\frac{n-1}{2}}$ for almost all $x \in \mathbb{R}^n \setminus \{0\}$. We evaluate two integrals in (4.2) as

$$\begin{aligned} \left\| \frac{P^\perp f}{|x|} \right\|_2^2 &= \int_0^\infty \int_{S^{n-1}} |\alpha \cdot \omega|^2 |g(r\omega)|^2 d\sigma(\omega) r^{n-1} dr \\ &= \int_{S^{n-1}} |\alpha \cdot \omega|^2 d\sigma(\omega) \int_0^\infty |u(r)|^2 dr \\ &= \frac{\sigma_{n-1}}{n} |\alpha|^2 \int_0^\infty |u(r)|^2 dr, \\ \|LP^\perp f\|_2^2 &= \int_0^\infty \int_{S^{n-1}} |\alpha - \omega(\alpha \cdot \omega)|^2 d\sigma(\omega) |u(r)|^2 dr \\ &= \int_{S^{n-1}} (|\alpha|^2 - |\alpha \cdot \omega|^2) d\sigma(\omega) \int_0^\infty |u(r)|^2 dr \\ &= \left(1 - \frac{1}{n} \right) \sigma_{n-1} |\alpha|^2 \int_0^\infty |u(r)|^2 dr, \end{aligned}$$

where we have used

$$\begin{aligned} \int_{S^{n-1}} |\alpha \cdot \omega|^2 d\sigma(\omega) &= \sum_{j=1}^n \sum_{k=1}^n \alpha_j \bar{\alpha}_k \int_{S^{n-1}} \omega_j \omega_k d\sigma(\omega) \\ &= \sum_{j=1}^n |\alpha_j|^2 \int_{S^{n-1}} \omega_j^2 d\sigma(\omega) \\ &= \sum_{j=1}^n |\alpha_j|^2 \frac{1}{n} \sigma_{n-1} = \frac{1}{n} \sigma_{n-1} |\alpha|^2. \end{aligned}$$

This proves (4.4). □

Proof of Theorem 6. The equality (4.5) follows from (??), (1.2), (4.1), and $\|LP^\perp f\|_2 = \|Lf\|_2$. □

Proof of Corollary 7. The equality in (4.6) holds if and only if (4.3) and

$$|x|^{1-\frac{n}{2}}\partial_r(|x|^{\frac{n}{2}-1}f) = \left(\partial_r + \frac{n-2}{2|x|}\right)f = 0.$$

Then f is written as $f(x) = |x|^{1-\frac{n}{2}}\psi\left(\frac{x}{|x|}\right)$ for some function $\psi : S^{n-1} \rightarrow \mathbb{C}$, which together with (4.3) implies that $f(x) = |x|^{1-\frac{n}{2}}\left(\alpha \cdot \frac{x}{|x|}\right)$ for some $\alpha \in \mathbb{C}^n$. In this case, $\frac{f}{|x|} \in L^2(\mathbb{R}^n)$ if and only if $\alpha = 0$, which means $f = 0$. □

Proof of Theorem 8. The equality (4.7) follows by substituting f by $P^\perp f$ in (4.5). □

Proof of Corollary 9. The equality (4.8) follows if and only if $P^\perp f = 0$, which means $f \in H_{\text{rad}}^1(\mathbb{R}^n)$. □

ACKNOWLEDGMENT

The first author was supported by JSPS KAKENHI Grant number 16H05995 and 16K1377, the second author was supported by JSPS KAKENHI Grant number JP16K05191, the third author was supported by JSPS KAKENHI Grant number 26247014, 16K13771 and 18KK0073.

REFERENCES

- [1] W. BECKNER, *Pitt's inequality and the fractional Laplacian: sharp error estimates*, Forum Math. **24** (2012), 177–209.
- [2] W. BECKNER, *Pitt's inequality and the uncertainty principle*, Proc. Amer. Math. Soc., **123**, (1995), 1897–1905.
- [3] W. BECKNER, *Pitt's inequality with sharp convolution estimates*, Proc. Amer. Math. Soc., **136**, (2008), 1871–1885.
- [4] N. BEZ, C. JEAVONS, T. OZAWA, AND M. SUGIMOTO, *Stability of trace theorems on the sphere*, J. Geom. Anal., **28** (2018), 1456–1476.
- [5] N. BEZ, S. MACHIHARA, AND M. SUGIMOTO, *Extremisers for the trace theorem on the sphere*, Math. Res. Lett., **23** (2016), 633–647.
- [6] N. BEZ, H. SAITO, AND M. SUGIMOTO, *Applications of the Funk–Hecke theorem to smoothing and trace estimates*, Adv. Math., **285** (2015), 1767–1795.
- [7] M. SH. BIRMAN, AND A. LAPTEV, *The negative discrete spectrum of a two-dimensional Schrödinger operator*, Comm. Pure Appl. Math., **49** (1996), 967–997.
- [8] K. BOGDAN, B. DYDA, AND P. KIM, *Hardy inequalities and non-explosion results for semigroups*, Potential Anal., **44**, (2016), 229–247.
- [9] D. G. COSTA, *Some new and short proofs for a class of Caffarelli-Kohn-Nirenberg type inequalities*, J. Math. Anal. Appl., **337** (2008), 311–317.
- [10] E. B. DAVIES AND A. M. HINZ, *Explicit constants for Rellich inequalities in $L^p(\Omega)$* , Math. Z., **227** (1998), 511–523.
- [11] J. DOLBEAULT AND B. VOLZONE, *Improved Poincaré inequalities*, Nonlinear Anal., **75** (2012), 5985–6001.
- [12] T. EKHOLOM AND R. L. FRANK, *On Lieb–Thirring inequalities for Schrödinger operators with virtual level*, Comm. Math. Phys., **264** (2006), 725–740.
- [13] N. GHOUSSOUB AND A. MORADIFAM, *Bessel pairs and optimal Hardy and Hardy-Rellich inequalities*, Math. Ann., **349** (2011), 1–57.

- [14] N. GHOUSSOUB AND A. MORADIFAM, *On the best possible remaining term in the Hardy inequality*, Proc. Natl. Acad. Sci. USA, **1305** (2008), 13746–13751.
- [15] N. IOKU, M. ISHIWATA, AND T. OZAWA, *Sharp remainder of a critical Hardy inequality*, Arch. Math. (Basel), **1**, (2016), 65–71.
- [16] N. IOKU, M. ISHIWATA, AND T. OZAWA, *Hardy type inequalities in L^p with sharp remainders*, J. Inequal. Appl., (2017), Paper No. 5, 7.
- [17] N. IOKU, AND M. ISHIWATA, *A scale invariant form of a critical Hardy inequality*, Int. Math. Res. Not. IMRN, (2015), 8830–8846.
- [18] E. H. LIEB, *Sharp constants in the Hardy–Littlewood–Sobolev and related inequalities*, Ann. of Math., **118**, (1983), 349–374.
- [19] S. MACHIHARA, T. OZAWA, AND H. WADADE, *Hardy type inequalities on balls*, Tohoku Math. J. (2), **65** (2013), 321–330.
- [20] S. MACHIHARA, T. OZAWA, AND H. WADADE, *Remarks on the Rellich inequality*, Math. Z., **286**, (2017), 1367–1373.
- [21] S. MACHIHARA, T. OZAWA, AND H. WADADE, *Remarks on the Hardy type inequalities with remainder terms in the framework of equalities*, arXiv:1611.03580 (2016), to appear in Adv. Stud. Pure Math.
- [22] M. RUZHANSKY, D. SURAGAN, *Hardy and Rellich inequalities, identities, and sharp remainders on homogeneous groups*, Adv. Math., **317** (2017), 799–822.
- [23] M. SANO, F. TAKAHASHI, *Scale invariance structures of the critical and the subcritical Hardy inequalities and their improvements*, Calc. Var. Partial Differential Equations, **56** (2017), Art. 69, 14.
- [24] E. M. STEIN AND G. WEISS, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton Mathematical Series, No. 32, Princeton University Press, Princeton, N.J., (1971).
- [25] F. TAKAHASHI, *A simple proof of Hardy’s inequality in a limiting case*, Arch. Math. (Basel). **104** (2015), 77–82.
- [26] B. WALTHER, *Regularity, decay, and best constants for dispersive equations*, J. Funct. Anal., **189** (2002), 325–335.
- [27] D. YAPAEV, *Sharp constants in the Hardy–Rellich inequalities*, J. Funct. Anal., **168**, (1999), 121–144.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, SAITAMA UNIVERSITY, SAITAMA 338-8570, JAPAN

E-mail address: machihara@rimath.saitama-u.ac.jp