

# Generalized cone-continuity of set-valued maps with scalarization\*

Premyuda Dechboon<sup>†</sup> and Tamaki Tanaka<sup>‡</sup>

Graduate School of Science and Technology, Niigata University<sup>§</sup>

Dedicated to the memory of Professor Kazimierz Goebel

## 1 Introduction

A set-valued map is a map that associates with a set depending on each element. Considering a singleton set, a set-valued mapping can be launched as a single-valued mapping. Because the concept of set-valued mappings includes single-valued maps, set-valued optimization presents a significant generalization along with unification of scalar and vector optimization problems.

On the other hand, transforming vectors or sets into real number, in other words scalarization, processes a quintessential methodology solving optimization problems with vector-valued or set-valued maps. One of challenging scalarizing functions ensues sublinear scalarization introduced by Tammer (see [2], [3], [4] and [5])

$$h_C(v; d) := \inf \{t \in \mathbb{R} \mid v \in td - C\}$$

where  $C$  is a convex cone in a real topological vector space and  $d \in C$ .

In general, composition is an operator analyzed a function from the results of another function. Several mathematical properties of each nested function are usually preserved by a composite operation. A continuous map composition, for example, is continuous on topological spaces. Under certain assumptions, we may characterize solutions for multicriteria questions using scalarization based on this attribute. This prompts us to explore how composite functions involving

---

\*This work is supported by Grant-in-Aid for Scientific Research (C) (JSPS KAKENHI Grant Number 21K03367) from Japan Society for the Promotion of Science (JSPS).

<sup>†</sup>A double degree student in Niigata University (Niigata, Japan) and King Mongkut's University of Technology Thonburi (Bangkok, Thailand); 新潟大学大学院自然科学研究科とモンクット王トンブリ工科大学 (タイ) の DDP 学生. E-mail: dechboon@m.sc.niigata-u.ac.jp

<sup>‡</sup>Faculty of Science, Niigata University; E-mail: tamaki@math.sc.niigata-u.ac.jp

<sup>§</sup>Niigata 950-2181, Japan; 〒 950-2181 新潟市西区五十嵐 2 の町 8050 番地

a set-valued map and a scalarization function convey continuity of primary set-valued mappings via various scalarization for sets.

Continuity notions of set-valued maps are significant in many branches of mathematics, including nonlinear functional analysis, optimization theory, and convex analysis. Cone-continuity is studied in several direction (see [16] and [9]).

Recently, Ike, Liu, Ogata and Tanaka [6] show certain results on the inheritance property of some kinds of continuity of set-valued maps via scalarization functions for sets: if a set-valued map has a kind of continuity (lower continuity or upper continuity; see [4]) then the composition of its set-valued map and a certain scalarization function assures a similar semicontinuity to that of its scalarization function defined on the family of nonempty subsets of a real topological vector space. Their results are generalizations of results in earlier study by Kuwano, Tanaka and Yamada [15].

The aim of this paper is to propose some idea how to obtain and apply the generalization in [1] by Dechboon and Tanaka of the inheritance property which is introduced by [6].

## 2 Basic Notations

Throughout the paper, let  $X$  be a topological space and  $Y$  a real topological vector space. Let  $\theta_Y$  be the zero vector in  $Y$  and  $\mathcal{P}(Y)$  denote the set of all nonempty subsets of  $Y$ . The topological interior, topological closure, convex hull, and complement of a set  $A \in \mathcal{P}(Y)$  are denoted by  $\text{int } A$ ,  $\text{cl } A$ ,  $\text{co } A$ , and  $A^c$ , respectively. Furthermore, we assume that  $C$  is a convex cone in  $Y$  with  $\text{int } C \neq \emptyset$  and  $\theta_Y \in C$ . Then,  $C + C = C$  holds, and  $\text{int } C$  and  $\text{cl } C$  are also convex cones. Accordingly, we can define a preorder  $\leq_C$  on  $Y$  induced by  $C$  as follows:

$$\text{for } y_1, y_2 \in Y, y_1 \leq_C y_2 \stackrel{\text{def}}{\iff} y_2 - y_1 \in C.$$

This preorder is compatible with the linear structure of  $Y$ :

$$\text{for all } y_1, y_2, y_3 \in Y, \quad y_1 \leq_C y_2 \implies y_1 + y_3 \leq_C y_2 + y_3; \quad (1)$$

$$\text{for all } y_1, y_2 \in Y \text{ and } t > 0, \quad y_1 \leq_C y_2 \implies ty_1 \leq_C ty_2. \quad (2)$$

When  $C$  is pointed (i.e.,  $C \cap (-C) = \{\theta_Y\}$ ),  $\leq_C$  is antisymmetric and then a partial order.

**Proposition 1.** *Let  $C, C'$  be convex cones in  $Y$  and  $d \in Y$ . Assume that  $C + (0, +\infty)d \subset C'$ . Then, for any  $v_1, v_2 \in Y$  and  $t, t' \in \mathbb{R}$  with  $t > t'$ ,*

$$v_1 + td \leq_C v_2 \implies v_1 + t'd \leq_{C'} v_2.$$

As generalizations of partial orderings for vectors, we give a definition of certain binary relations between sets in  $Y$ , called set relations. This is a modified version of the original one proposed in [12].

**Definition 2** (set relations, [12]). For  $A, B \in \mathcal{P}(Y)$ , we define the following eight types of binary relations on  $\mathcal{P}(Y)$ .

- (i)  $A \leq_C^{(1)} B \stackrel{\text{def}}{\iff} \forall a \in A, \forall b \in B, a \leq_C b \iff A \subset \bigcap_{b \in B} (b - C)$   
 $\iff B \subset \bigcap_{a \in A} (a + C)$ ;
- (ii)  $A \leq_C^{(2L)} B \stackrel{\text{def}}{\iff} \exists a \in A \text{ s.t. } \forall b \in B, a \leq_C b \iff A \cap (\bigcap_{b \in B} (b - C)) \neq \emptyset$ ;
- (iii)  $A \leq_C^{(2U)} B \stackrel{\text{def}}{\iff} \exists b \in B \text{ s.t. } \forall a \in A, a \leq_C b \iff (\bigcap_{a \in A} (a + C)) \cap B \neq \emptyset$ ;
- (iv)  $A \leq_C^{(2)} B \stackrel{\text{def}}{\iff} A \leq_C^{(2L)} B \text{ and } A \leq_C^{(2U)} B \iff A \cap (\bigcap_{b \in B} (b - C)) \neq \emptyset$   
 $\text{and } (\bigcap_{a \in A} (a + C)) \cap B \neq \emptyset$ ;
- (v)  $A \leq_C^{(3L)} B \stackrel{\text{def}}{\iff} \forall b \in B, \exists a \in A \text{ s.t. } a \leq_C b \iff B \subset A + C$ ;
- (vi)  $A \leq_C^{(3U)} B \stackrel{\text{def}}{\iff} \forall a \in A, \exists b \in B \text{ s.t. } a \leq_C b \iff A \subset B - C$ ;
- (vii)  $A \leq_C^{(3)} B \stackrel{\text{def}}{\iff} A \leq_C^{(3L)} B \text{ and } A \leq_C^{(3U)} B \iff B \subset A + C \text{ and } A \subset B - C$ ;
- (viii)  $A \leq_C^{(4)} B \stackrel{\text{def}}{\iff} \exists a \in A, \exists b \in B \text{ s.t. } a \leq_C b \iff A \cap (B - C) \neq \emptyset$   
 $\iff (A + C) \cap B \neq \emptyset$ .

In the above definition, the letters  $L$  and  $U$  stand for “lower” and “upper,” respectively. Each relation  $\leq_C^{(j)}$  is transitive for  $j = 1, 2L, 2U, 3L, 3U$  and not transitive for  $j = 4$ . Since  $\theta_Y \in C$ ,  $\leq_C^{(j)}$  is reflexive for  $j = 3L, 3U, 4$  and hence a preorder for  $j = 3L, 3U$ . Besides, for each  $j = 1, 2L, 2U, 3L, 3U, 4$ , the relation  $\leq_C^{(j)}$  satisfies certain similar properties to conditions (1) and (2) for all  $A, B \in \mathcal{P}(Y)$ ,

- (i)  $A \leq_C^{(j)} B \implies A + y \leq_C^{(j)} B + y$  for  $y \in Y$ ;
- (ii)  $A \leq_C^{(j)} B \implies tA \leq_C^{(j)} tB$  for  $t > 0$ .

Also, we easily obtain the following implications:

$$\begin{cases} A \leq_C^{(1)} B \implies A \leq_C^{(2L)} B \implies A \leq_C^{(3L)} B \implies A \leq_C^{(4)} B; \\ A \leq_C^{(1)} B \implies A \leq_C^{(2U)} B \implies A \leq_C^{(3U)} B \implies A \leq_C^{(4)} B; \\ A \leq_C^{(1)} B \implies A \leq_C^{(2)} B \implies A \leq_C^{(3)} B \implies A \leq_C^{(4)} B \end{cases} \quad (3)$$

for  $A, B \in \mathcal{P}(Y)$ .

**Proposition 3** ([6]). *Let  $C'$  and  $C$  be two nonempty convex cones in  $Y$  and  $d \in Y$ . Assume that  $C' + (0, +\infty)d \subset C$ . Then, for each  $j = 1, 2L, 3L, 2U, 3U, 4$ , any  $A, B \in \mathcal{P}(Y)$ ,  $s, s' \in \mathbb{R}$  with  $s' < s$  and  $t, t' \in \mathbb{R}$  with  $t < t'$ ,*

$$A \leq_{C'}^{(j)} B + s'd \implies A \leq_C^{(j)} B + sd,$$

$$\text{and } A + t'd \leq_{C'}^{(j)} B \implies A + td \leq_C^{(j)} B.$$

### 3 Unification of Scalarizing Functions

Now, we recall the scalarization scheme [13] for sets in a real vector space related to the set relations, which are certain generalizations as unification of several nonlinear scalarizations proposed in [5].

**Definition 4** ([7, 13]). For each  $j = 1, 2L, 3L, 2U, 3U, 4$ , we define

$$I_C^{(j)}(A; V, d) := \inf \left\{ t \in \mathbb{R} \mid A \leq_C^{(j)} (V + td) \right\}, \quad (4)$$

$$S_C^{(j)}(A; V, d) := \sup \left\{ t \in \mathbb{R} \mid (V + td) \leq_C^{(j)} A \right\}, \quad (5)$$

for any  $A, V \in \mathcal{P}(Y)$  and  $d \in Y$ .

The idea of these scalarization functions is introduced in [13], which originates from the idea of Gerstewitz's (Tammer's) sublinear scalarizing functional in [2]; see [4, 7]. This type of scalarization measures how far a given reference set needs to be moved toward a specific direction to fulfill each set relation between a target set and its moved reference set. Note that  $V$  and  $d$  in (4) and (5) are index parameters for scalarization which play key roles as a reference set and a reference direction, respectively.

**Proposition 5** ([7]). *Let  $C$  be a convex cone in  $V$ . The following inequalities hold between each scalarizing function for sets:*

$$\begin{aligned} I_C^{(4)}(A; W, d) &\leq I_C^{(3L)}(A; W, d) \leq I_C^{(2L)}(A; W, d) \leq I_C^{(1)}(A; W, d); \\ I_C^{(4)}(A; W, d) &\leq I_C^{(3U)}(A; W, d) \leq I_C^{(2U)}(A; W, d) \leq I_C^{(1)}(A; W, d); \\ I_C^{(4)}(A; W, d) &\leq I_C^{(3)}(A; W, d) \leq I_C^{(2)}(A; W, d) \leq I_C^{(1)}(A; W, d); \\ S_C^{(1)}(A; W, d) &\leq S_C^{(2L)}(A; W, d) \leq S_C^{(3L)}(A; W, d) \leq S_C^{(4)}(A; W, d); \\ S_C^{(1)}(A; W, d) &\leq S_C^{(2U)}(A; W, d) \leq S_C^{(3U)}(A; W, d) \leq S_C^{(4)}(A; W, d); \\ S_C^{(1)}(A; W, d) &\leq S_C^{(2)}(A; W, d) \leq S_C^{(3)}(A; W, d) \leq S_C^{(4)}(A; W, d) \end{aligned}$$

for  $A, W \in \mathcal{P}(V) \setminus \{\emptyset\}$  and  $d \in C$ .

**Proposition 6** ([7]). *Let  $C$  be a convex cone in  $V$ . There are certain relations among the scalarizations of types (2L), (2U), (2) as well as (3L), (3U), (3):*



- (i)  $I_C^{(2)}(A; W, d) = \max \left\{ I_C^{(2L)}(A; W, d), I_C^{(2U)}(A; W, d) \right\};$
- (ii)  $I_C^{(3)}(A; W, d) = \max \left\{ I_C^{(3L)}(A; W, d), I_C^{(3U)}(A; W, d) \right\};$
- (iii)  $S_C^{(2)}(A; W, d) = \min \left\{ S_C^{(2L)}(A; W, d), S_C^{(2U)}(A; W, d) \right\};$
- (iv)  $S_C^{(3)}(A; W, d) = \min \left\{ S_C^{(3L)}(A; W, d), S_C^{(3U)}(A; W, d) \right\}$

for  $A, W \in \mathcal{P}(V) \setminus \{\emptyset\}$  and  $d \in C$ .

**Proposition 7** ([6]). *Let  $A, V \in \mathcal{P}(Y)$  and  $d \in Y$ . Then the following statements hold*

$$\begin{aligned}
-I_C^{(1)}(-A; -V, d) &= S_C^{(1)}(A; V, d), \\
-I_C^{(2L)}(-A; -V, d) &= S_C^{(2U)}(A; V, d), \\
-I_C^{(3L)}(-A; -V, d) &= S_C^{(3U)}(A; V, d), \\
-I_C^{(2U)}(-A; -V, d) &= S_C^{(2L)}(A; V, d), \\
-I_C^{(3U)}(-A; -V, d) &= S_C^{(3L)}(A; V, d), \\
-I_C^{(4)}(-A; -V, d) &= S_C^{(4)}(A; V, d).
\end{aligned}$$

For each  $j$  without  $j = 4$ , scalarizing functions  $I_C^{(j)}(\cdot; W, d)$  and  $S_C^{(j)}(\cdot; W, d)$  with a nonempty reference set  $W$  and a direction  $d$  have the following monotonicity with respect to  $\leq_C^{(j)}$ , which is referred to as “ $j$ -monotonicity” in [10]:

$$\begin{cases} A \leq_C^{(j)} B \implies I_C^{(j)}(A; W, d) \leq I_C^{(j)}(B; W, d); \\ A \leq_C^{(j)} B \implies S_C^{(j)}(A; W, d) \leq S_C^{(j)}(B; W, d). \end{cases} \quad (6)$$

## 4 Generalized cone-continuity

Let  $\mathcal{N}(x)$  and  $\preceq$  be a neighborhood system of a point  $x \in X$  and a binary relation on  $\mathcal{P}(Y)$ , respectively.

**Definition 8** (Definition 12 in [1]). Let  $F : X \rightarrow \mathcal{P}(Y)$ ,  $x_0 \in X$ ,  $\preceq$  a binary relation on  $\mathcal{P}(Y)$  and  $C \subset Y$  a convex cone. We say that  $F$  is  $(\preceq, C)$ -continuous at  $x_0$  if

$$\forall W \subset Y, W \text{ open}, W \preceq F(x_0), \exists V \in \mathcal{N}_X(x_0) \text{ s.t. } W + C \preceq F(x), \forall x \in V.$$

As special cases, for  $A, B \in \mathcal{P}(Y)$ , we consider binary relations  $\text{int } A \cap B \neq \emptyset$  and  $B \subset \text{int } A$  by  $A \preceq_1 B$  and  $A \preceq_2 B$ , respectively. Accordingly,  $(\preceq_1, C)$ -continuity and  $(\preceq_2, C)$ -continuity coincide with “ $C$ -lower continuity” and “ $C$ -upper continuity” for set-valued maps, respectively. Indeed,  $F : X \rightarrow \mathcal{P}(Y)$  is  $(\preceq_1, C)$ -continuous at  $x_0$  if and only if

$$\forall W \subset Y, W \text{ open}, W \cap F(x_0) \neq \emptyset, \exists V \in \mathcal{N}_X(x_0) \text{ s.t. } (W + C) \cap F(x) \neq \emptyset, \forall x \in V,$$

that is,  $F$  is  $C$ -lower continuous at  $x_0$ . Similarly,  $F$  is  $(\preceq_2, C)$ -continuous at  $x_0$  if and only if

$$\forall W \subset Y, W \text{ open}, F(x_0) \subset W, \exists V \in \mathcal{N}_X(x_0) \text{ s.t. } F(x) \subset W + C, \forall x \in V,$$

that is,  $F$  is  $C$ -upper continuous at  $x_0$ ; see Definition 2.5.16 of [4].

**Remark 9.** If  $C = \{0\}$  then  $(\preceq, C)$ -continuity for set-valued maps becomes  $\preceq$ -continuity in Definition 3.2 in [6]. Moreover,  $\preceq_1$ -continuity and  $\preceq_2$ -continuity coincide with the classical notions of lower continuity and upper continuity for set-valued maps, respectively.

**Definition 10** (Definition 14 in [1]). Let  $\varphi : \mathcal{P}(Y) \rightarrow \mathbb{R} \cup \{\pm\infty\}$ ,  $A_0 \in \mathcal{P}(Y)$ ,  $\preceq$  a binary relation on  $\mathcal{P}(Y)$ , and  $C$  a convex cone in  $Y$  with  $C \neq Y$ . Then, we say that  $\varphi$  is

- (i)  $(\preceq, C)$ -lower semicontinuous at  $A_0$  if  $\forall r < \varphi(A_0), \exists W \in \mathcal{P}(Y), W$  open, s.t.  $W \preceq A_0$  and  $r < \varphi(A), \forall A \in U(W + C, \preceq)$ ;
- (ii)  $(\preceq, C)$ -upper semicontinuous at  $A_0$  if  $\forall r > \varphi(A_0), \exists W \in \mathcal{P}(Y), W$  open, s.t.  $W \preceq A_0$  and  $r > \varphi(A), \forall A \in U(W + C, \preceq)$ ,

where  $U(V, \preceq) := \{A \in \mathcal{P}(Y) \mid V \preceq A\}$ .

**Remark 11.** When  $C = \{0\}$ ,  $(\preceq, C)$ -lower and  $(\preceq, C)$ -upper semicontinuities are coincident with  $\preceq$ -lower and  $\preceq$ -upper semicontinuities, respectively, which are introduced in Definition 3.3 of [6]. In Definition 10, we adopt that if  $\varphi(A_0) = -\infty$  (resp.  $+\infty$ ) then  $\varphi$  is  $(\preceq, C)$ -lower (resp. upper) semicontinuous at  $A_0$ .

Therefore, we can easily show the following results as generalizations of Theorems 3.1 and 3.2 in [6].

**Theorem 12** (Theorem 16 in [1]). *Let  $F : X \rightarrow \mathcal{P}(Y)$ ,  $\varphi : \mathcal{P}(Y) \rightarrow \mathbb{R} \cup \{\pm\infty\}$ ,  $x_0 \in X$ ,  $\preceq$  a binary relation on  $\mathcal{P}(Y)$ , and  $C \subset Y$  a convex cone. If  $F$  is  $(\preceq, C)$ -continuous at  $x_0$  and  $\varphi$  is  $(\preceq, C)$ -lower semicontinuous at  $F(x_0)$ , then  $\varphi \circ F$  is lower semicontinuous at  $x_0$ .*

**Theorem 13** (Theorem 17 in [1]). *Let  $F : X \rightarrow \mathcal{P}(Y)$ ,  $\varphi : \mathcal{P}(Y) \rightarrow \mathbb{R} \cup \{\pm\infty\}$ ,  $x_0 \in X$ ,  $\preceq$  a binary relation on  $\mathcal{P}(Y)$ , and  $C \subset Y$  a convex cone. If  $F$  is  $(\preceq, C)$ -continuous at  $x_0$  and  $\varphi$  is  $(\preceq, C)$ -upper semicontinuous at  $F(x_0)$ , then  $\varphi \circ F$  is upper semicontinuous at  $x_0$ .*

## 5 Continuity of Scalarization and Consequences

**Proposition 14.** *Let  $A_0, V \in \mathcal{P}(Y)$ ,  $C'$  and  $C$  two nonempty convex cones in  $Y$  such that  $C' \subset C \neq Y$ ,  $d \in \text{int} C$ . Then, the following statements hold:*

- (i)  $I_C^{(j)}(\cdot; V, d)$  is  $(\preceq_1, C')$ -lower semicontinuous at  $A_0$  for  $j = 1, 3U$ .
- (ii)  $S_C^{(j)}(\cdot; V, d)$  is  $(\preceq_1, C')$ -upper semicontinuous at  $A_0$  for  $j = 1, 3L$ .
- (iii)  $I_C^{(j)}(\cdot; V, d)$  is  $(\preceq_1, -C')$ -upper semicontinuous at  $A_0$  for  $j = 2L, 4$ .
- (iv)  $S_C^{(j)}(\cdot; V, d)$  is  $(\preceq_1, -C')$ -lower semicontinuous at  $A_0$  for  $j = 2U, 4$ .
- (v)  $I_C^{(j)}(\cdot; V, d)$  is  $(\preceq_2, -C')$ -upper semicontinuous at  $A_0$  for  $j = 1, 3U$ .
- (vi)  $S_C^{(j)}(\cdot; V, d)$  is  $(\preceq_2, -C')$ -lower semicontinuous at  $A_0$  for  $j = 1, 3L$ .
- (vii)  $I_C^{(j)}(\cdot; V, d)$  is  $(\preceq_2, C')$ -lower semicontinuous at  $A_0$  for  $j = 2L, 4$ .
- (viii)  $S_C^{(j)}(\cdot; V, d)$  is  $(\preceq_2, C')$ -upper semicontinuous at  $A_0$  for  $j = 2U, 4$ .

The following examples determine that  $I_C^{(j)}(\cdot; V, d)$  and  $S_C^{(j)}(\cdot; V, d)$  do not satisfy continuity for some  $j$ .

**Example 15.** Let  $Y = \mathbb{R}^2$ ,  $C = \mathbb{R}_+^2$ ,  $A_0 = \{(x, y) : x + y = 12 \text{ and } 2 \leq x \leq 10\}$ ,  $V = \{(x, y) : x + y = 6 \text{ and } 1 \leq x \leq 5\}$  and  $d = (1, 1)$ .

Then  $I_C^{(3L)}(\cdot; V, d)$  is not  $(\preceq_1, C')$ -lower semicontinuous at  $A_0$ . Additionally, this example can illustrate that  $I_C^{(3L)}(\cdot; V, d)$  is not  $(\preceq_1, -C')$ -lower semicontinuous at  $A_0$ . Besides,  $S_C^{(3U)}(\cdot; -V, d)$  is neither  $(\preceq_1, C')$ -upper semicontinuous nor  $(\preceq_1, -C')$ -upper semicontinuous at  $-A_0$ .

Moreover, we have  $I_C^{(3L)}(\cdot; V, d)$  is not  $(\preceq_1, C')$ -upper semicontinuous at  $A_0$ . Additionally, this example can illustrate that  $I_C^{(3L)}(\cdot; V, d)$  is not  $(\preceq_1, -C')$ -upper semicontinuous at  $A_0$ . Besides,  $S_C^{(3U)}(\cdot; -V, d)$  is neither  $(\preceq_1, C')$ -lower semicontinuous nor  $(\preceq_1, -C')$ -lower semicontinuous at  $-A_0$ .

**Example 16.** Let  $Y = \mathbb{R}^2$ ,  $C = \mathbb{R}_+^2$ ,  $A_0 = \{(x, y) : x + y = 12 \text{ and } 4 \leq x \leq 8\}$ ,  $V = \{(x, y) : x + y = 6 \text{ and } 1 \leq x \leq 5\}$  and  $d = (1, 1)$ .

Therefore  $I_C^{(2U)}(\cdot; V, d)$  is not  $(\preceq_1, C')$ -lower semicontinuous at  $A_0$ . Additionally, this example can illustrate that  $I_C^{(2U)}(\cdot; V, d)$  is not  $(\preceq_1, -C')$ -lower semicontinuous at  $A_0$ . Besides,  $S_C^{(2L)}(\cdot; -V, d)$  is neither  $(\preceq_1, C')$ -upper semicontinuous nor  $(\preceq_1, -C')$ -upper semicontinuous at  $-A_0$ .

**Example 17.** Let  $Y = \mathbb{R}^2$ ,  $C = \mathbb{R}_+^2$ ,  $A_0 = \{(x, y) : x + y = 12 \text{ and } 2 \leq x \leq 5\}$ ,  $V = \{(x, y) : x + y = 6 \text{ and } 1 \leq x \leq 5\}$  and  $d = (1, 1)$ . Therefore  $I_C^{(2U)}(\cdot; V, d)$  is not  $(\preceq_1, C')$ -upper semicontinuous at  $A_0$ . Additionally, this example can illustrate that  $I_C^{(2U)}(\cdot; V, d)$  is not  $(\preceq_1, -C')$ -upper semicontinuous at  $A_0$ . Besides,  $S_C^{(2L)}(\cdot; -V, d)$  is neither  $(\preceq_1, C')$ -lower semicontinuous nor  $(\preceq_1, -C')$ -lower semicontinuous at  $-A_0$ .

For  $I_C^{(2U)}(\cdot; V, d)$ ,  $S_C^{(2L)}(\cdot; V, d)$ ,  $I_C^{(3L)}(\cdot; V, d)$  and  $S_C^{(3U)}(\cdot; V, d)$ , the continuity properties are shown using the compactness assumptions.

**Proposition 18.** *Let  $A_0, V \in \mathcal{P}(Y)$ ,  $C'$  and  $C$  two nonempty convex cones such that  $C' \subset C \neq Y$  and  $d \in \text{int } C$ . Assume that  $A_0$  and  $V$  are compact. Then, the following statements hold:*

$I_C^{(2U)}(\cdot; V, d)$  is  $(\preceq_2, -C')$ -upper semicontinuous at  $A_0$ .

$S_C^{(2L)}(\cdot; V, d)$  is  $(\preceq_2, -C')$ -lower semicontinuous at  $A_0$ .

$I_C^{(3L)}(\cdot; V, d)$  is  $(\preceq_2, C')$ -lower semicontinuous at  $A_0$ .

$S_C^{(3U)}(\cdot; V, d)$  is  $(\preceq_2, C')$ -upper semicontinuous at  $A_0$ .

By Theorems 12 and 13, the following results are obtained.

**Theorem 19.** *Let  $F : X \rightarrow \mathcal{P}(Y)$ ,  $x_0 \in X$ ,  $V \in \mathcal{P}(Y)$ ,  $C'$  and  $C$  two nonempty convex cones in  $Y$  such that  $C' \subset C \neq Y$  and  $d \in \text{int } C$ . The following statements hold:*

- (a) *If  $F$  is  $(\preceq_1, C')$ -continuous at  $x_0$ , then*
  - (i)  $I_C^{(j)}(F(\cdot); V, d)$  is lower semicontinuous at  $x_0$ , for all  $j = 1, 3U$ ,
  - (ii)  $S_C^{(j)}(F(\cdot); V, d)$  is upper semicontinuous at  $x_0$ , for all  $j = 1, 3L$ .
- (b) *If  $F$  is  $(\preceq_1, -C')$ -continuous at  $x_0$ , then*
  - (i)  $I_C^{(j)}(F(\cdot); V, d)$  is upper semicontinuous at  $x_0$ , for all  $j = 2L, 4$ ,
  - (ii)  $S_C^{(j)}(F(\cdot); V, d)$  is lower semicontinuous at  $x_0$ , for all  $j = 2U, 4$ .
- (c) *If  $F$  is  $(\preceq_2, -C')$ -continuous at  $x_0$ , then*
  - (i)  $I_C^{(j)}(F(\cdot); V, d)$  is upper semicontinuous at  $x_0$ , for all  $j = 1, 3U$ ,
  - (ii)  $S_C^{(j)}(F(\cdot); V, d)$  is lower semicontinuous at  $x_0$ , for all  $j = 1, 3L$ .
- (d) *If  $F$  is  $(\preceq_2, C')$ -continuous at  $x_0$ , then*
  - (i)  $I_C^{(j)}(F(\cdot); V, d)$  is lower semicontinuous at  $x_0$ , for all  $j = 2L, 4$ ,
  - (ii)  $S_C^{(j)}(F(\cdot); V, d)$  is upper semicontinuous at  $x_0$ , for all  $j = 2U, 4$ .

Moreover, by Proposition 18, the consequent result can be implied by assuming compactness.

**Theorem 20.** *Let  $F : X \rightarrow \mathcal{P}(Y)$ ,  $x_0 \in X$ ,  $V \in \mathcal{P}(Y)$ ,  $C'$  and  $C$  two nonempty convex cones such that  $C' \subset C \neq Y$  and  $d \in \text{int } C$ . Assume that  $F(x_0)$  and  $V$  are compact.*

- (a) *If  $F$  is  $(\preceq_2, -C')$ -continuous at  $x_0$ , then*

- (i)  $I_C^{(2U)}(F(\cdot); V, d)$  is upper semicontinuous at  $x_0$ ,
- (ii)  $S_C^{(2L)}(F(\cdot); V, d)$  is lower semicontinuous at  $x_0$ ,
- (b) If  $F$  is  $(\preceq_2, C')$ -continuous at  $x_0$ , then
  - (i)  $I_C^{(3L)}(F(\cdot); V, d)$  is lower semicontinuous at  $x_0$ ,
  - (ii)  $S_C^{(3U)}(F(\cdot); V, d)$  is upper semicontinuous at  $x_0$ .

Finally, from the Theorems 19 and 20 together with Remark 11, we can summarize the previous results as follows.

Table 1: Continuity properties of the composite functions.

$F$	$(\preceq_1, C')$ -conti.	$(\preceq_2, C')$ -conti.	$(\preceq_1, -C')$ -conti.	$(\preceq_2, -C')$ -conti.	l.c. ( $C' = \{0\}$ )	u.c. ( $C' = \{0\}$ )
$I_C^{(1)} \circ F$	l.s.c.	-	-	u.s.c.	l.s.c.	u.s.c.
$I_C^{(2L)} \circ F$	-	l.s.c.	u.s.c.	-	u.s.c.	l.s.c.
$I_C^{(3L)} \circ F$	-	l.s.c. (*)	-	-	-	l.s.c. (*)
$I_C^{(2U)} \circ F$	-	-	-	u.s.c. (*)	-	u.s.c. (*)
$I_C^{(3U)} \circ F$	l.s.c.	-	-	u.s.c.	l.s.c.	u.s.c.
$I_C^{(4)} \circ F$	-	l.s.c.	u.s.c.	-	u.s.c.	l.s.c.
$S_C^{(1)} \circ F$	u.s.c.	-	-	l.s.c.	u.s.c.	l.s.c.
$S_C^{(2L)} \circ F$	-	-	-	l.s.c. (*)	-	l.s.c. (*)
$S_C^{(3L)} \circ F$	u.s.c.	-	-	l.s.c.	u.s.c.	l.s.c.
$S_C^{(2U)} \circ F$	-	u.s.c.	l.s.c.	-	l.s.c.	u.s.c.
$S_C^{(3U)} \circ F$	-	u.s.c. (*)	-	-	-	u.s.c. (*)
$S_C^{(4)} \circ F$	-	u.s.c.	l.s.c.	-	l.s.c.	u.s.c.

where (\*) means the compactness assumptions are required and “l.c.” and “u.c.” denote lower continuity and upper continuity of  $F$ , respectively.

## References

- [1] P. Dechboon, T. Tanaka, Some kinds of continuity properties on composite functions of set-valued maps and scalarizing functions, RIMS Kôkyûroku **2214** (2022), 76–85.
- [2] C. Gerstewitz (C. Tammer), Nichtkonvexe dualität in der vektoroptimierung, Wiss. Z. Tech. Hochsch. Leuna-Merseburg **25**(3) (1983), 357–364.
- [3] C. Gerth (Tammer), P. Weidner, Nonconvex separation theorems and some applications in vector optimization, J. Optim. Theory Appl., **67** (1990), 297–320.

- [4] A. Göpfert, H. Riahi, C. Tammer, C. Zălinescu, *Variational Methods in Partially Ordered Space*, Springer, New York, 2003.
- [5] A. H. Hamel, A. Löhne, Minimal element theorems and Ekeland's principle with set relations. *J. Nonlinear Convex Anal.* **7** (2006), 19–37.
- [6] K. Ike, M. Liu, Y. Ogata, T. Tanaka, Semicontinuity of the composition of set-valued map and scalarization function for sets, *J. Appl. Numer. Optim.*, **1** (2019), 267–276.
- [7] K. Ike, Y. Ogata, T. Tanaka, H. Yu, Sublinear-like scalarization scheme for sets and its applications to set-valued inequalities, In: A.A.Khan, E.Köbis, C.Tammer (eds.), *Variational Analysis and Set Optimization: Developments and Applications in Decision Making*, pp.91–113, CRC Press, Boca Raton, FL, 2019.
- [8] K. Ike, T. Tanaka, Convex-cone-based comparisons of and difference evaluations for fuzzy sets, *Optimization*, **67** (2018), 1051–1066.
- [9] A. A. Khan, C. Tammer, C. Zălinescu, *Set-valued Optimization*, Springer, Berlin, 2015.
- [10] S. Kobayashi, Y. Saito, T. Tanaka, Convexity for compositions of set-valued map and monotone scalarizing function, *Pacific J. Optim.* **12** (2016), 43–54.
- [11] E. Köbis, M. A. Köbis, Treatment of set order relations by means of a nonlinear scalarization functional: a full characterization, *Optimization* **65** (2016), 1805–1827.
- [12] D. Kuroiwa, T. Tanaka, T. X. D. Ha, On cone convexity of set-valued maps, *Nonlinear Anal.* **30** (1997), 1487–1496.
- [13] I. Kuwano, T. Tanaka, S. Yamada, Characterization of nonlinear scalarizing functions for set-valued maps, In: S.Akashi, W.Takahashi, T.Tanaka (eds.), *Proceedings of the Asian Conference on Nonlinear Analysis and Optimization*, pp.193–204, Yokohama Publishers, Yokohama, 2009.
- [14] I. Kuwano, T. Tanaka, S. Yamada, Inherited properties of nonlinear scalarizing functions for set-valued maps, In: Edited by S.Akashi, Y.Kimura, T.Tanaka (eds.), *Proceedings of the Sixth International Conference on Nonlinear Analysis and Convex Analysis*, pp.161–177, Yokohama Publishers, Yokohama, 2009.
- [15] I. Kuwano, T. Tanaka, S. Yamada, Unified scalarization for sets and set-valued Ky Fan minimax inequality, *J. Nonlinear Convex Anal.* **11** (2010), 513–525.

- [16] D.T. Luc, Theory of vector optimization, In: Lecture Notes in Economics and Mathematical Systems, vol. 319, Springer, Berlin, 1989.
- [17] S. Nishizawa, T. Tanaka, P. G. Georgiev, On inherited properties of set-valued maps, Proceedings of the Third International Conference on Nonlinear Analysis and Convex Analysis, Yokohama Publishers, Yokohama, 2003, 341–350.
- [18] Y. Ogata, T. Tanaka, Y. Saito, G. M. Lee, J. H. Lee, An alternative theorem for set-valued maps via set relations and its application to robustness of feasible sets, Optimization **67** (2018), 1067–1075.
- [19] Y. Saito, T. Tanaka, S. Yamada, On generalization of Ricceri’s theorem for Fan-Takahashi minimax inequality into set-valued maps via scalarization, J. Nonlinear Convex Anal. **16** (2015), 513–525.
- [20] Y. Sawaragi, H. Nakayama, T. Tanino, Theory of Multiobjective Optimization, Academic Press, Orlando, Florida, 1985.
- [21] Y. Sonda, I. Kuwano, T. Tanaka, Cone-semicontinuity of set-valued maps by analogy with real-valued semicontinuity, Nihonki Math. J., **21** (2010), 91–103.