

Two-microlocal analysis の新たな展開に向けて

奈良女子大学 森藤 紳哉

Shinya Moritoh
Nara Women's University

1 Introduction

This report is based on the author's talk given at RIMS on December 9th, 2019. Two-microlocal ideas in wavelet analysis are considered. Sections 2 and 3 are taken from [JM] and [MY], respectively. Section 4 deals with some recent results obtained in [Mo].

2 What is “two-microlocal estimate” ?

We first give a brief survey of Jaffard-Meyer (1996). See [JM]. The determination of the pointwise regularity of a function f requires the use of some tools introduced by Bony (1986). See [Bo].

Let S_j be the “low-pass filter” which, after performing the Fourier transform, is the multiplication by $\widehat{\varphi}(2^{-j}\xi)$, where $\widehat{\varphi}(\xi) = 1$ if $|\xi| \leq 1/2$ and $\widehat{\varphi}(\xi) = 0$ if $|\xi| \geq 1$. Define $\Delta_j = S_{j+1} - S_j$. Thus we have the Littlewood-Paley decomposition:

$$Id = S_0 + \Delta_0 + \Delta_1 + \cdots .$$

The Fourier transform of $\Delta_j(f)$ is supported by the set $2^{j-1} \leq |\xi| \leq 2^{j+1}$.

Definition 2.1 (Jaffard-Meyer). *Let $s, s' \in \mathbb{R}$. Then $f \in \mathcal{S}'(\mathbb{R}^n)$ is said to belong to $C_{x_0}^{s, s'}$ if*

$$|S_0(f)(x)| \leq C(1 + |x - x_0|)^{-s'}$$

and

$$|\Delta_j(f)(x)| \leq C2^{-js}(1 + 2^j|x - x_0|)^{-s'}$$

Definition 2.2 (Bony). *Let $s, s' \in \mathbb{R}$. Then $f \in \mathcal{S}'(\mathbb{R}^n)$ is said to belong to $H_{x_0}^{s, s'}$ if*

$$\|2^{js}(1 + 2^j|x - x_0|)^{s'} \Delta_j(f)\|_{L^2} \leq c_j$$

with $\sum |c_j|^2 < \infty$.

Remark 2.3. We have the following fact: $u \in H_{x_0}^{s, -k}$, with k being a positive integer, if and only if $u = \sum_{|\alpha| \leq k} (x - x_0)^\alpha u_\alpha$, where $u_\alpha \in H^{s-|\alpha|}(\mathbb{R}^n)$.

Let us now consider an orthonormal wavelet basis on \mathbb{R}^n . Such a basis is composed by translations and dilations of $2^n - 1$ functions $\psi^{(i)}$. Recall the usual notation

$$\psi_{j,k}^{(i)}(x) = 2^{nj/2} \psi^{(i)}(2^j x - k), \quad j \in \mathbb{Z}, \quad k \in \mathbb{Z}^n.$$

The wavelet decomposition of a function f will be written

$$f = \sum_{i,j,k} C_{j,k} 2^{nj/2} \psi^{(i)}(2^j x - k).$$

We will usually forget the index i . The following result is easy to check:

Proposition 2.4. $f \in \mathcal{S}'(\mathbb{R}^n)$ belongs to $C_{x_0}^{s,s'}$ if and only if

$$|C_{j,k}| \leq C 2^{-(s+n/2)j} (1 + 2^j |k 2^{-j} - x_0|)^{-s'}.$$

The following characterization also holds:

Proposition 2.5. $f \in \mathcal{S}'(\mathbb{R}^n)$ belongs to $H_{x_0}^{s,s'}$ if and only if

$$\sum_{j,k} 2^{2js} (1 + 2^j |k 2^{-j} - x_0|)^{2s'} |C_{j,k}|^2 < \infty.$$

Our next purpose is to characterize the two-microlocal spaces in terms of local ‘‘Hölder type’’ conditions. In order to state these conditions, we need the Hölder-Zygmund spaces $\dot{C}^s(\mathbb{R}^n)$. If $0 < s < 1$, then $f \in \dot{C}^s(\mathbb{R}^n)$ is characterized by

$$|f(x) - f(y)| \leq C|x - y|^s.$$

If $s = 1$, then $f \in \dot{C}^s(\mathbb{R}^n)$ is characterized by

$$|f(x+h) - 2f(x) + f(x-h)| \leq C|h|.$$

The definition of the case where $s > 1$ needs higher order differences and is omitted.

It is easily checked that $f \in \dot{C}^s(\mathbb{R}^n)$ if and only if its wavelet coefficients satisfy the condition

$$|C_{j,k}| \leq C 2^{-(s+n/2)j}.$$

Let $A \subset \mathbb{R}^n$. By definition, a function f belongs to $C^s(A)$ if it is the restriction to A of a function F in $\dot{C}^s(\mathbb{R}^n)$. The norm of f is then the infimum of all possible norms of F in $\dot{C}^s(\mathbb{R}^n)$. Let B_ρ be the ball $|x - x_0| \leq \rho$, and Γ_ρ the annulus $\rho \leq |x - x_0| \leq 3\rho$. The following characterizations are the starting point of the talk:

Theorem 2.6. If $s' < 0$, then $f \in \mathcal{S}'(\mathbb{R}^n)$ belongs to $C_{x_0}^{s,s'}$ if and only if

$$\|f|_{C^{s+s'}(B_\rho)}\| \leq C\rho^{-s'}. \quad (1)$$

If $s' > 0$, then $f \in \mathcal{S}'(\mathbb{R}^n)$ belongs to $C_{x_0}^{s,s'}$ if and only if $f \in \dot{C}^s(\mathbb{R}^n)$ and

$$\|f|_{C^{s+s'}(\Gamma_\rho)}\| \leq C\rho^{-s'}. \quad (2)$$

Proof. We assume that the wavelet ψ is compactly supported and that $0 \in \text{supp } \psi$. See [D]. We denote by C' the diameter of $\text{supp } \psi$. We first suppose that f belongs to $\dot{C}_{x_0}^{s,s'}$ so that its wavelet coefficients satisfy

$$|C_{j,k}| \leq C2^{-(s+n/2)j}(1 + |2^j x_0 - k|)^{-s'}. \quad (3)$$

Note first that if $s' > 0$, then (3) implies that $|C_{j,k}| \leq C2^{-(s+n/2)j}$, and so f belongs to $\dot{C}^s(\mathbb{R}^n)$. We split the wavelet decomposition

$$f = \sum C_{j,k} \psi_{j,k}$$

into three sums: $f = f_1 + f_2 + f_3$: The first, f_1 , corresponds to the wavelets whose supports do not intersect the ball B_ρ (or the annulus Γ_ρ), and we can forget this sum.

Next we consider the sum f_2 whose coefficients satisfy $2^j \rho \leq 10C'$; in that case, because $2^j |k2^{-j} - x_0|$ can be estimated from above by some constant comparable to $10C'$, (3) becomes

$$|C_{j,k}| \leq C2^{-(s+n/2)j},$$

and so $\|f_2 | \dot{C}^s(\mathbb{R}^n)\| \leq C$. The inequalities (1) and (2) for f_2 follow from this. (The details are omitted.)

Finally we consider the remaining sum f_3 whose coefficients satisfy $2^j \rho \geq 10C'$.

The case where $s' > 0$: (3) becomes

$$|C_{j,k}| \leq C2^{-(s+s'+n/2)j} \rho^{-s'},$$

because the supports of the wavelets are inside the annulus Γ_ρ so that $|x_0 - k2^{-j}| \geq \rho$. The corresponding sum f_3 satisfies

$$\|f_3 | \dot{C}^{s+s'}(\mathbb{R}^n)\| \leq C\rho^{-s'}.$$

The case where $s' < 0$: (3) implies that

$$|C_{j,k}| \leq C2^{-(s+n/2)j}(1 + 2^j \rho)^{-s'} \leq C2^{-(s+s'+n/2)j} \rho^{-s'}.$$

We have the same conclusion as before.

Conversely let us assume that (1) or (2) holds. We consider a given wavelet $\psi_{j,k}$.

The case where $s' < 0$: We take for ρ the smallest number such that the support of $\psi_{j,k}$ is completely included in B_ρ so that any function extending f outside B_ρ has the same wavelet coefficient $C_{j,k}$ and (1) implies that

$$2^{(s+s'+n/2)j} |C_{j,k}| \leq C\rho^{-s'}.$$

The case where $s' > 0$: Suppose first that $|x_0 - k2^{-j}| > 2C'2^{-j}$. Then the support of $\psi_{j,k}$ is completely included in Γ_ρ when $\rho = |x_0 - k2^{-j}|/2$ so that any function extending f outside Γ_ρ has the same wavelet coefficient $C_{j,k}$, and (2) implies that

$$2^{(s+s'+n/2)j} |C_{j,k}| \leq C\rho^{-s'}.$$

If $|x_0 - k2^{-j}| \leq 2C'2^{-j}$, then we have to prove that $|C_{j,k}| \leq C2^{-(s+n/2)j}$, which is implied by the assumption that $f \in \dot{C}^s(\mathbb{R}^n)$. \square

3 Two-microlocal Besov spaces and wavelets

Two-microlocal Besov spaces are considered by Moritoh-Yamada (2004), which is a natural extension of Jaffard-Meyer (1996). See [MY].

We first give the following definition and proposition. See [M] and [T].

Definition 3.1 (homogeneous Besov space). *Let $s > 0$ and $1 \leq p, q \leq \infty$. Then the homogeneous Besov $\dot{B}_{p,q}^s(\mathbb{R}^n)$ is defined as the set of all tempered distributions f (modulo polynomials) satisfying*

$$\|f | \dot{B}_{p,q}^s(\mathbb{R}^n)\| = \left(\sum_{j \in \mathbb{Z}} 2^{j\tilde{s}q} \|\mathcal{F}^{-1}(\varphi_j \mathcal{F}f) | L_p(\mathbb{R}^n)\|^q \right)^{1/q} < \infty.$$

Here, $\mathcal{F}f(\xi)$ denotes the Fourier transform of $f(x)$, and $\{\varphi_j\}_{j \in \mathbb{Z}}$ is a smooth resolution of unity.

Proposition 3.2. *$f \in \dot{B}_{p,q}^s(\mathbb{R}^n)$ if and only if*

$$\sum_{j \in \mathbb{Z}} 2^{j\tilde{s}q} \left(\sum_{k \in \mathbb{Z}^n} |C_{j,k}|^p \right)^{q/p} < \infty,$$

where $\tilde{s} = s + n/2 - n/p$.

We can define the local Besov spaces $B_{p,q}^s(U)$ by restriction (see the previous section), and we now give the definition of the two-microlocal Besov spaces $B_{p,q}^{s,s'}(U)$, where U is an open subset of \mathbb{R}^n .

Definition 3.3 (two-microlocal Besov space). *Let $s > 0$, $s' \in \mathbb{R}$ and $1 \leq p, q \leq \infty$. Then $f \in \mathcal{S}'(\mathbb{R}^n)$ is said to belong to the two-microlocal Besov space $B_{p,q}^{s,s'}(U)$ if the following two-microlocal estimate holds:*

$$\|f | B_{p,q}^{s,s'}(U)\| = \left[\sum_{j \in \mathbb{Z}} 2^{j\tilde{s}q} \left\{ \sum_{k \in \mathbb{Z}^n} \left| (1 + 2^j d(k2^{-j}, U))^{s'} C_{j,k} \right|^p \right\}^{\frac{q}{p}} \right]^{\frac{1}{q}} < \infty,$$

where $d(k2^{-j}, U)$ denotes the distance from $k2^{-j}$ to U .

In order to state the local Besov type conditions in our theorem below, we shall use the following notation as an analogue of Hörmander's notation [H]: If $g(\rho)$ is a function of the real variable ρ , defined for all positive ρ , we write $g(\rho) = \mathcal{O}^{(p)}(\rho^{-s})$ if and only if

$$\int_0^R (g(\rho)\rho^s)^p \frac{d\rho}{\rho} = \int_0^R g(\rho)^p \rho^{sp-1} d\rho < \infty \quad \text{for every } R > 0.$$

Theorem 3.4. *Let $s > 0$, $s' < 0$ and $1 \leq p \leq \infty$. Let U be an open subset in \mathbb{R}^n and $A_\rho = \{x \in \mathbb{R}^n; d(x, U) < \rho, x \notin U\}$. Then $f \in \mathcal{S}'(\mathbb{R}^n)$ belongs to $B_{p,p}^{s,s'}(U)$ if and only if there exists a decomposition $f = f_1 + f_2$ such that*

$$f_1 \in \dot{B}_{p,p}^s(\mathbb{R}^n),$$

and

$$\|f_2\|_{B_{p,p}^{s+s'}(A_\rho)} = \mathcal{O}^{(p)}(\rho^{-s'}).$$

Proof. We assume that the wavelet ψ is compactly supported and that $0 \in \text{supp } \psi$. We denote by C' the diameter of the support of the wavelet ψ . Let $f \in B_{p,p}^{s,s'}(U)$. Then its wavelet coefficients satisfy

$$\sum_{j \in \mathbb{Z}} 2^{j\tilde{s}p} \sum_{k \in \mathbb{Z}^n} \left| (1 + 2^j d(k2^{-j}, U))^{s'} C_{j,k} \right|^p < \infty. \quad (4)$$

We write f as

$$f = \sum_{\text{supp } \psi_{j,k} \cap U \neq \emptyset} C_{j,k} \psi_{j,k} + \sum_{\text{supp } \psi_{j,k} \cap U = \emptyset} C_{j,k} \psi_{j,k} =: f_1 + f_2.$$


If $\text{supp } \psi_{j,k} \cap U \neq \emptyset$, then $2^j d(2^{-j}k, U)$ is estimated from above by some constant comparable to C' . Therefore $f_1 \in \dot{B}_{p,p}^s(\mathbb{R}^n)$.

Next we split the wavelet decomposition of f_2 into three sums $f_2 = \sum_1 + \sum_2 + \sum_3$: Let $R > 0$ be fixed. The first, \sum_1 , corresponds to the wavelets whose supports do not intersect A_R , and we can forget this sum.

Next we consider the sum \sum_2 whose coefficients satisfy $2^j R \leq 10C'$; in that case, because $2^j d(2^{-j}k, U)$ can be estimated from above by some constant comparable to $10C'$, we have that $\sum_2 \in \dot{B}_{p,p}^s(\mathbb{R}^n)$.

Finally we consider the remaining sum \sum_3 whose coefficients satisfy $2^j R \geq 10C'$. We decompose A_R into the ‘‘curved annuli’’ as follows:

$$A_R = \bigcup_{m \in \mathbb{Z}; 2^{-m} \leq R} \{x \in \mathbb{R}^n; 2^{-m-1} \leq d(x, U) \leq 2^{-m}\} = \bigcup_{m; 2^m R \geq 1} D_m. \quad (5)$$



By using this decomposition (5), we can write (4) as follows:

$$\sum_{j; 2^j R \geq 10C'} 2^{j\tilde{s}p} \sum_{m; 2^m R \geq 1} (1 + 2^{j-m})^{s'p} \sum_{k; k2^{-j} \in D_m} |C_{j,k}|^p < \infty. \quad (6)$$

The case where $m > j + L(C')$, $L(C')$ being an integer dependent only on C' , is negligible because $\text{supp } \psi_{j,k} \cap U = \emptyset$. Therefore we obtain from (6) that

$$\begin{aligned} & \sum_{j; 2^j R \geq 10C'} \sum_{\substack{m; 2^m R \geq 1 \\ m \leq j + L(C')}} 2^{j\tilde{s}p} 2^{(j-m)s'p} \sum_{k; k2^{-j} \in D_m} |C_{j,k}|^p = \\ & = \sum_{m; 2^m R \geq 1} 2^{-ms'p} \sum_{\substack{j; 2^j R \geq 10C' \\ j \geq m - L(C')}} 2^{jp(\tilde{s}+s')} \sum_{k; k2^{-j} \in D_m} |C_{j,k}|^p < \infty. \end{aligned} \quad (7)$$

On the other hand, the $\mathcal{O}^{(p)}$ -condition that for every $R > 0$,

$$\int_0^R \left(\rho^{s'} \|f_2 | B_{p,p}^{s+s'}(A_\rho)\| \right)^p \frac{d\rho}{\rho} < \infty$$

follows from the condition that

$$\sum_{u \in \mathbb{Z}; 2^{-u} \leq R} 2^{-us'p} \sum_{j; 2^j R \geq 10C'} 2^{jp(\tilde{s}+s')} \sum_{v \in \mathbb{Z}; v \geq u} \sum_{k; k2^{-j} \in D_v} |C_{j,k}|^p < \infty. \quad (8)$$

Because $\text{supp } \psi_{j,k} \cap U = \emptyset$, and the geometric series $\sum_{u; u \leq v} 2^{-us'p}$ is estimated from above by some constant comparable to $2^{-vs'p}$ (note that $s' < 0$), this last condition (8) follows from that

$$\sum_{v; 2^v R \geq 1} 2^{-vs'p} \sum_{\substack{j; 2^j R \geq 10C' \\ j \geq v - L(C')}} 2^{jp(\tilde{s}+s')} \sum_{k; k2^{-j} \in D_v} |C_{j,k}|^p < \infty. \quad (9)$$

It follows from (7) and (9) that the remaining sum \sum_3 satisfies the local Besov $\mathcal{O}^{(p)}$ -condition, as desired.

Conversely let us assume that $f = f_1 + f_2$ satisfies the following conditions:

$$f_1 \in \dot{B}_{p,p}^s(\mathbb{R}^n), \quad (10)$$

and

$$\|f_2 | B_{p,p}^{s+s'}(A_\rho)\| = \mathcal{O}^{(p)}(\rho^{-s'}). \quad (11)$$

We note that if the support of the wavelet $\psi_{j,k}$ is completely included in A_ρ , then any function extending f_2 outside A_ρ has the same wavelet coefficient $C_{j,k}$. From this remark and (11), we have that for any $R > 0$,

$$\sum_{u; 2^u R \geq 1} 2^{-us'p} \sum_{j \in \mathbb{Z}} 2^{jp(\tilde{s}+s')} \sum_{k; k2^{-j} \in A_{2^{-u}}} |C_{j,k}|^p < \infty. \quad (12)$$

The condition (12) is equivalent to that

$$\sum_{j \in \mathbb{Z}} 2^{jp(\tilde{s}+s')} \sum_{k \in \mathbb{Z}^n} |C_{j,k}|^p \sum_{\substack{u; 2^u R \geq 1 \\ 2^u d(k2^{-j}, U) \leq 1}} 2^{-us'p} < \infty.$$

After the calculation of the geometric sum, we arrive at the following:

$$\sum_{j \in \mathbb{Z}} 2^{jp(\tilde{s}+s')} \sum_{k \in \mathbb{Z}^n} |C_{j,k}|^p \left(d(k2^{-j}, U)^{s'p} - R^{s'p} \right) < \infty. \quad (13)$$

Note that $s' < 0$. Then as $R \rightarrow \infty$ in (13), we obtain that

$$\sum_{j \in \mathbb{Z}} 2^{j\tilde{s}p} \sum_{k \in \mathbb{Z}^n} \left| (1 + 2^j d(k2^{-j}, U))^{s'} C_{j,k} \right|^p < \infty,$$

that is $f_2 \in B_{p,p}^{s,s'}(U)$. Taking into account the assumption (10) that $f_1 \in \dot{B}_{p,p}^s(\mathbb{R}^n)$, we conclude that $f = f_1 + f_2 \in B_{p,p}^{s,s'}(U)$. \square

4 Two-microlocal Besov spaces with dominating mixed smoothness

Moritoh (2016) considers “two-microlocal Besov spaces with dominating mixed smoothness” as a natural extension of Jaffard-Meyer (1996) and Moritoh-Yamada (2004) by taking account of uncertainty functions given by Weyl-Hörmander calculus (Bony-Lerner, 1989). See [Mo] and [BL].

We treat only the case where $n = 2$. Let us now consider an orthonormal wavelet basis on \mathbb{R}^2 composed by translations and dilations of $\psi(x_1)\psi(x_2)$, where $\psi(x)$ is a one-dimensional compactly supported smooth wavelet. Let $\psi_{j,k}(x) = 2^{j/2}\psi(2^jx - k)$ for $j \in \mathbb{Z}, k \in \mathbb{Z}$. Then every $f \in \mathcal{S}'(\mathbb{R}^2)$ will be written

$$f(x) = \sum_{\mathbf{j} \in \mathbb{Z}^2} \sum_{\mathbf{k} \in \mathbb{Z}^2} C_{\mathbf{j},\mathbf{k}} \psi_{j_1,k_1}(x_1)\psi_{j_2,k_2}(x_2),$$

where $\mathbf{j} = (j_1, j_2)$ and $\mathbf{k} = (k_1, k_2)$.

Let $s_1, s_2 > 0$ and $1 \leq p_1, p_2, q_1, q_2 \leq \infty$. Then the homogeneous Besov space with dominating mixed smoothness $S\dot{B}_{\mathbf{p},\mathbf{q}}^{\mathbf{s}}(\mathbb{R}^2)$ is defined as the set of all tempered distributions f (modulo polynomials) satisfying

$$\begin{aligned} & \|f|S\dot{B}_{\mathbf{p},\mathbf{q}}^{\mathbf{s}}(\mathbb{R}^2)\| \\ &= \left[\sum_{j_2 \in \mathbb{Z}} \left(\int_{\mathbb{R}} \left(\sum_{j_1 \in \mathbb{Z}} \left(\int_{\mathbb{R}} \left| 2^{j_1 s_1 + j_2 s_2} (\varphi_{j_1} \varphi_{j_2} \hat{f})^\vee(x_1, x_2) \right|^{p_1} dx_1 \right)^{q_1} dx_2 \right)^{p_2} \right)^{\frac{1}{q_2}} < \infty, \end{aligned}$$

where $\mathbf{s} = (s_1, s_2)$, $\mathbf{p} = (p_1, p_2)$, $\mathbf{q} = (q_1, q_2)$, and

$$(\varphi_{j_1} \varphi_{j_2} \hat{f})^\vee(x_1, x_2) = (\varphi_{j_1}(\xi_1) \varphi_{j_2}(\xi_2) \hat{f}(\xi_1, \xi_2))^\vee(x_1, x_2).$$

See Schmeisser-Triebel's book [ST].

Let us recall the fact that $f \in S\dot{B}_{\mathbf{p},\mathbf{q}}^{\mathbf{s}}(\mathbb{R}^2)$ if and only if

$$\left(\sum_{j_2 \in \mathbb{Z}} \left(\sum_{k_2 \in \mathbb{Z}} \left(\sum_{j_1 \in \mathbb{Z}} \left(\sum_{k_1 \in \mathbb{Z}} |2^{j_1 \tilde{s}_1 + j_2 \tilde{s}_2} C_{\mathbf{j},\mathbf{k}}|^{p_1} \right)^{\frac{q_1}{p_1}} \right)^{\frac{p_2}{q_1}} \right)^{\frac{q_2}{p_2}} \right)^{\frac{1}{q_2}} < \infty,$$

where $\tilde{s}_i = s_i + 1/2 - 1/p_i$ ($i = 1, 2$). See [B] and [V]. We treat only the case where $\mathbf{p} = \mathbf{q} = (p, p)$, $1 \leq p \leq \infty$. We can define the local Besov space $SB_{p,p}^{\mathbf{s}}(\mathbb{R}_{x_1} \times A_\rho)$ as usual, where $\mathbb{R}_{x_1} \times A_\rho$ denotes the horizontal strip $\{(x_1, x_2); x_1 \in \mathbb{R}, |x_2| < \rho\}$ for $\rho > 0$. We can also give the definition of the two-microlocal Besov space with dominating mixed smoothness $SB_{p,p}^{(s_1, s_2), s_3}(\mathbb{R}_{x_1} \times \{0\})$ as follows:

Definition 4.1. *Let $s_1, s_2 > 0$, $s_3 \in \mathbb{R}$, and $1 \leq p \leq \infty$. Then $f \in \mathcal{S}'(\mathbb{R}^2)$ is said to belong to the two-microlocal Besov space with dominating mixed smoothness $SB_{p,p}^{(s_1, s_2), s_3}(\mathbb{R}_{x_1} \times \{0\})$ if the following two-microlocal estimate holds:*

$$\begin{aligned} & \|f \mid SB_{p,p}^{(s_1, s_2), s_3}(\mathbb{R}_{x_1} \times \{0\})\| \\ & := \left[\sum_{\mathbf{j} \in \mathbb{Z}^2} \sum_{\mathbf{k} \in \mathbb{Z}^2} 2^{(j_1 \tilde{s}_1 + j_2 \tilde{s}_2)p} (1 + 2^{j_1} + (|k_2| + 1)2^{-j_2} 2^{j_1 \vee j_2})^{s_3 p} |C_{\mathbf{j},\mathbf{k}}|^p \right]^{\frac{1}{p}} < \infty, \end{aligned}$$

where $j_1 \vee j_2 = \max\{j_1, j_2\}$.


Our main theorem of this section is the following:

Theorem 4.2. *Let $s_i > 0$, $s_3 < 0$, $s_i + s_3 > 0$ ($i = 1, 2$), and $1 \leq p \leq \infty$. Then $f \in \mathcal{S}'(\mathbb{R}^2)$ belongs to $SB_{p,p}^{(s_1, s_2), s_3}(\mathbb{R}_{x_1} \times \{0\})$ if and only if there exists a decomposition $f = f_1 + f_2 + f_3 + f_4$ such that*

$$\begin{aligned} f_1 & \in S\dot{B}_{p,p}^{(s_1, s_2)}(\mathbb{R}^2), \quad f_2 \in S\dot{B}_{p,p}^{(s_1 + s_3, s_2)}(\mathbb{R}^2), \\ f_3 & \in S\dot{B}_{p,p}^{(s_1 + s_3, s_2 - s_3)}(\mathbb{R}^2), \end{aligned}$$

and

$$\|f_4 \mid SB_{p,p}^{(s_1, s_2 + s_3)}(\mathbb{R}_{x_1} \times A_\rho)\| = \mathcal{O}^{(p)}(\rho^{-s_3}).$$



Skech of the proof: We employ the method used in the proof of Theorem 4.2. Let $f \in SB_{p,p}^{(s_1,s_2),s_3}(\mathbb{R}_{x_1} \times \{0\})$. Then its wavelet coefficients satisfy

$$\sum_{\mathbf{j} \in \mathbb{Z}^2} \sum_{\mathbf{k} \in \mathbb{Z}^2} 2^{(j_1 \bar{s}_1 + j_2 \bar{s}_2)p} (1 + 2^{j_1} + (|k_2| + 1)2^{-j_2} 2^{j_1 \vee j_2})^{s_3 p} |C_{\mathbf{j}, \mathbf{k}}|^p < \infty. \quad (14)$$

We decompose f as follows:

$$f = f_1 + f_2,$$

where f_1 and f_2 correspond to the cases where $0 \in \text{supp } \psi_{j_2, k_2}$ and $0 \notin \text{supp } \psi_{j_2, k_2}$, respectively.


First: We decompose f_1 into three parts according to $\{j_1 > 0, j_2 > 0\}$, $\{j_2 < 0, j_1 > j_2\}$, and $\{j_1 < 0, j_2 > j_1\}$.

Second: We decompose f_2 into three parts, among which the case where $2^{j_2} R \geq 10C'$ is the most important.

Third: We decompose this important term into three parts according to $\{j_1 < 0, j_2 < m\}$, $\{j_1 > 0, j_2 < j_1 + m\}$, and $\{j_2 > m, j_2 > j_1 + m\}$. The last term yields the function f_4 characterized by the local Besov type condition with dominating mixed smoothness.

Summing up, the case where $2^{j_2} R \geq 10C'$ (R is a fixed positive number), $j_2 > m$, $j_2 > j_1 + m$ ($m > -\log_2 R$) in the wavelet decomposition of f yields the most singular part f_4 .


We finally remark that the case where $j_1 > j_2$ and $j_2 < 0$ in the wavelet decomposition of f_1 yields the function $f_3 \in \dot{SB}_{p,p}^{(s_1+s_3, s_2-s_3)}(\mathbb{R}^2)$.



Remark 4.3. The idea of this theorem is that every f belonging to the generalized function space $SB_{p,p}^{(s_1,s_2),s_3}(\mathbb{R}_{x_1} \times \{0\})$ has a good decomposition $f = \sum_{i=1}^4 f_i$, where the term f_4 represents the singularities of the function f along the line \mathbb{R}_{x_1} ; they satisfy the local Besov type conditions in the neighborhood of the x_1 -axis. (As we have seen in section 2, every $f \in B_{p,p}^{s,s'}(x_0)$ has a good decomposition $f = f_1 + f_2$, where the term f_2 represents the singularities of the function f at the point x_0 .) Our future research is a more complete theory of two-microlocal spaces using Weyl-Hörmander calculus.

Remark 4.4. The typical examples considered by Jaffard-Meyer are an indefinitely oscillating function of the form $x^\alpha \sin(1/x^\beta)$, and Riemann's nondifferentiable function $\sigma(x) = \sum_{n=1}^{\infty} (1/n^2) \sin(\pi n^2 x)$, where the Hölder regularity at a point x_0 depends on the Diophantine approximation properties of x_0 . Higher dimensional singularities will be studied in our future research.

Remark 4.5. The two-microlocal Besov spaces of product type are easily introduced and characterized. It is associated with the uncertainty functions $\lambda_i = 1 + |x_i| |\xi_i|$ ($i = 1, 2$); the norm of the wavelet coefficients $C_{\mathbf{j},\mathbf{k}}$ is defined by means of the weighted coefficients $2^{(j_1 \tilde{s}_1 + j_2 \tilde{s}_2)} (1 + |k_1|)^{s'_1} (1 + |k_2|)^{s'_2} |C_{\mathbf{j},\mathbf{k}}|$.



$$\mathbb{R}_{x_1} \times A_{\rho_2} \text{ and } A_{\rho_1} \times \mathbb{R}_{x_2}$$

References

- [B] Bazarkhanov, D. B., *Wavelet representations and equivalent normings for some function spaces of generalized mixed smoothness*, (Russian), Mat. Zh. **5** (2005), no. 2 (16), 12–16.
- [Bo] Bony, J.-M., *Second microlocalization and propagation of singularities for semilinear hyperbolic equations*, Hyperbolic equations and related topics (Katata/Kyoto, 1984), 11–49, Academic Press, Boston, MA, 1986.
- [BL] Bony, J.-M. and Lerner, N., *Quantification asymptotique et microlocalisations d'ordre supérieur. I*, Ann. Sci. École Norm. Sup. (4) **22** (1989), no. 3, 377–433.
- [D] Daubechies, I., *Orthonormal bases of compactly supported wavelets*, Comm. Pure Appl. Math. **41** (1988), no. 7, 909–996.

- [H] Hörmander, L., *On interior regularity of the solutions of partial differential equations*, Comm. Pure Appl. Math. **11** (1958), 197–218.
- [JM] Jaffard, S. and Meyer, Y., Wavelet methods for pointwise regularity and local oscillations of functions, Mem. Amer. Math. Soc. **123** (1996), no. 587.
- [M] Meyer, Y., *Ondelettes et Opérateurs, I*, Actualités Mathématiques, Hermann, Paris, 1990.
- [Mo] Moritoh, S., *Detection of singularities in wavelet and ridgelet analyses*, RIMS Kôkyûroku Bessatsu **B57** (2016), 1–13.
- [MY] Moritoh, S. and Yamada, T., *Two-microlocal Besov spaces and wavelets*, Rev. Mat. Iberoamericana **20** (2004), no. 1, 277–283.
- [ST] Schmeisser, H.-J. and Triebel, H., *Topics in Fourier analysis and function spaces*, A Wiley-Interscience Publication, John Wiley & Sons, Ltd., Chichester, 1987.
- [T] Triebel, H., *Theory of Function Spaces*, Monographs in Mathematics **78**, Birkhäuser, Basel, 1983.
- [V] Vybiral, J., *Function spaces with dominating mixed smoothness*, Dissertationes Math. (Rozprawy Mat.) **436** (2006).