

Spectral projective schemes and their applications

Mariko Ohara (Joint work with Takeshi Torii)
Oshima College

Acknowledgement. I would like to appreciate Professor Maki Nakasuji and her Research Center Project "Zeta functions and their representations". This work was supported by the Research Institute for Mathematical Sciences, an International Joint Usage / Research Center located in Kyoto University.

An \mathbb{E}_∞ -ring is a spectrum equipped with commutative multiplicative law up to coherent homotopy.

Lurie refined the notation of \mathbb{E}_∞ -ring by using ∞ -operad $N_\Delta(\mathcal{F}in_*)$. In 2001, Mandell, May, Schwede and Shipley considered diagram spectra, especially \mathbb{Z} -indexed ring spectra. Lurie consider \mathbb{Z} -grading on \mathbb{E}_∞ -ring by using \mathbb{Z} with order preserving morphisms. I think some professionals already noticed that the symmetric monoidal structure on discrete graded rings and modules are obtained by the Day convolution.

We have symmetric monoidal ∞ -categories $N_\Delta(\mathcal{O}_{\mathbb{N}}^{\otimes}) \rightarrow N_\Delta(\mathcal{F}in_*)$ and $N_\Delta(\mathcal{O}_{\mathbb{Z}}^{\otimes}) \rightarrow N_\Delta(\mathcal{F}in_*)$. By using these symmetric monoidal ∞ -categories, we concretely define graded \mathbb{E}_∞ -rings and graded modules over them, and study their properties. We construct projective schemes associated to connective \mathbb{N} -graded \mathbb{E}_∞ -rings in spectral algebraic geometry. Actually, Torii and I gave these construction in more general situation, please see [8]. In this paper, I use an expedient ; I state their properties by using ∞ -operads.

Under some finiteness conditions, we show that the ∞ -category of almost perfect quasi-coherent sheaves over a spectral projective scheme $\text{Proj}(A)$ associated to a connective \mathbb{N} -graded \mathbb{E}_∞ -ring A can be described in terms of \mathbb{Z} -graded A -modules.

1. GRADED \mathbb{E}_∞ -RINGS AND GRADED MODULES

For a symmetric monoidal category consisting of one object $\{0\}$ with the unique isomorphism, we have a category $\mathcal{F}in_*$. The map $\{0\} \rightarrow \mathbb{N}$ and $\{0\} \rightarrow \mathbb{Z}$ are the symmetric monoidal functors.

Definition 1.1. Let $\mathcal{C}^{\otimes} \rightarrow N_\Delta(\mathcal{F}in_*)$ be a symmetric monoidal ∞ -category. We obtain a symmetric monoidal ∞ -category $\text{Fun}_{N_\Delta(\mathcal{F}in_*)}(N_\Delta(\mathbb{Z}), \mathcal{C})^{\otimes} \rightarrow N_\Delta(\mathcal{F}in_*)$ by the Day convolution which is denoted by $\hat{\otimes}$.

We use the notation $\text{Alg}_{\mathcal{O}'/\mathcal{O}}(\mathcal{M})$ for the fibration $\mathcal{M} \rightarrow \mathcal{O}$ of ∞ -operads and given ∞ -operads $\mathcal{O}' \rightarrow \mathcal{O}$.

For $X \in \text{Fun}_{N_\Delta(\mathcal{F}in_*)}(N_\Delta(\mathbb{Z}), \mathcal{C})^{\hat{\otimes}}$, X_i is the value at $i \in \mathbb{Z}$ of the underlying functor of X .

We take the symmetric monoidal ∞ -categories Sp^{\otimes} and Mod_R^{\otimes} , where R is an \mathbb{E}_∞ -ring.

Key diagram. Consider the following diagram

$$\begin{array}{ccccc} N_\Delta(\mathcal{F}in_*) & \xleftarrow{i} & N_\Delta(\mathcal{O}_{\mathbb{Z}}^{\otimes}) & \xleftarrow{0} & N_\Delta(\mathcal{F}in_*) \\ & \searrow A(\infty) & \downarrow A & \swarrow A_0 & \\ & & \text{Sp}^{\otimes} & & \end{array},$$

where the map 0 is induced from $\{0\} \rightarrow \mathbb{Z}$, the map i is the structure map of symmetric monoidal structure and $A(\infty)$ is the operadic left Kan extensions of A along i . Then,

- (i) A_0 is an \mathbb{E}_∞ -ring, i.e., A_0 is an object in $\text{Alg}_{N_\Delta(\mathcal{F}in_*)/N_\Delta(\mathcal{F}in_*)}(\text{Sp}^{\otimes})$,
- (ii) $A(\infty)$ is an \mathbb{E}_∞ -ring, i.e., an object in $\text{Alg}_{N_\Delta(\mathcal{F}in_*)/N_\Delta(\mathcal{F}in_*)}(\text{Sp}^{\otimes})$.

Especially, the functor $(-)_0$ commutes with the graded tensor. We call $A(\infty)$ the underlying \mathbb{E}_∞ -ring of A . Roughly speaking, $A(\infty)$ is such as a form of direct sum with respect to grading.

Definition 1.2. • We define the ∞ -category of \mathbb{Z} -graded \mathbb{E}_∞ -rings by

$$\text{Alg}_{N_\Delta(\mathcal{F}in_*)/N_\Delta(\mathcal{F}in_*)}(\text{Fun}_{N_\Delta(\mathcal{F}in_*)}(N_\Delta(\mathbb{Z}), \text{Sp})^{\hat{\otimes}}),$$

and call its object a \mathbb{Z} -graded \mathbb{E}_∞ -rings.

- We say that a \mathbb{Z} -graded \mathbb{E}_∞ -ring A is connective if each A_i for $i \in \mathbb{Z}$ is a connective spectrum.
- Let R be an \mathbb{E}_∞ -ring. We define the ∞ -category of \mathbb{Z} -graded \mathbb{E}_∞ -rings over R by

$$\text{Alg}_{N_\Delta(\mathcal{F}in_*)/N_\Delta(\mathcal{F}in_*)}(\text{Fun}_{N_\Delta(\mathcal{F}in_*)}(N_\Delta(\mathbb{Z}), \text{Mod}_R)^{\hat{\otimes}}),$$

and call its object a \mathbb{Z} -graded \mathbb{E}_∞ -rings over R .

We denote by $\text{CAlg}_R(\mathbb{Z})$ and $\text{CAlg}_R(\mathbb{N})$ the ∞ -category of \mathbb{Z} -graded and \mathbb{N} -graded \mathbb{E}_∞ -rings over R , respectively. We identify objects of $\text{CAlg}(\mathbb{N})$ with that of $\text{CAlg}(\mathbb{Z})$.

1.1. Modules over graded \mathbb{E}_∞ -rings.

Definition 1.3. For a \mathbb{Z} -graded \mathbb{E}_∞ -ring A and an \mathbb{E}_∞ -ring R , the ∞ -category of \mathbb{Z} -graded A -modules is

$$\text{Mod}_A(\text{Fun}_{N_\Delta(\mathcal{F}in_*)}(N_\Delta(\mathbb{Z}), \text{Sp})^{\hat{\otimes}}),$$

where the notation $\text{Mod}_A(-)$ is in the sense of Lurie. Let us denote the ∞ -category of \mathbb{Z} -graded A -modules by $\text{Mod}_A(\mathbb{Z})$.

We call a morphism in $\mathrm{CAlg}(\mathbb{Z})$ and $\mathrm{Mod}_A(\mathbb{Z})$ a morphism of degree 0 or a morphism of graded \mathbb{E}_∞ -rings and of graded A -modules.

Remark 1.4. The ∞ -category of \mathbb{Z} -graded A -modules over R can be defined as

$$\mathrm{Mod}_A(\mathrm{Fun}_{N_\Delta(\mathcal{F}\mathrm{in}_*)}(N_\Delta(\mathbb{Z}), \mathrm{Mod}_R)^{\hat{\otimes}}).$$

1.2. Localizations of graded \mathbb{E}_∞ -rings. For X in $\mathrm{Sp}(\mathbb{Z})$ and $g \in \mathbb{Z}$, we define a twisting $X(g)$ in $\mathrm{Sp}(\mathbb{Z})$ by $X(g)_{g'} \simeq X_{g+g'}$ for $g' \in \mathbb{Z}$.

Let A be a \mathbb{Z} -graded \mathbb{E}_∞ -ring and let $a \in \pi_0(A)$ be homogeneous of degree $g \in \mathbb{Z}$. We regard a as a morphism $a : A \rightarrow A(g)$ of \mathbb{Z} -graded A -modules. Since $\mathrm{Mod}_A(\mathbb{Z})$ is a presentable ∞ -category, there exists a localization functor

$$L : \mathrm{Mod}_A(\mathbb{Z}) \longrightarrow \mathrm{Mod}_A(\mathbb{Z})$$

with respect to the map $a : A \rightarrow A(g)$. As in the nongraded case, $L(M)$ is equivalent to $M[a^{-1}]$, where $M[a^{-1}]$ is a colimit of the sequence

$$M \xrightarrow{a} M(g) \xrightarrow{a} M(2g) \xrightarrow{a} \dots$$

in $\mathrm{Mod}_A(\mathbb{Z})$. The localization L is smashing given by $L(-) \simeq A[a^{-1}] \otimes_A (-)$ and is compatible with the symmetric monoidal structure on $\mathrm{Mod}_A(\mathbb{Z})$. We can regard $l : A \rightarrow A[a^{-1}]$ a morphism in $\mathrm{CAlg}(\mathbb{Z})$. We obtain an adjunction

$$l_! : \mathrm{Mod}_A(\mathbb{Z}) \rightleftarrows \mathrm{Mod}_{A[a^{-1}]}(\mathbb{Z}) : l^*,$$

where the left adjoint $l_!$ is a symmetric monoidal functor given by $M \mapsto A[a^{-1}] \otimes_A M$, and the right adjoint l^* is a fully faithful lax symmetric monoidal functor.

By [3, Remark 7.3.2.13], this adjunction induces an adjunction

$$l_! : \mathrm{CAlg}_A \rightleftarrows \mathrm{CAlg}_{A[a^{-1}]} : l^*,$$

where the right adjoint l^* is fully faithful. Hence $l_! : \mathrm{CAlg}_A(\mathbb{Z}) \rightarrow \mathrm{CAlg}_{A[a^{-1}]}(\mathbb{Z})$ is a localization functor.

If a is invertible in $\pi_0(B)$, we have an equivalence

$$\mathrm{Map}_{\mathrm{CAlg}_A(\mathbb{Z})}(A[a^{-1}], B) \xrightarrow{\simeq} \mathrm{Map}_{\mathrm{CAlg}_A(\mathbb{Z})}(A, B)$$

since there is an equivalence $l_!A \simeq A[a^{-1}]$ in $\mathrm{CAlg}_A(\mathbb{Z})$.

We define an ∞ -category

$$\mathrm{CAlg}_A^{\mathrm{Zar}}(\mathbb{Z})$$

to be the full subcategory of $\mathrm{CAlg}_A(\mathbb{Z})$ spanned by those objects of the form $A[a^{-1}]$ for some homogeneous element $a \in \pi_0(A)$. We define $\mathrm{CAlg}_{\pi_0(A)}^{\heartsuit\mathrm{Zar}}(\mathbb{Z})$ to be the full subcategory of $\mathrm{CRing}_{\pi_0(A)}(\mathbb{Z})$ spanned by those objects of the form $\pi_0(A)[a^{-1}]$ for some $a \in \pi_0(A)$.

Definition 1.5. A spectral scheme X is a projective spectral scheme if there exists a collection $\{U_a\}$ such that U_a covers X and there exists $A \in \text{CAlg}(\mathbb{Z})$ such that $(U_a, \mathcal{O}_X) \simeq (\text{Spec}(A[\alpha_a^{-1}]_0), \mathcal{O}_{\text{Spec}(A[\alpha_a^{-1}]_0)})$ for each U_a and for degree (more than) 1 elements $\alpha_a \in \pi_0(A(\infty))$.

2. QUASI-COHERENT SHEAVES ON PROJECTIVE SPECTRAL SCHEMES

Definition 2.1. Let A be a connective \mathbb{N} -graded \mathbb{E}_∞ -ring. We say that A is Noetherian if $\pi_0(A)$ is a Noetherian \mathbb{N} -graded commutative ring and $\pi_n(A)$ is a finitely generated \mathbb{N} -graded $\pi_0(A)$ -module for any $n \in \mathbb{Z}$.

Let A be a connective Noetherian \mathbb{N} -graded \mathbb{E}_∞ -ring. In this section we assume that A satisfies the following condition:

There are finitely many elements of $\pi_0(A_1)$ which generate $\pi_0(A)$ as an \mathbb{N} -graded commutative ring over $\pi_0(A_0)$.

Let A be a connective Noetherian \mathbb{N} -graded \mathbb{E}_∞ -ring. Set $X = \text{Proj}(A)$. We take a set $\{a_i\}_{i=1}^r \subset \pi_0(A_1)$ of generators of $\pi_0(A)$ as an \mathbb{N} -graded commutative ring over $\pi_0(A_0)$. We define a \mathbb{Z} -graded \mathbb{E}_∞ -ring B by $B = A[a_1^{-1}] \times \cdots \times A[a_r^{-1}]$. Let $g : A \rightarrow B$ be the canonical morphism of \mathbb{Z} -graded \mathbb{E}_∞ -rings. We take a Čech nerve

$$C(g)_+^\bullet$$

of g in the opposite ∞ -category of $\text{CAlg}(\mathbb{Z})$. Then $C(g)_+^\bullet$ is an augmented cosimplicial object of $\text{CAlg}(\mathbb{Z})$ such that $C(g)_+^{-1} \simeq A$ and $C(g)_+^n \simeq B^n$ for $n \geq 0$, where B^n is given by

$$B^n = \overbrace{B \otimes_A \cdots \otimes_A B}^{n+1}.$$

By using the functor $(-)_0 : \text{CAlg}(\mathbb{Z}) \rightarrow \text{CAlg}$, we obtain $C(g)_0^\bullet$ as the composite of $(-)_0$ with the restriction $C(g)_+^\bullet = C(g)_+^\bullet|_\Delta$.

Note that there is a faithfully flat affine morphism $f : U \rightarrow X$, where $U = \text{Spec}(B_0)$, and we have an equivalence $U_\bullet \simeq \text{Spec}(C(g)_0^\bullet)$ of simplicial objects of affine spectral schemes. Since $f : U \rightarrow X$ is an effective epimorphism, we have an equivalence

$$|U_\bullet| \xrightarrow{\simeq} X$$

in $\widehat{\text{Shv}}_{\text{fpqc}}$, where the left hand side is the geometric realization of the simplicial object U_\bullet .

We have an ∞ -category $\text{QCoh}(X)$ of quasi-coherent sheaves of \mathcal{O}_X -modules on X , which is stable presentable symmetric monoidal with unit \mathcal{O}_X by [5, Proposition 2.2.4.2].

There is an equivalence

$$\text{QCoh}(X) \simeq \lim_{\Delta} \text{Mod}_{B_0^\bullet}$$

of symmetric monoidal stable ∞ -categories.

Recall that $\mathrm{Sp}(\mathbb{Z})$ is a symmetric monoidal stable presentable ∞ -category, so that it is a commutative algebra object of $\mathcal{P}r_{\mathrm{St}}^{\mathrm{L}}$, where $\mathcal{P}r_{\mathrm{St}}^{\mathrm{L}}$ is the ∞ -category of stable presentable ∞ -categories and colimit-preserving functors. We denote by $\mathrm{CAlg}(\mathcal{P}r_{\mathrm{St}}^{\mathrm{L}})$ the ∞ -category of commutative algebra objects of $\mathcal{P}r_{\mathrm{St}}^{\mathrm{L}}$. By [3, Theorem 4.8.5.16], we have a functor

$$\mathrm{Mod}_{(-)}(\mathbb{Z}) : \mathrm{CAlg}(\mathbb{Z}) \longrightarrow \mathrm{CAlg}(\mathcal{P}r_{\mathrm{St}}^{\mathrm{L}}),$$

which assigns to $D \in \mathrm{CAlg}(\mathbb{Z})$ the symmetric monoidal ∞ -category $\mathrm{Mod}_D(\mathbb{Z})$ of \mathbb{Z} -graded D -modules.

By applying the functor $\mathrm{Mod}_{(-)}(\mathbb{Z})$ to $C(g)_+^{\bullet}$ and using the equivalences $\mathrm{Mod}_{B^n}(\mathbb{Z}) \simeq \mathrm{Mod}_{B_0^n}$ for $n \geq 0$, we obtain a symmetric monoidal functor

$$\mathrm{Mod}_A(\mathbb{Z}) \longrightarrow \lim_{\Delta} \mathrm{Mod}_{B_0^{\bullet}}.$$

Definition 2.2. We define a functor

$$\widetilde{(-)} : \mathrm{Mod}_A(\mathbb{Z}) \longrightarrow \mathrm{QCoh}(X)$$

to be the composite of the functor $\mathrm{Mod}_A(\mathbb{Z}) \rightarrow \lim_{\Delta} \mathrm{Mod}_{B_0^{\bullet}}$ with the equivalence between $\lim_{\Delta} \mathrm{Mod}_{B_0^{\bullet}}$ and $\mathrm{QCoh}(X)$. We call \widetilde{M} the quasi-coherent sheaf on X associated to a \mathbb{Z} -graded A -modules M .

By the construction, the functor $\widetilde{(-)} : \mathrm{Mod}_A(\mathbb{Z}) \rightarrow \mathrm{QCoh}(X)$ is symmetric monoidal.

Recall that we have defined the shifting functor $(q) : \mathrm{Mod}_A(\mathbb{Z}) \rightarrow \mathrm{Mod}_A(\mathbb{Z})$ given by $M(q)_n \simeq M_{q+n}$ for $M \in \mathrm{Mod}_A(\mathbb{Z})$ and $q, n \in \mathbb{Z}$. For $q \in \mathbb{Z}$, we define $\mathcal{O}_X(q)$ to be the quasi-coherent sheaf $\widetilde{A(q)}$ on X .

Proposition 2.3. *The quasi-coherent sheaf $\mathcal{O}_X(q)$ is locally free of rank 1 for any $q \in \mathbb{Z}$.*

Proof. It suffices to show that the restriction $\widetilde{A(q)}|_V$ is equivalent to $\widetilde{A}|_V$ for any affine open set $V = \mathrm{Spec}(\pi_0(A)[f^{-1}]_0)$ of the underlying projective scheme $\mathrm{Proj}(\pi_0(A))$, where f is an element of $\pi_0(A)$ of degree 1. The restriction $\widetilde{A(q)}|_V$ corresponds to an $A[f^{-1}]_0$ -module $A[f^{-1}]_q$. The multiplication by f^q induces an equivalence of $A[f^{-1}]_0$ -modules between $A[f^{-1}]_q$ and $A[f^{-1}]_0$. Thus, there is an equivalence of quasi-coherent sheaves between $\widetilde{A(q)}|_V$ and $\widetilde{A}|_V$. This completes the proof. \square

For a quasi-coherent sheaf \mathcal{F} of \mathcal{O}_X -modules on X and $q \in \mathbb{Z}$, we define $\mathcal{F}(q) = \mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{O}_X(q)$.

If \mathcal{F} is the quasi-coherent sheaf associated to a \mathbb{Z} -graded A -module M , then we have an equivalence $\mathcal{F}(q) \simeq \widetilde{M(q)}$ of quasi-coherent sheaves.

The functor $\widetilde{(-)}$ is obtained from the augmented cosimplicial diagram $\mathrm{Mod}_{(-)}(\mathbb{Z}) \circ C(g)_+^{\bullet} : \Delta_+ \rightarrow \mathrm{CAlg}(\mathbb{Z}) \rightarrow \widehat{\mathrm{Cat}}_{\infty}$. Since the functor $\mathrm{Mod}_{(-)}(\mathbb{Z}) : \mathrm{CAlg}(\mathbb{Z}) \rightarrow \widehat{\mathrm{Cat}}_{\infty}$ factors

through the ∞ -category $\mathcal{P}r_{\text{St}}^{\text{L}}$ of stable presentable ∞ -categories and colimit-preserving functors, we see that the functor $\widetilde{(-)} : \text{Mod}_A(\mathbb{Z}) \rightarrow \text{QCoh}(X)$ is a morphism in $\mathcal{P}r_{\text{St}}^{\text{L}}$.

Thus, there exists a right adjoint $\Gamma_*(X, -) : \text{QCoh}(X) \rightarrow \text{Mod}_A(\mathbb{Z})$ to $\widetilde{(-)}$.

The equivalence $\text{QCoh}(X) \xrightarrow{\simeq} \lim_{\Delta} \text{Mod}_{B^\bullet}(\mathbb{Z})$ of ∞ -categories implies an equivalence

$$\text{Map}_{\text{QCoh}(X)}(\widetilde{M}, \mathcal{F}) \simeq \lim_{\Delta} \text{Map}_{\text{Mod}_{B^\bullet}(\mathbb{Z})}(B^\bullet \otimes_A M, \Gamma(U_\bullet, \mathcal{F}(*)))$$

of mapping spaces. Since we have an equivalence

$$\text{Map}_{\text{Mod}_{B^\bullet}(\mathbb{Z})}(B^\bullet \otimes_A M, \Gamma(U_\bullet, \mathcal{F}(*))) \simeq \text{Map}_{\text{Mod}_A(\mathbb{Z})}(M, \Gamma(U_\bullet, \mathcal{F}(*)))$$

of cosimplicial spaces, there is a natural equivalence

$$\text{Map}_{\text{QCoh}(X)}(\widetilde{M}, \mathcal{F}) \simeq \text{Map}_{\text{Mod}_A(\mathbb{Z})}(M, \lim_{\Delta} \Gamma(U_\bullet, \mathcal{F}(*))).$$

Hence we obtain $\Gamma_*(X, \mathcal{F}) \simeq \lim_{\Delta} \Gamma(U_\bullet, \mathcal{F}(*))$.

We show that the functor $\Gamma_*(X, -)$ is fully faithful. Recall that $B = A[a_1^{-1}] \times \cdots \times A[a_r^{-1}]$, where $\{a_i\}_{i=1}^r \subset \pi_0(A_1)$ is the set of generators of $\pi_0(A)$ as an \mathbb{N} -graded commutative ring over $\pi_0(A_0)$. We have the faithfully flat affine morphism $f : U \rightarrow X$, where $U = \text{Spec}(B_0)$. Note that there is an equivalence $\Gamma(U, f^* \mathcal{O}_X(*)) \simeq B$ of \mathbb{Z} -graded \mathbb{E}_∞ -rings and hence that $\Gamma(U, f^* \mathcal{F}(*))$ is a \mathbb{Z} -graded B -module for a quasi-coherent sheaf \mathcal{F} of \mathcal{O}_X -modules on X . We have the restriction map $\Gamma_*(X, \mathcal{F}) \rightarrow \Gamma(U, f^* \mathcal{F}(*))$, which induces a map

$$B \otimes_A \Gamma_*(X, \mathcal{F}) \rightarrow \Gamma(U, f^* \mathcal{F}(*))$$

of \mathbb{Z} -graded B -modules. We shall compare $\Gamma(U, f^* \mathcal{F}(*))$ with the scalar extension $B \otimes_A \Gamma_*(X, \mathcal{F})$.

Lemma 2.4 (cf. [8]). *Let \mathcal{F} be a quasi-coherent sheaf of \mathcal{O}_X -module on X . There is a natural equivalence*

$$B \otimes_A \Gamma_*(X, \mathcal{F}) \xrightarrow{\simeq} \Gamma(U, f^* \mathcal{F}(*))$$

of \mathbb{Z} -graded B -modules.

Proof. We have $U = V_1 \times \cdots \times V_r$, where $V_i = \text{Spec}(A[a_i^{-1}]_0)$ for $1 \leq i \leq r$. This implies a decomposition

$$\Gamma(U, f^* \mathcal{F}(*)) \simeq \Gamma(V_1, \mathcal{F}(*)) \times \cdots \times \Gamma(V_r, \mathcal{F}(*)),$$

where $\Gamma(V_i, \mathcal{F}(*))$ is a \mathbb{Z} -graded $A[a_i^{-1}]$ -module for $1 \leq i \leq r$. Since $\Gamma(V_i, \mathcal{F}(*))$ is a \mathbb{Z} -graded $A[a_i^{-1}]$ -module, the restriction map $\Gamma_*(X, \mathcal{F}) \rightarrow \Gamma(U, f^* \mathcal{F}(*))$ induces a map $\Gamma_*(X, \mathcal{F})[a_i^{-1}] \rightarrow \Gamma_*(V_i, \mathcal{F})$ of \mathbb{Z} -graded $A[a_i^{-1}]$ -modules. It suffices to show that this map is an equivalence for any i with $1 \leq i \leq r$.

Let P be the partially ordered set of all nonempty finite subsets of $\{1, \dots, r\}$. We set $V_I = \bigcap_{i \in I} V_i$ for $I \in P$. By [5, Proposition 1.1.4.4], we have an equivalence

$$\Gamma_*(X, \mathcal{F}) \simeq \lim_{I \in P} \Gamma(V_I, \mathcal{F}(*))$$

of \mathbb{Z} -graded A -modules. Note that the right hand side is a finite limit indexed by P . Since filtered colimits commute with finite limits, we obtain an equivalence

$$\Gamma_*(X, \mathcal{F})[a_i^{-1}] \simeq \lim_{I \in P} (\Gamma(V_I, \mathcal{F}(*))[a_i^{-1}])$$

of \mathbb{Z} -graded $A[a_i^{-1}]$ -modules. By definition, we have an equivalence

$$\Gamma(V_I, \mathcal{F}(*))[a_i^{-1}] \simeq \Gamma(V_{I \cup \{i\}}, \mathcal{F}(*))$$

for any $I \in P$. We consider a functor $\theta : P \rightarrow \text{Mod}_{A[a_i^{-1}]}(\mathbb{Z})$ which assigns to $I \in P$ the \mathbb{Z} -graded $A[a_i^{-1}]$ -module $\Gamma(V_{I \cup \{i\}}, \mathcal{F}(*))$. Let P_i be the subset of P consisting of finite subsets of $\{1, \dots, r\}$ which contain i . Since the functor θ is a right Kan extension of the restriction to P_i , we have an equivalence

$$\lim_{I \in P} \Gamma(V_{I \cup \{i\}}, \mathcal{F}(*)) \simeq \lim_{J \in P_i} \Gamma(V_J, \mathcal{F}(*)).$$

By [5, Proposition 1.1.4.4], we have an equivalence

$$\Gamma(V_i, \mathcal{F}(*)) \simeq \lim_{J \in P_i} \Gamma(V_J, \mathcal{F}(*)).$$

Thus, we have an equivalence $\Gamma(V_i, \mathcal{F}(*)) \simeq \Gamma_*(X, \mathcal{F})[a_i^{-1}]$ of \mathbb{Z} -graded $A[a_i^{-1}]$ -modules. \square

Especially, we can see that the functor $\Gamma_*(X, -) : \text{QCoh}(X) \rightarrow \text{Mod}_A(\mathbb{Z})$ is fully faithful.

We have the adjunction $\widetilde{(-)} : \text{Mod}_A(\mathbb{Z}) \rightleftarrows \text{QCoh}(X) : \Gamma_*(X, -)$ of ∞ -categories. Since the left adjoint $\widetilde{(-)}$ is a symmetric monoidal functor, the right adjoint $\Gamma_*(X, -)$ is a lax symmetric monoidal functor. In particular, $\Gamma_*(X, \mathcal{O}_X)$ is a \mathbb{Z} -graded \mathbb{E}_∞ -ring and there is a map

$$A \longrightarrow \Gamma_*(X, \mathcal{O}_X)$$

of \mathbb{Z} -graded \mathbb{E}_∞ -rings. For a quasi-coherent sheaf \mathcal{F} of \mathcal{O}_X -modules on X , $\Gamma_*(X, \mathcal{F})$ is a \mathbb{Z} -graded $\Gamma_*(X, \mathcal{O}_X)$ -module. We note that the \mathbb{Z} -graded A -module structure on $\Gamma_*(X, \mathcal{F})$ is obtained from the \mathbb{Z} -graded $\Gamma_*(X, \mathcal{O}_X)$ -module structure through the map $A \rightarrow \Gamma_*(X, \mathcal{O}_X)$ of \mathbb{Z} -graded \mathbb{E}_∞ -rings.

3. THE PROPERTIES OF QUASI-COHERENT SHEAVES ON SPECTRAL PROJECTIVE SCHEMES

The definition of projective schemes by using quasi-coherent sheaves may be valuable in the non-commutative geometry. For example, for a field k and finitely generated commutative graded k -algebra which is generated by degree 1 elements, Artin and Zhan shows that there is an categorical equivalence between the category of certain coherent sheaves on projective scheme of A and the category of finite graded right A -modules. Vervekin also studied injective objects and Ext-groups in the category of finite graded A -modules.

Lemma 3.1. *For $M \in \text{Mod}_A(\mathbb{Z})$, we have $\widetilde{M} \simeq 0$ if and only if $\pi_n(M)$ is Zariski locally bounded above for any $n \in \mathbb{Z}$.*

Proof. We take a set $\{a_i\}_{i=1}^r \subset \pi_0(A_1)$ of generators of $\pi_0(A)$ as an \mathbb{N} -graded commutative ring over $\pi_0(A_0)$. Then there is an affine open covering $\{V_i\}_{i=1}^r$ of X , where $V_i = \text{Spec}(A[a_i^{-1}]_0)$. We have $\widetilde{M} \simeq 0$ if and only if $\widetilde{M}|_{V_i} \simeq 0$ for $i = 1, \dots, r$. Under the equivalence $\text{QCoh}(V_i) \simeq \text{Mod}_{A[a_i^{-1}]_0}$, the restriction $\widetilde{M}|_{V_i}$ corresponds to an $A[a_i^{-1}]_0$ -module $M[a_i^{-1}]_0$. Hence $\widetilde{M} \simeq 0$ if and only if $M[a_i^{-1}]_0 \simeq 0$ for $i = 1, \dots, r$. This is equivalent to the condition that $\pi_n(M)[a_i^{-1}]_0 = 0$ for any $n \in \mathbb{Z}$ and $i = 1, \dots, r$. \square

Definition 3.2. We say that a \mathbb{Z} -graded A -module M is locally bounded above in homotopy groups if the \mathbb{Z} -graded $\pi_0(A)$ -module $\pi_n(M)$ is locally bounded above for each $n \in \mathbb{Z}$. We define $\text{Mod}_A^{\text{lbah}}(\mathbb{Z})$ to be the full subcategory of $\text{Mod}_A(\mathbb{Z})$ spanned by those objects that are locally bounded above in homotopy groups.

We have the adjunction $\widetilde{(-)} : \text{Mod}_A(\mathbb{Z}) \rightleftarrows \text{QCoh}(X) : \Gamma_*(X, -)$ of stable presentable ∞ -categories, where the left adjoint $\widetilde{(-)}$ is symmetric monoidal and the right adjoint $\Gamma_*(-)$ is lax symmetric monoidal and fully faithful. By Lemma 3.1, we have $\widetilde{M} \simeq 0$ if and only if $\pi_n(M)$ is locally bounded above for any $n \in \mathbb{Z}$.

Hence we obtain the following proposition.

Proposition 3.3. *The functor $\widetilde{(-)} : \text{Mod}_A(\mathbb{Z}) \rightarrow \text{QCoh}(X)$ induces an equivalence*

$$\text{Mod}_A(\mathbb{Z}) / \text{Mod}_A^{\text{lbah}}(\mathbb{Z}) \xrightarrow{\simeq} \text{QCoh}(X)$$

of stable symmetric monoidal presentable ∞ -categories. Here, W be the class of all morphisms in $\text{Mod}_A(\mathbb{Z})$ whose cofiber lies in $\text{Mod}_A^{\text{lbah}}(\mathbb{Z})$ and the left hand side is the localization with respect to the class W .

\square

3.1. The equivalence of Serre theorem. In this subsection, we give a short survey of a generalization of Serre theorem as in [8]. It is given by restricting the equivalence in Proposition 3.3 to "finitely generated part". We prepare some notation for "finiteness".

Definition 3.4. Let A be a connective \mathbb{N} -graded \mathbb{E}_∞ -ring.

- (i) Let R be a connective Noetherian \mathbb{E}_∞ -ring. Recall that an R -module M is almost perfect if $\pi_n(M)$ is a finitely generated $\pi_0(R)$ -module for any $n \in \mathbb{Z}$ and if $\pi_n(M) = 0$ for $n \ll 0$ [3, Proposition 7.2.4.17].
- (ii) We let $\text{Mod}_A^{\text{perf}}(\mathbb{Z})$ be the smallest stable subcategory of $\text{Mod}_A(\mathbb{Z})$ which contains $A(q)$ for all $q \in \mathbb{Z}$ and is closed under retracts. We say that a \mathbb{Z} -graded A -module M is perfect if it belongs to the full subcategory $\text{Mod}_A^{\text{perf}}(\mathbb{Z})$.
- (iii) The quasi-coherent sheaf \widetilde{M} is almost perfect if a \mathbb{Z} -graded A -module M is almost finitely generated.
- (iv) We say that a \mathbb{Z} -graded A -module M is almost finitely generated if the following conditions are satisfied: for each $n \in \mathbb{Z}$, the \mathbb{Z} -graded $\pi_0(A)$ -module $\pi_n(M)$ is finitely generated, and, for $n \ll 0$, $\pi_n(M) = 0$. We define an ∞ -category $\text{Mod}_A^{\text{afg}}(\mathbb{Z})$ to be the full subcategory of $\text{Mod}_A(\mathbb{Z})$ spanned by almost finitely generated \mathbb{Z} -graded A -modules.
- (v) Let M be an almost finitely generated \mathbb{Z} -graded A -module. We say that M is almost torsion if the \mathbb{Z} -graded $\pi_0(A)$ -module $\pi_n(M)$ is bounded above for each $n \in \mathbb{Z}$. We define an ∞ -category $\text{Mod}_A^{\text{ator}}(\mathbb{Z})$ to be the full subcategory of $\text{Mod}_A^{\text{afg}}(\mathbb{Z})$ spanned by almost torsion \mathbb{Z} -graded A -modules.

We give a characterization of M in $\text{Mod}_A^{\text{afg}}(\mathbb{Z})$ satisfying $\widetilde{M} \simeq 0$ in terms of the \mathbb{Z} -graded $\pi_0(A)$ -modules $\pi_n(M)$ for $n \in \mathbb{Z}$. By Lemma 3.1, we have $\widetilde{M} \simeq 0$ if and only if M is almost torsion.

If we restrict the symmetric monoidal functor $\widetilde{(-)} : \text{Mod}_A(\mathbb{Z}) \rightarrow \text{QCoh}(X)$ to the full ∞ -subcategory $\text{Mod}_A^{\text{afg}}(\mathbb{Z})$, it factors through $\text{QCoh}(\text{Proj}(A))^{\text{aperf}}$. By the above argument, it also induces a symmetric monoidal exact functor

$$\widetilde{(-)} : \text{Mod}_A^{\text{afg}}(\mathbb{Z}) / \text{Mod}_A^{\text{ator}}(\mathbb{Z}) \longrightarrow \text{QCoh}(X)^{\text{aperf}}.$$

The Serre theorem describes the ∞ -category $\text{QCoh}(X)^{\text{aperf}}$ of almost perfect quasi-coherent sheaves on X in terms of \mathbb{Z} -graded A -modules. It requires that the above symmetric monoidal exact functor gives an equivalence of symmetric monoidal ∞ -categories. To see this, especially, we need the essentially surjectivity of this functor.

The key proposition is the following, which is proved by calculating the spectral sequence.

Proposition 3.5 ([8]). *If $\mathcal{F} \in \text{QCoh}(X)$ is almost perfect, then we have*

- the \mathbb{Z} -graded $\pi_0(A)$ -module $\pi_n(\Gamma_*(X, \mathcal{F}))$ is strongly quasi-finitely generated for each $n \in \mathbb{Z}$, and
- $\pi_n(\Gamma_*(X, \mathcal{F})) = 0$ for $n \ll 0$.

□

The ∞ -category $\mathrm{QCoh}(X)^{\mathrm{aperf}}$ of almost perfect quasi-coherent sheaves on X can be related with the ∞ -category $\mathrm{Mod}_A^{\mathrm{afg}}(\mathbb{Z})$ of almost finitely generated \mathbb{Z} -graded A -modules.

Let $f \in \pi_0(A)$ be homogeneous of positive degree and let $U = \mathrm{Spec}(A[f^{-1}]_0)$ be an affine open subscheme of X . The restriction $\widetilde{M}|_U$ corresponds to the $A[f^{-1}]_0$ -module $M[f^{-1}]_0$. We see that $M[f^{-1}]_0$ is almost perfect since M is almost finitely generated.

Conversely, by Proposition 3.5, we obtain

- If $\mathcal{F} \in \mathrm{QCoh}(X)$ is almost perfect, then $\pi_n \mathcal{F}$ is a coherent sheaf of \mathcal{O}_X -modules on X for any $n \in \mathbb{Z}$.
- $\mathcal{F} \in \mathrm{QCoh}(X)$ is almost perfect, then $\pi_n \mathcal{F} = 0$ for $n \ll 0$.

By proceeding local argument, we have the following.

Theorem 3.6 ([8]). *The functor $\widetilde{(-)} : \mathrm{Mod}_A(\mathbb{Z}) \rightarrow \mathrm{QCoh}(X)$ induces an equivalence*

$$\mathrm{Mod}_A^{\mathrm{afg}}(\mathbb{Z}) / \mathrm{Mod}_A^{\mathrm{ator}}(\mathbb{Z}) \xrightarrow{\simeq} \mathrm{QCoh}(X)^{\mathrm{aperf}}$$

of small stable symmetric monoidal ∞ -categories.

REFERENCES

- [1] S. Glasman, Day convolution for ∞ -categories, *Math. Res. Lett.* 23 (2016), no. 5, 1369–1385.
- [2] J. Lurie, *Higher topos theory*, *Annals of Mathematics Studies*, 170, Princeton University Press, Princeton, NJ, 2009.
- [3] J. Lurie, *Higher Algebra* (version September 18, 2017), preprint, available at <https://www.math.ias.edu/~lurie/>.
- [4] J. Lurie, *Structured Spaces*, preprint, 2011, available at <https://www.math.ias.edu/~lurie/>.
- [5] J. Lurie, *Spectral Algebraic Geometry* (version February 3, 2018), preprint, available at <https://www.math.ias.edu/~lurie/>.
- [6] J.-P. Serre, *Faisceaux algébriques cohérents*, *Ann. of Math. (2)* 61 (1955), 197–278.
- [7] The Stacks project authors, *The Stacks project*, 2020, <https://stacks.math.columbia.edu/>.
- [8] M. Ohara, and T. Torii, On graded \mathbb{E}_∞ -rings and projective schemes in spectral algebraic geometry. *Journal of Homotopy and Related Structures*, Vol.17 (2022), 105–144.
- [9] B. Toën, and G. Vezzosi, *Homotopical algebraic geometry. I. Topos theory*, *Adv. Math.* 193 (2005), no. 2, 257–372.
- [10] B. Toën, and G. Vezzosi, *Homotopical algebraic geometry. II. Geometric stacks and applications*, *Mem. Amer. Math. Soc.* 193 (2008).

OSHIMA COLLEGE ASSISTANT PROFESSOR

Email address: `primarydecomposition@gmail.com`