

# Segre cubic 4-fold

Shigeru MUKAI

6/17/22(F)

@Roma Univ. Tre

Abstract: The cyclic triple covering of the projective 4-space with branch the Segre cubic is characterized by 10 cusps as the Segre 3-fold is so by 10 nodes. The Fano variety of lines is birationally equivalent to the Hilbert square of a K3 surface studied by Vinberg(1983). I will discuss the binational automorphism group of this holomorphic symplectic 4-fold.

§1 Main motivation and side job

§2  $K3^{[2]}$  4-folds

§3 Vinberg's theorem and generalization to  $K3^{[2]}$  4-folds

§4 Review of Vinberg's proof

§5 Proof of main theorem

References (8 items)

## §1 Main motivation and side job

Main motivation: generalize  $\text{Aut}(K3)$  to  $\text{Bir-Aut}(HK)$  (replacing nef cone with movable cone).

Side job: Kummer theory


(I start with) Kummer quartic surface,  $\Phi: A/\pm 1 \xrightarrow{|\mathbb{Z}/2|} P^3$

(1)  $A = J(C)$ ,  $C$ : curve of genus 2,  $\Phi$ : embedding

( $16_6 - 16_6$ ) configuration of nodes and tropes

(2)  $A = E_1 \times E_2$ :  $\Phi$  is of degree 2 onto a smooth quadric  $P^1 \times P^1$

$$\tau^2 = f_4(x)g_4(y)$$



branch locus

## Order 3 analogue

$(A, \sigma)$  : Abelian surface & symplectic automorphism of order 3


Quotient  $A/\sigma$  is a K3 surface with 9 cusps,  $9A_2$

(1)  $A=J(C)$ ,  $C: \tau^2 = q(x^3)$ ,  $\sigma: x \mapsto \omega x$ ,  $\omega^3 = 1$

(Barth-)Bertin-Vanhaecke sextic.  $A/\sigma \subset \mathbb{P}^4$ , Image = (2, 3) c.i.  
 $(9_4 - 9_4)$  configuration of 9 cusps and 9 conics

(2)  $A = E \times E$ ,  $E$  with CM of order 3

$A/\sigma \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ ,  $\tau^3 = (x^2 - x)(y^2 - y)$



branch

$[A/\sigma \hookrightarrow \mathbb{P}^4] = (\text{quadric cone}) \cap (3)$  c.i.

## Mini history of Aut S and Picard lattice in $U +$ (Leech)

	(1)	(2)
Kummer	Kondo(1998), $(A_3 + 6A_1)^\perp$	Keum-Kondo(2000), $(2D_4)^\perp$
ord 3 ver.	Not yet, $(A_5 + A_1)^\perp$	Vinberg(1983), $E_6^\perp$

## §2. K3<sup>[2]</sup> 4-folds

Simplest higher dimensional analogue of a K3 surface is the Hilbert square of a K3 (and its deformations)

Let

$S^{[2]} \rightarrow S^{(2)}$  be the minimal resolution of symmetric product.

$S^{[2]}$  = moduli of ideal sheaves of 2 points (allowing inf. nears)  
 $= M_S(1, 0, -1)$

$\text{Pic } S^{[2]} = \text{Pic } S \oplus \mathbb{Z}\delta, \quad \delta = (1, 0, 1)$   
 $= (1, 0, -1) \text{ in } \mathbb{Z} \oplus \text{Pic } S \oplus \mathbb{Z}$

Notation.  $\alpha$  on  $S$  is identified with  $(0, \alpha, 0)$  on  $S^{[2]}$ .

Relation with cubic 4-folds and K3 sextic  $S = (2, 3)$ , c.i. in  $P^4$ .

$S : q(x, y, z, u, v) = d(x, y, z, u, v) = 0$  in  $P^4$

$X = X_S : q(x, y, z, u, v)w + d(x, y, z, u, v) = 0$  in  $P^5$

This cubic 4-fold is singular at (100000).

Fact: Fano variety of lines  $F(X_S)$  is birational with  $S^{[2]}$

(1)  $S$  : Bertin-Vanhaecke  $\Rightarrow X_S$  has  $A_1 + 9A_2$

(2)  $S$  : Vinberg  $\Rightarrow X_S$  has  $10A_2$  since the quadric hull of  $S$  is a cone

Observation:  $X_S$  is a 4-dimensional analogue of Segre cubic 3-fold when  $S$  : Vinberg.

$$Y : \sum_1^6 y_i = \sum_1^6 y_i^3 = 0 \subset \mathbb{P}^5$$

$$X_S : \left( \sum_1^6 x_i \right) \left( \sum_1^6 x_i^2 \right) = 2 \sum_1^6 x_i^3 \subset \mathbb{P}^5$$

Segre cubic 3-fold  $Y$  is characterized by  $10A_1$   
 \_\_\_\_\_ 4-fold  $X$  \_\_\_\_\_  $10A_2$

$X_S$  is a triple cyclic covering of  $P^4$  with branch Segre cubic.

Both have an action of symmetric group  $G_6$  of degree 6

$$\begin{aligned}
 \text{Aut}(S \subset \mathbb{P}^4) &\cong C_3 \cdot [ (G_3 \times G_3) \cdot C_2 ] \\
 &\quad \cap \quad \quad \quad \cap \quad \quad \quad \text{Normalizer of } \sigma \\
 \text{Aut}(X_S \subset \mathbb{P}^5) &\cong G_6 \ni (123)(456) =: \sigma
 \end{aligned}$$

§3 Vinberg's theorem and generalization to  $K3^{[2]}$  4-folds

Theorem (Vinberg)

$$\text{Aut } S = (\text{Free product of 12 involutions}) \rtimes \text{Aut}(S \subset \mathbb{P}^4)$$

Main Theorem  $\text{Bir-Aut}(S^{[2]})$  is semi-direct product of  $\langle 84+120 \text{ involutions} \rangle$  by symmetric group  $G_7$  of degree 7.

Surprisingly,

$G_6$ -action on  $S$  extends to a birational  $G_7$ -action on  $S^{[2]}$ .

Reason: a symmetry of degree 7 induced by Lagrangian fibration

$$S^{[2]} \dashrightarrow \mathbb{P}^2 \quad (\text{of } 3\tilde{A}_6\text{-type}).$$

Mordell-Weil group has a 7-torsion (birational) section.

§4 Review of Vinberg's proof

Quick review on  $\text{Aut } S$ , Picard number 20,

$\text{Pic } S = U + 2 E_8 + A_2$


Orthogonal group  $O^+(U+2E_8+A_2)$

$= \langle 24 (-2)\text{-reflections, } 12 (-6)\text{-reflections} \rangle \rtimes \text{Aut}(S \subset P^4)$


$24 = 6 + 9 \cdot 2$

$$T_S = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$$

$$U = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, A_2 = \begin{pmatrix} -2 & 1 \\ 1 & -2 \end{pmatrix}$$



Dual graph of 24  $P^1$ 's



6 circuits of length 18

$S \rightarrow P^1$  elliptic fibration

with  $I_{18}$  fiber, MW group  $Z \oplus Z/6$


$\Rightarrow$  2 anti-symplectic involutions

$(-1)$ -mult. &

its composite with 2-torsion translation

Modulo  $\text{Aut}(S \subset P^4)$ , these act as

$(-6)$ -reflection.



## §5 Proof of main theorem

Look at the action of  $\text{Bir-Aut } S^{[2]}$  on  $\text{Pic } S \oplus \mathbb{Z}\delta$ , and on the movable cone in it. The rest is basically the same as (Vinberg's) K3, but get complicated in two points:

- (1) the orthogonal group is no more more reflective
- (2) Divisibility should be taken into account. There are two types of  $(-2)$ -divisors:

- a)  $(-2)$  effective divisor  $\text{Im}[E \times S \rightarrow S^{[2]}]$  for  $(-2)$ -curve  $E$  on  $S$  (divisibility 1)
- b) half  $\delta$  of exceptional divisor class (divisibility 2)


(1) is overcome by Conway-Borcherds domain CB in the positive cone.

CB domain is surrounded by 309 walls:

$$309 = \underbrace{35 + 70}_{\substack{\text{(-2)-walls} \\ \text{(divisibility 1 \& 2,} \\ \text{respectively)}}} + \underset{\uparrow}{84} + \underset{\uparrow}{120}$$

① ②
③
④

$$\star O^+(U+2E_8+A_2+A_1) = \left\langle \begin{array}{l} 105 \text{ (-2)-reflections} \\ 84 \text{ (-6)-reflections} \\ 120 \text{ quasi-reflections} \end{array} \right\rangle \times \mathcal{G}_7$$


  
 symmetry of  
 CB domain


Geometrization of ★

① Basic (-2)-divisors (of divisibility 1) # = 35 = 24 + 2 + 9

24 are "pull-back" of (-2)'s from S.

Extra 11(=2+9) are (1, f, 1) for elliptic pencil |f| on S of minimal Coxeter number (= 12). Since f is isotopic, this divisor has Beauville-square (-2). Geometrically, (1, f, 1) is the Zariski closure of locus of {a, b} with a ≠ b and Φ(a) = Φ(b).

Two f's are of type 3E\_6 and 9 of type D\_7+A\_11.



The dual graph of these 35 (-2) divisors is the 4-valent odd graph O\_4.

② (-2) divisors of divisibility 2  $\Leftrightarrow$  70 edges of  $O_4$

③ (-6)-walls  $\Leftrightarrow$  Induced automorphism from  $S$

④ (-42)-walls: Non-induced automorphism  $\Leftrightarrow$  (-1) multiplication of Lagrangian fibration of type  $3\tilde{A}_6 \pmod{\mathfrak{S}_7}$

Final answer:  $\text{Aut } S^{[2]} = \langle 84+120 \text{ involutions} \rangle \rtimes \mathfrak{S}_7$ .



## §1 Main motivation and side job

Bertin, J. and Vanhaecke, P.: The even system and generalized Kummer surfaces, *Math. Proc. Camb. Phil. Soc.*, **116**(1994), 131-142.

Barth, W.: K3 surfaces with nine cusps, *Geom. Dedicata*, **72**(1998), 171-178.

Kondo, S.: The automorphism group of a generic Jacobian Kummer surface, *J. Alg. Geom.* **7**(1998), 589-609.

Keum, J-H. and Kondo, S.: The automorphism groups of Kummer surfaces associated with the product of two elliptic curves, *Trans. Amer. Math. Soc.*, **353**(2000), 1469-1487.

Vinberg, E.B.: The two most algebraic K3 surfaces, *Math. Ann.* **265**(1983), 1-27.

## §2 $K3^{[2]}$ 4-folds

Dolgachev, I.: Corrado Segre and nodal cubic threefolds, in *“From classical to modern algebraic geometry; Corrado Segre’s mastership and legacy”*, Springer, 2016, pp. 429-450.

## §3 Vinberg's theorem and generalization to $K3^{[2]}$ 4-folds

Oguiso, K.: Picard number of the generic fiber of an abelian variety fingered hyperkahler manifold, *Math. Ann.*, **344**(2009), 929-937.

## §5 Proof of main theorem

Borcherds, R.: Automorphism groups of Lorentzian lattices, *J. Algebra*, **111**(1987), 133-153.