

## SOME RESULTS OF $F$ -BIHARMONIC MAPS

YINGBO HAN AND SHUXIANG FENG

ABSTRACT. In this paper, we give the notion of  $F$ -biharmonic maps, which is a generalization of biharmonic maps. We derive the first variation formula which yields  $F$ -biharmonic maps. Then we investigate the harmonicity of  $F$ -biharmonic maps under the curvature conditions on the target manifold  $(N, h)$ . We also introduce the stress  $F$ -bienergy tensor  $S_{F,2}$ . Then, by using the stress  $F$ -bienergy tensor  $S_{F,2}$ , we obtain some nonexistence results of proper  $F$ -biharmonic maps under the assumption that  $S_{F,2} = 0$ . Moreover, we derive some monotonicity formulas for the special case of the biharmonic map, i.e., where  $F$ -biharmonic map with  $F(t) = t$ . Then, by using these monotonicity formulas, we obtain new results on the non existence of proper biharmonic isometric immersions from complete manifolds.

### 1. INTRODUCTION

Harmonic maps play a central roll in variational problems for smooth maps between manifolds  $u: (M, g) \rightarrow (N, h)$  as the critical points of the energy functional  $E(u) = \frac{1}{2} \int_M \|du\|^2 dv_g$ . On the other hand, in 1981, J. Eells and L. Lemaire [7] proposed the problem to consider the  $k$ -harmonic maps which are critical maps of the functional

$$E_k(u) = \int_M \frac{\|(d + \delta)^k u\|^2}{2} dv_g$$

---

Received January 10, 2013; revised July 15, 2013.

2010 *Mathematics Subject Classification*. Primary 58E20, 53C21.

*Key words and phrases*.  $F$ -biharmonic map; stress  $F$ -bienergy tensor; harmonic map.




Go back

Full Screen

Close

Quit



for smooth maps  $u : M \rightarrow N$ . G. Y. Jiang [9] studied the first and second variation formulas of the bienergy  $E_2$  where critical maps of  $E_2$  are called biharmonic maps. There have been extensive studies on biharmonic maps (for instance, see [9, 13, 14, 15, 16, 18, 19]).

Let  $F : [0, \infty) \rightarrow [0, \infty)$  be a  $C^3$  function such that  $F' > 0$  on  $(0, \infty)$ . For a smooth map  $u : (M, g) \rightarrow (N, h)$  between Riemannian manifolds  $(M, g)$  and  $(N, h)$ , we define the  $F$ - $k$ -energy  $E_{F,k}(u)$  of  $u$  by

$$E_{F,k}(u) = \int_M F\left(\frac{\|(d + \delta)^k u\|^2}{2}\right) dv_g,$$

which is  $E_k(u)$  if  $F(t) = t$ . When  $k = 1$ , we have

$$E_{F,1}(u) = \int_M F\left(\frac{\|du\|^2}{2}\right) dv_g = E_F(u),$$

which was introduced by M. Ara in [1]. The critical maps of  $E_F(u)$  are called  $F$ -harmonic maps which are the generalization of harmonic maps,  $p$ -harmonic maps or exponentially harmonic maps. There have been extensive studies in this area (for instance, [4, 5, 11, 12]). When  $k = 2$ , we have

$$E_{F,2}(u) = \int_M F\left(\frac{\|\tau(u)\|^2}{2}\right) dv_g,$$

where  $\tau(u) = -\delta du = \text{trace } \tilde{\nabla}(du)$ . It is the bienergy of G. Y. Jiang [9], the  $p$ -bienergy of P. Hornung and R. Moser [6] or exponentially bienergy when  $F(t) = t$ ,  $F(t) = (2t)^{\frac{p}{2}}$  or  $F(t) = e^t$ . We say that  $u$  is an  $F$ -biharmonic map if

$$\frac{d}{dt} E_{F,2}(u_t)|_{t=0} = 0$$




Go back

Full Screen

Close

Quit



for any compactly supported variation  $u_t: M \rightarrow N$  with  $u_0 = u$ . In this note, we derive the first variation formula which yields  $F$ -biharmonic maps. Then we investigate the harmonicity of  $F$ -biharmonic maps under the curvature conditions on the target manifold  $(N, h)$ . We also introduce the stress  $F$ -bienergy tensor  $S_{F,2}$ . Then, by using the stress  $F$ -bienergy tensor  $S_{F,2}$ , we obtain some non existence results of proper  $F$ -biharmonic maps under the assumption  $S_{F,2} = 0$ . Also, we derive some monotonicity formulas for the special case of a biharmonic map, i.e., an  $F$ -biharmonic map with  $F(t) = t$ . Then, by using these monotonicity formulas, we investigate the harmonicity of biharmonic isometric maps from complete manifolds.

*Remark 1.1.* In [17], the authors introduced  $f$ -biharmonic maps which are critical points of the bi- $f$ -energy functional

$$E_f^2(u) = \frac{1}{2} \int_M \|\tau_f(u)\|^2 dv_g,$$

where  $\tau_f(u) = f\tau(u) + du(\text{grad } f)$  and  $f \in C^\infty(M)$ . We think that it is more reasonable to call them “bi- $f$ -harmonic maps” as parallel to “biharmonic maps”.



Go back


Full Screen

Close

Quit

## 2. THE FIRST VARIATION FORMULA

Let  $\nabla$  and  ${}^N\nabla$  always denote the Levi-Civita connections of  $M$  and  $N$ , respectively. Let  $\tilde{\nabla}$  be the induced connection on  $u^{-1}TN$  defined by  $\tilde{\nabla}_X W = {}^N\nabla_{du(X)} W$ , where  $X$  is a tangent vector of  $M$  and  $W$  is a section of  $u^{-1}TN$ . We choose a local orthonormal frame field  $\{e_i\}$  on  $M$ . We define



the  $F$ -bitension field  $\tau_{F,2}(u)$  of  $u$  by

$$\begin{aligned} \tau_{F,2}(u) &= -J(F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u)) \\ &= -\tilde{\Delta}(F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u)) - \sum_i R^N(du(e_i), F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u)) du(e_i), \end{aligned}$$

where  $J$  is the Jacobi operator of the second variation for the energy  $E(u) = \frac{1}{2} \int_M \|du\|^2 dv_g$ ,  $\tilde{\Delta} = -\sum_i (\tilde{\nabla}_{e_i} \tilde{\nabla}_{e_i} - \tilde{\nabla}_{\nabla_{e_i} e_i})$  is the rough Laplacian on the section of  $u^{-1}TN$  and  $R^N(X, Y) = [{}^N\nabla_X, {}^N\nabla_Y] - {}^N\nabla_{[X, Y]}$  is the curvature operator on  $N$ .

Under the notation above we have the following theorem

**Theorem 2.1** (The first variation formula). *Let  $u: M \rightarrow N$  be a smooth map. Then*

$$(1) \quad \frac{d}{dt} E_{F,2}(u_t) |_{t=0} = \int_M h(\tau_{F,2}(u), V) dv_g,$$

where  $V = \frac{d}{dt} u_t |_{t=0}$ .

*Proof.* Let  $\Psi: (-\varepsilon, \varepsilon) \times M \rightarrow N$  be defined by  $\Psi(t, x) = u_t(x)$ , where  $(-\varepsilon, \varepsilon) \times M$  is equipped with the product metric. We extend the vector fields  $\frac{\partial}{\partial t}$  on  $(-\varepsilon, \varepsilon)$ ,  $X$  on  $M$  naturally on  $(-\varepsilon, \varepsilon) \times M$ , and denote those also by  $\frac{\partial}{\partial t}$ ,  $X$ . Then

$$d\Psi \left( \frac{\partial}{\partial t} \right) = \frac{d}{dt} u_t |_{t=0} = V.$$

We shall use the same notations  $\nabla$  and  $\tilde{\nabla}$  for the Levi-Civita connection on  $(-\varepsilon, \varepsilon) \times M$  and the induced connection on  $\Psi^{-1}TN$ , respectively.



Go back

Full Screen

Close

Quit

We compute

$$\begin{aligned}
 & \frac{\partial}{\partial t} F \left( \frac{\|\tau(u_t)\|^2}{2} \right) \\
 &= F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \frac{1}{2} \frac{\partial}{\partial t} \|\tau(u_t)\|^2 \\
 &= F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) h \left( \tilde{\nabla}_{\frac{\partial}{\partial t}} \tau(u_t), \tau(u_t) \right) \\
 (2) \quad &= \sum_i F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) h \left( \tilde{\nabla}_{\frac{\partial}{\partial t}} [(\tilde{\nabla}_{e_i} d\Psi)(e_i)], \tau(u_t) \right) \\
 &= \sum_i h \left( \tilde{\nabla}_{\frac{\partial}{\partial t}} \tilde{\nabla}_{e_i} d\Psi(e_i) - \tilde{\nabla}_{\frac{\partial}{\partial t}} d\Psi(\nabla_{e_i} e_i), F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right) \\
 &= \sum_i h \left( R^N \left( d\Psi \left( \frac{\partial}{\partial t} \right), d\Psi(e_i) \right) d\Psi(e_i), F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right) \\
 &\quad + \sum_i h \left( \tilde{\nabla}_{e_i} \tilde{\nabla}_{e_i} d\Psi \left( \frac{\partial}{\partial t} \right) - \tilde{\nabla}_{\nabla_{e_i} e_i} d\Psi \left( \frac{\partial}{\partial t} \right), F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right),
 \end{aligned}$$

where we use

$$\tilde{\nabla}_{\frac{\partial}{\partial t}} d\Psi(e_i) - \tilde{\nabla}_{e_i} d\Psi \left( \frac{\partial}{\partial t} \right) = d\Psi \left[ \frac{\partial}{\partial t}, e_i \right] = 0$$

and

$$\tilde{\nabla}_{\frac{\partial}{\partial t}} d\Psi(\nabla_{e_i} e_i) - \tilde{\nabla}_{\nabla_{e_i} e_i} d\Psi \left( \frac{\partial}{\partial t} \right) = d\Psi \left[ \frac{\partial}{\partial t}, \nabla_{e_i} e_i \right] = 0$$

for the fifth equality.



Go back

Full Screen

Close

Quit



Let  $X_t$  and  $Y_t$  be two compactly supported vector fields on  $M$  such that  $g(X_t, Z) = h(\tilde{\nabla}_Z d\Psi \left( \frac{\partial}{\partial t} \right), F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t))$  and  $g(Y_t, Z) = h(d\Psi \left( \frac{\partial}{\partial t} \right), \tilde{\nabla}_Z \left( F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right))$  for any vector field  $Z$  on  $M$ . Then the divergence of  $X_t$  and  $Y_t$  are given by the following:

$$\begin{aligned}
 \text{div}(X_t) &= \sum_k g(\nabla_{e_k} X_t, e_k) = \sum_k e_k g(X_t, e_k) - \sum_k g(X_t, \nabla_{e_k} e_k) \\
 &= \sum_k e_k h \left( \tilde{\nabla}_{e_k} d\Psi \left( \frac{\partial}{\partial t} \right), F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right) \\
 &\quad - \sum_k h \left( \tilde{\nabla}_{\nabla_{e_k} e_k} d\Psi \left( \frac{\partial}{\partial t} \right), F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right) \\
 (3) \quad &= \sum_k h \left( \tilde{\nabla}_{e_k} \tilde{\nabla}_{e_k} d\Psi \left( \frac{\partial}{\partial t} \right) \right. \\
 &\quad \left. - \tilde{\nabla}_{\nabla_{e_k} e_k} d\Psi \left( \frac{\partial}{\partial t} \right), F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right) \\
 &\quad + \sum_k h \left( \tilde{\nabla}_{e_k} d\Psi \left( \frac{\partial}{\partial t} \right), \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right] \right)
 \end{aligned}$$



Go back

Full Screen

Close

Quit

and

$$\begin{aligned}
 \operatorname{div}(Y_t) &= \sum_k g(\nabla_{e_k} Y_t, e_k) = \sum_k e_k g(Y_t, e_k) - \sum_k g(Y_t, \nabla_{e_k} e_k) \\
 &= \sum_k e_k h \left( d\Psi \left( \frac{\partial}{\partial t} \right), \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right] \right) \\
 &\quad - \sum_k h \left( d\Psi \left( \frac{\partial}{\partial t} \right), \tilde{\nabla}_{\nabla_{e_k} e_k} \left( F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right) \right) \\
 (4) \quad &= \sum_k h \left( d\Psi \left( \frac{\partial}{\partial t} \right), \tilde{\nabla}_{e_k} \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right] \right) \\
 &\quad - \tilde{\nabla}_{\nabla_{e_k} e_k} \left[ F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right] \\
 &\quad + \sum_k h \left( \tilde{\nabla}_{e_k} d\Psi \left( \frac{\partial}{\partial t} \right), \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right] \right).
 \end{aligned}$$

From (2), (3) and (4), we have

$$\begin{aligned}
 \frac{\partial}{\partial t} F \left( \frac{\|\tau(u_t)\|^2}{2} \right) &= \sum_i h \left( R^N \left( d\Psi \left( \frac{\partial}{\partial t} \right), d\Psi(e_i) \right) d\Psi(e_i), F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right) \\
 (5) \quad &+ \sum_i h \left( d\Psi \left( \frac{\partial}{\partial t} \right), \tilde{\nabla}_{e_i} \tilde{\nabla}_{e_i} \left[ F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right] \right) \\
 &\quad - \tilde{\nabla}_{\nabla_{e_i} e_i} \left[ F' \left( \frac{\|\tau(u_t)\|^2}{2} \right) \tau(u_t) \right] + \operatorname{div}(X_t) - \operatorname{div}(Y_t).
 \end{aligned}$$




Go back

Full Screen

Close

Quit



By (5) and Green's theorem, we have

$$\begin{aligned}
 & \frac{d}{dt} E_{F,2}(u_t)|_{t=0} \\
 &= \int_M \frac{\partial}{\partial t} F\left(\frac{\|\tau(u_t)\|^2}{2}\right) \Big|_{t=0} dv_g \\
 &= \int_M h\left(-\tilde{\Delta}\left[F'\left(\frac{\|\tau(u)\|^2}{2}\right)\tau(u)\right] \right. \\
 &\quad \left. - \sum_i R^N\left(du(e_i), \left[F'\left(\frac{\|\tau(u)\|^2}{2}\right)\tau(u)\right]\right) du(e_i), V\right) dv_g \\
 &= \int_M h(\tau_{F,2}(u), V) dv_g.
 \end{aligned}$$

This proves Theorem 2.1. □

The first variation formula allows us to define the notion of an  $F$ -biharmonic map for the functional  $E_{F,2}(u)$ .

**Definition 2.2.** A smooth map  $u$  is called an  $F$ -biharmonic map for the functional  $E_{F,2}(u)$  if it is a solution of the Euler-Lagrange equation  $\tau_{F,2}(u) = 0$ .

*Remark 2.3.* By Definition 2.2, we know that any harmonic map is an  $F$ -biharmonic map.

**Proposition 2.4.** Let  $u: M \rightarrow N$  be a smooth map. If  $\|\tau(u)\|^2$  is constant, then  $u$  is  $F$ -biharmonic if and only if it is biharmonic.



Go back

Full Screen

Close

Quit



*Proof.* Since  $\|\tau(u)\|^2$  is constant, we have

$$\begin{aligned}\tau_{F,2}(u) &= F' \left( \frac{\|\tau(u)\|^2}{2} \right) \left[ -\tilde{\Delta}(\tau(u)) - \sum_i R^N(du(e_i), \tau(u))du(e_i) \right] \\ &= F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau_2(u),\end{aligned}$$

so we know that  $u$  is  $F$ -biharmonic if and only if it is biharmonic. □

*Remark 2.5.* When  $\|\tau(u)\|^2$  is non-constant, we have

$$\begin{aligned}\tau_{F,2}(u) &= F' \left( \frac{\|\tau(u)\|^2}{2} \right) \left[ -\tilde{\Delta}(\tau(u)) - \sum_i R^N(du(e_i), \tau(u))du(e_i) \right] \\ &\quad - \left[ \tilde{\Delta}F' \left( \frac{\|\tau(u)\|^2}{2} \right) \right] \tau(u) + \tilde{\nabla}_{\text{grad } F' \left( \frac{\|\tau(u)\|^2}{2} \right)} \tau(u) \\ &= F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau_2(u) - \left[ \tilde{\Delta}F' \left( \frac{\|\tau(u)\|^2}{2} \right) \right] \tau(u) + \tilde{\nabla}_{\text{grad } F' \left( \frac{\|\tau(u)\|^2}{2} \right)} \tau(u).\end{aligned}$$

From this equation, we know that there are many differences between  $F$ -biharmonic maps and biharmonic maps when  $F(t) = (2t)^{\frac{p}{2}}$ , ( $p > 2$ ) or  $F(t) = e^t$ .

### 3. NON-EXISTENCE RESULTS FOR $F$ -BIHARMONIC MAPS

From the definition of an  $F$ -biharmonic map, we know that a harmonic map is  $F$ -biharmonic map, so a basic question in theory is to understand under what conditions the converse is true. A first general answer to this problem for  $F(t) = t$ , proved by G. Y. Jiang [9], is the following theorem



Go back

Full Screen

Close

Quit

**Theorem 3.1** ([9]). *Let  $u: (M, g) \rightarrow (N, h)$  be a smooth map. If  $M$  is compact, orientable and the sectional curvature of  $(N, h)$  is non-positive, i.e.,  $\text{Riem}^N \leq 0$ , then  $u$  is a biharmonic map if and only if it is harmonic.*

In this section, we will obtain the following results

**Theorem 3.2.** *Let  $u: (M, g) \rightarrow (N, h)$  be a smooth map. If  $M$  is compact, orientable and the sectional curvature of  $(N, h)$  is non-positive, i.e.,  $\text{Riem}^N \leq 0$ , then  $u$  is an  $F$ -biharmonic map if and only if it is harmonic.*

*Proof.* Computing the Laplacian of the function  $\|F'(\frac{\|\tau(u)\|^2}{2})\tau(u)\|^2$ , we have

$$\begin{aligned}
 & \Delta \left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2 \\
 (6) \quad & = 2 \sum_k h \left( \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right], \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \\
 & \quad + 2h \left( -\tilde{\Delta} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right], F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right).
 \end{aligned}$$

Since  $u$  is an  $F$ -biharmonic map, we have

$$\begin{aligned}
 (7) \quad \tau_{F,2}(u) & = -\tilde{\Delta} \left( F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right) \\
 & \quad - \sum_i R^N \left( du(e_i), F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right) du(e_i) = 0.
 \end{aligned}$$



Go back

Full Screen

Close

Quit

From (6) and (7), we have

$$\begin{aligned}
 & \Delta \left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2 \\
 &= 2 \sum_k h(\tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right], \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right]) \\
 (8) \quad &+ 2 \sum_i h(R^N \left( du(e_i), F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right) du(e_i), F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u)).
 \end{aligned}$$

Since the section curvature of  $N$  is non-positive, i.e.,  $\text{Riem}^N \leq 0$  and by (8), we have

$$(9) \quad \Delta \left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2 \geq 0$$

By the Green's theorem  $\int_M \Delta \left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2 dv_g = 0$  and (9), we have

$$\Delta \left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2 = 0,$$

so then  $\left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2$  is constant. From (8), we have

$$(10) \quad \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] = 0, \quad \text{for } k = 1, \dots, m.$$




Go back

Full Screen

Close

Quit



Setting  $X = \sum_i h \left( du(e_i), F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right) e_i$ , we have

$$\begin{aligned}
 \operatorname{div}(X) &= \sum_k g(\nabla_{e_k} X, e_k) \\
 &= h \left( \tau(u), F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right) \\
 (11) \quad &+ \sum_i h \left( du(e_i), \tilde{\nabla}_{e_i} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \\
 &= h \left( \tau(u), F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right) \\
 &= F' \left( \frac{\|\tau(u)\|^2}{2} \right) \|\tau(u)\|^2.
 \end{aligned}$$

Integrating (11) over  $M$ , we have

$$(12) \quad 0 = \int_M \operatorname{div}(X) dv_g = \int_M F' \left( \frac{\|\tau(u)\|^2}{2} \right) \|\tau(u)\|^2 dv_g.$$

From  $F'(t) > 0$  on  $(0, \infty)$  and (12), we have  $\tau(u) = 0$ . □

When  $u$  is a Riemannian immersion and  $\dim M = \dim N - 1$ , we can replace the hypothesis  $\operatorname{Riem}^N \leq 0$  with the hypothesis  $\operatorname{Ricci}^N \leq 0$ , and we obtain the following theorem.

**Theorem 3.3.** *Let  $u: (M, g) \rightarrow (N, h)$  be a Riemannian immersion. If  $M$  is compact, orientable,  $\operatorname{Ricci}^N \leq 0$  and  $\dim M = \dim N - 1$ , then  $u$  is an  $F$ -biharmonic map if and only if it is harmonic.*



Go back

Full Screen

Close

Quit

*Proof.* Since  $u$  is a Riemannian immersion and  $\dim M = \dim N - 1$ , we have

$$\begin{aligned}
 (13) \quad & \sum_i h \left( R^N \left( du(e_i), F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right) du(e_i), F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right) \\
 & = -\text{Ricci}^N \left( F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u), F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right).
 \end{aligned}$$

From (8), (13) and  $\text{Ricci}^N \leq 0$ , we have

$$\Delta \left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2 \geq 0.$$

Applying the same argument as in the proof of Theorem 3.2, we get the result.  $\square$

**Theorem 3.4.** *Let  $(M, g)$  be an  $m$ -dimensional complete manifold with  $\text{Vol}(M, g) = \infty$ . If  $u: (M, g) \rightarrow (N, h)$  is an  $F$ -biharmonic map, the sectional curvature of  $(N, h)$  is non-positive, i.e.,  $\text{Riem}^N \leq 0$  and  $\int_M \left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2 dv_g < \infty$ , then  $u$  is harmonic.*

*Proof.* Since  $u$  is an  $F$ -biharmonic map, we have

$$\begin{aligned}
 (14) \quad \tau_{F,2}(u) & = -\tilde{\Delta} \left( F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right) \\
 & \quad - \sum_i R^N \left( du(e_i), F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right) du(e_i) = 0.
 \end{aligned}$$



Go back

Full Screen

Close

Quit

Take any point  $x_0 \in M$  and for every  $r > 0$ , let us consider the following cut off function  $\lambda(x)$  on  $M$ :

$$(15) \quad \begin{cases} 0 \leq \lambda(x) \leq 1, & x \in M, \\ \lambda(x) = 1, & x \in B_r(x_0), \\ \lambda(x) = 0, & x \in M - B_{2r}(x_0), \\ |\nabla \lambda| \leq \frac{2}{r}, & x \in M, \end{cases}$$

where  $B_r(x_0) = \{x \in M : d(x, x_0) < r\}$  and  $d$  is the distance of  $(M, g)$ .

Let  $X$  be a compactly supported vector field on  $M$  such that

$$g(X, Y) = h \left( \tilde{\nabla}_Y \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right], \lambda^2 \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right).$$

Then the divergence of  $X$  is given by the following expression

$$(16) \quad \begin{aligned} & \operatorname{div}(X) \\ &= \sum_k g(\nabla_{e_k} X, e_k) = \sum_k e_k g(X, e_k) - \sum_k g(X, \nabla_{e_k} e_k) \\ &= \sum_k e_k h \left( \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right], \lambda^2 \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \\ &\quad - \sum_k h \left( \tilde{\nabla}_{\nabla_{e_k} e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right], \lambda^2 \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \\ &= h \left( -\tilde{\Delta} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right], \lambda^2 \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \\ &\quad + \sum_k h \left( \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right], \tilde{\nabla}_{e_k} \left( \lambda^2 \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \right). \end{aligned}$$



Go back

Full Screen

Close

Quit

From (14) and (16), we have

$$\begin{aligned}
 & \operatorname{div}(X) \\
 (17) \quad &= \sum_k h \left( R^N \left( du(e_k), F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right) du(e_k), \lambda^2 \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \\
 & \quad + \sum_k h \left( \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right], \tilde{\nabla}_{e_k} \left( \lambda^2 \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \right).
 \end{aligned}$$

Integrating (17) over  $M$  and  $\operatorname{Riem}^N \leq 0$ , we get

$$\begin{aligned}
 & \sum_k \int_M h \left( \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right], \tilde{\nabla}_{e_k} \left( \lambda^2 \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \right) dv_g \\
 (18) \quad &= - \sum_k \int_M h \left( R^N \left( du(e_k), F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right) du(e_k), \lambda^2 \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) dv_g \\
 &= \sum_k \int_M h \left( R^N \left( F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u), du(e_k) \right) du(e_k), \lambda^2 \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) dv_g \\
 &\leq 0.
 \end{aligned}$$

From (18), we have

$$\begin{aligned}
 & 0 \geq \sum_k \int_M h \left( \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right], \tilde{\nabla}_{e_k} \left( \lambda^2 \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \right) dv_g \\
 (19) \quad &= \sum_k \int_M \lambda^2 \left\| \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right\|^2 dv_g \\
 & \quad + 2 \sum_k \int_M \lambda e_k(\lambda) h \left( \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right], \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) dv_g.
 \end{aligned}$$



Go back

Full Screen

Close

Quit

Therefore, we have

$$\begin{aligned}
 & \sum_k \int_M \lambda^2 \left\| \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right\|^2 dv_g \\
 & \leq -2 \sum_k \int_M \lambda e_k(\lambda) h \left( \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right], \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) dv_g \\
 (20) \quad & = - \sum_k \int_M 2h \left( \lambda \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right], e_k(\lambda) \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) dv_g \\
 & \leq \sum_k \int_M \left\{ \frac{1}{2} \lambda^2 \left\| \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right\|^2 + 2[e_k(\lambda)]^2 \left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2 \right\} dv_g,
 \end{aligned}$$

where we use the following Cauchy-Schwarz inequality

$$\pm 2h(V, W) \leq \varepsilon \|V\|^2 + \frac{1}{\varepsilon} \|W\|^2$$

for the second inequality and  $\varepsilon = \frac{1}{2}$ .

From (20), we have

$$\begin{aligned}
 (21) \quad & \sum_k \int_M \lambda^2 \left\| \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right\|^2 dv_g \leq 4 \int_M \sum_k [e_k(\lambda)]^2 \left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2 dv_g, \\
 & \leq \frac{16}{r^2} \int_M \left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2 dv_g.
 \end{aligned}$$

Since  $\int_M \left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2 dv_g < \infty$  and  $(M, g)$  is complete, then we have  $(r \rightarrow \infty)$

$$\int_M \sum_k \left\| \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right\|^2 dv_g = 0.$$




Go back

Full Screen

Close

Quit



For every vector field  $X$  on  $M$ , we have

$$\tilde{\nabla}_X \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] = 0.$$

So we know that  $\left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2$  is constant, say  $C$ . Therefore, if  $\text{Vol}(M, g) = \infty$  and  $C \neq 0$ , then

$$\int_M \left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2 dv_g = C^2 \text{Vol}(M, g) = \infty,$$

which yields a contradiction. Thus, we have  $\left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2 = C = 0$ . From  $F'(t) > 0$  on  $(0, \infty)$  and  $\left\| F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right\|^2 = 0$ , we know that  $\tau(u) = 0$ , i.e.  $u$  is harmonic.  $\square$

From Theorem 3.4, we have the following corollaries:

**Corollary 3.5.** *Let  $(M, g)$  be an  $m$ -dimensional complete manifold with  $\text{Vol}(M, g) = \infty$ . If  $u: (M, g) \rightarrow (N, h)$  is an exponentially biharmonic map, the sectional curvature of  $(N, h)$  is non-positive, i.e.,*

$$\text{Riem}^N \leq 0 \quad \text{and} \quad \int_M \|\tau(u)\|^2 e^{\|\tau(u)\|^2} dv_g < \infty,$$

then  $u$  is harmonic.

**Corollary 3.6.** *Let  $(M, g)$  be an  $m$ -dimensional complete manifold with  $\text{Vol}(M, g) = \infty$ . If  $u: (M, g) \rightarrow (N, h)$  is a  $p$ -biharmonic map, the sectional curvature of  $(N, h)$  is non-positive, i.e.,  $\text{Riem}^N \leq 0$  and  $\int_M \|\tau(u)\|^{2p-2} dv_g < \infty$ , then  $u$  is harmonic.*



Go back

Full Screen

Close

Quit



**Corollary 3.7 ([15]).** *Let  $(M, g)$  be an  $m$ -dimensional complete manifold with  $\text{Vol}(M, g) = \infty$ . If  $u: (M, g) \rightarrow (N, h)$  is a biharmonic map, the sectional curvature of  $(N, h)$  is non-positive, i.e.,  $\text{Riem}^N \leq 0$  and  $\int_M \|\tau(u)\|^2 dv_g < \infty$ , then  $u$  is harmonic.*

#### 4. STRESS $F$ -BIENERGY TENSOR

The stress bienergy tensor and the conservation law of a biharmonic map between Riemannian manifolds were first studied by G.Y. Jiang in [10]. Following Jiang's notion, we define the stress  $F$ -bienergy tensor of a smooth map as follows.

**Definition 4.1.** Let  $u: (M, g) \rightarrow (N, h)$  be a smooth map between two Riemannian manifolds. The stress  $F$ -bienergy tensor of  $u$  is defined by

$$S_{F,2}(X, Y) = F\left(\frac{\|\tau(u)\|^2}{2}\right)g(X, Y) + \sum_k h\left(du(e_k), \tilde{\nabla}_{e_k}\left[F'\left(\frac{\|\tau(u)\|^2}{2}\right)\tau(u)\right]\right)g(X, Y) \\ - h\left(du(X), \tilde{\nabla}_Y\left[F'\left(\frac{\|\tau(u)\|^2}{2}\right)\tau(u)\right]\right) - h\left(du(Y), \tilde{\nabla}_X\left[F'\left(\frac{\|\tau(u)\|^2}{2}\right)\tau(u)\right]\right)$$

for any  $X, Y \in \Gamma(TM)$ .

*Remark 4.2.* When  $F(t) = t$ , we have  $S_{F,2}(X, Y) = S_2(X, Y)$ , where  $S_2$  is stress bienergy tensor in [10].

**Theorem 4.3.** *For any smooth map  $u: (M, g) \rightarrow (N, h)$*

$$(\text{div } S_{F,2})(X) = -h(\tau_{F,2}(u), du(X)) - F''\left(\frac{\|\tau(u)\|^2}{2}\right)X\left(\frac{\|\tau(u)\|^4}{4}\right)$$

for any vector field  $X \in \Gamma(TM)$ .

*Proof.* We choose a local orthonormal frame field  $\{e_i\}$  on  $M$  with  $\nabla_{e_i} e_i|_x = 0$  at a point  $x \in M$ . Let  $X$  be a vector field on  $M$ . At  $x$ , we compute



Go back

Full Screen

Close

Quit

$$\begin{aligned}
(\operatorname{div} S_{F,2})(X) &= \sum_i (\nabla_{e_i} S_{F,2})(e_i, X) = \sum_i e_i S_{F,2}(e_i, X) - S_{F,2}(e_i, \nabla_{e_i} X) \\
&= \sum_i e_i \left[ F \left( \frac{\|\tau(u)\|^2}{2} \right) g(e_i, X) + \sum_k h \left( du(e_k), \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) g(e_i, X) \right. \\
&\quad \left. - h \left( du(e_i), \tilde{\nabla}_X \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) - h \left( du(X), \tilde{\nabla}_{e_i} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \right] \\
&\quad - \sum_i \left[ F \left( \frac{\|\tau(u)\|^2}{2} \right) g(e_i, \nabla_{e_i} X) + \sum_k h \left( du(e_k), \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) g(e_i, \nabla_{e_i} X) \right. \\
&\quad \left. - h \left( du(e_i), \tilde{\nabla}_{\nabla_{e_i} X} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) - h \left( du(\nabla_{e_i} X), \tilde{\nabla}_{e_i} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \right] \\
&= X \left( F \left( \frac{\|\tau(u)\|^2}{2} \right) \right) + \sum_k h \left( (\tilde{\nabla} du)(X, e_k), \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \\
&\quad + \sum_k h \left( du(e_k), \tilde{\nabla}_X \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) - h \left( \tau(u), \tilde{\nabla}_X \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \\
&\quad - \sum_i h \left( du(e_i), \tilde{\nabla}_{e_i} \tilde{\nabla}_X \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) - \sum_k h \left( (\tilde{\nabla} du)(X, e_k), \tilde{\nabla}_{e_k} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \\
&\quad + \sum_i h \left( du(e_i), \tilde{\nabla}_{\nabla_{e_i} X} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) - \sum_i h \left( du(X), \tilde{\nabla}_{e_i} \tilde{\nabla}_{e_i} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) \\
&= -F'' \left( \frac{\|\tau(u)\|^2}{2} \right) X \left( \frac{\|\tau(u)\|^4}{4} \right) \\
&\quad + h \left( \tilde{\Delta} \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] + \sum_i R^N \left( du(e_i), \left[ F' \left( \frac{\|\tau(u)\|^2}{2} \right) \tau(u) \right] \right) du(e_i), du(X) \right) \\
&= -h(\tau_{F,2}(u), du(X)) - F'' \left( \frac{\|\tau(u)\|^2}{2} \right) X \left( \frac{\|\tau(u)\|^4}{4} \right).
\end{aligned}$$




Go back

Full Screen

Close

Quit



□

From Theorem 3.1, we know that if  $u: M \rightarrow N$  is an  $F$ -biharmonic map, then

$$(22) \quad (\operatorname{div} S_{F,2})(X) = -F''\left(\frac{\|\tau(u)\|^2}{2}\right)X\left(\frac{\|\tau(u)\|^4}{4}\right).$$

**Proposition 4.4.** *Let  $c: I \subset \mathbb{R} \rightarrow (N, h)$  be a curve parametrized by arc-length. Assume that  $S_{F,2} = 0$  and  $l_F = \inf_{t \geq 0} \frac{tF'(t)}{F(t)} > 0$ . Then  $c$  is geodesic.*

*Proof.* A direct computation shows that

$$\begin{aligned} 0 &= S_{F,2}\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right) = F\left(\frac{\|\tau(c)\|^2}{2}\right) - h\left(dc\left(\frac{\partial}{\partial t}\right), \tilde{\nabla}_{\frac{\partial}{\partial t}}\left[F'\left(\frac{\|\tau(c)\|^2}{2}\right)\tau(c)\right]\right) \\ &= F\left(\frac{\|\tau(c)\|^2}{2}\right) + h\left(\tau(c), \left[F'\left(\frac{\|\tau(c)\|^2}{2}\right)\tau(c)\right]\right), > (1 + 2l_F)F\left(\frac{\|\tau(c)\|^2}{2}\right). \end{aligned}$$

If  $F\left(\frac{\|\tau(c)\|^2}{2}\right) = 0$ , then  $\tau(c) = 0$ . □

**Proposition 4.5.** *Let  $u: (M^2, g) \rightarrow (N, h)$  be a map from a surface. Then  $S_{F,2} = 0$  implies  $u$  is harmonic.*

*Proof.* The trace of  $S_{F,2}$  gives the equality

$$\begin{aligned} 0 &= \operatorname{trace} S_{F,2} = F\left(\frac{\|\tau(u)\|^2}{2}\right) + 2\left\langle du, \tilde{\nabla}\left[F'\left(\frac{\|\tau(u)\|^2}{2}\right)\tau(u)\right] \right\rangle \\ &\quad - 2\left\langle du, \tilde{\nabla}\left[F'\left(\frac{\|\tau(u)\|^2}{2}\right)\tau(u)\right] \right\rangle = F\left(\frac{\|\tau(u)\|^2}{2}\right), \end{aligned}$$

so we have  $\tau(u) = 0$ . □



Go back

Full Screen

Close

Quit

**Proposition 4.6.** *Let  $u: (M^m, g) \rightarrow (N, h)$ ,  $m \neq 2$ . Then  $S_{F,2} = 0$  if and only if*

$$(23) \quad \begin{aligned} & \frac{2}{m-2} F\left(\frac{\|\tau(u)\|^2}{2}\right) g(X, Y) + h\left(du(X), \tilde{\nabla}_Y \left[F'\left(\frac{\|\tau(u)\|^2}{2}\right) \tau(u)\right]\right) \\ & + h\left(du(Y), \tilde{\nabla}_X \left[F'\left(\frac{\|\tau(u)\|^2}{2}\right) \tau(u)\right]\right) = 0 \end{aligned}$$

for any  $X, Y \in \Gamma(TM)$ .

*Proof.* Since  $S_{F,2} = 0$ , we have  $\text{trace} S_{F,2} = 0$ . Therefore,

$$\sum_k h\left(du(e_k), \tilde{\nabla}_{e_k} \left[F'\left(\frac{\|\tau(u)\|^2}{2}\right) \tau(u)\right]\right) = -\frac{m}{m-2} F\left(\frac{\|\tau(u)\|^2}{2}\right).$$

Substituting it into the definition of  $S_{F,2}$ , we obtain

$$\begin{aligned} 0 = S_{F,2}(X, Y) &= -\frac{2}{m-2} F\left(\frac{\|\tau(u)\|^2}{2}\right) g(X, Y) \\ & - h\left(du(X), \tilde{\nabla}_Y \left[F'\left(\frac{\|\tau(u)\|^2}{2}\right) \tau(u)\right]\right) - h\left(du(Y), \tilde{\nabla}_X \left[F'\left(\frac{\|\tau(u)\|^2}{2}\right) \tau(u)\right]\right). \end{aligned}$$

□

**Proposition 4.7.** *A map  $u: (M^m, g) \rightarrow (N, h)$ ,  $m > 2$ , with  $S_{F,2} = 0$  and  $\text{rank } u \leq m - 1$  is harmonic.*

*Proof.* Take  $p \in M$ . Since  $\text{rank } u(p) \leq m - 1$ , there exists a unit vector  $X_p \in \text{Ker } du_p$  and for  $X = Y = X_p$ , (23) becomes  $F\left(\frac{\|\tau(u)\|^2}{2}\right) = 0$ , so  $\tau(u) = 0$ . □

**Corollary 4.8.** *Let  $u: (M^m, g) \rightarrow (N^n, h)$  be a submersion ( $m > n$ ), if  $S_{F,2} = 0$ , then  $u$  is harmonic.*




Go back

Full Screen

Close

Quit



Recall that for two 2-tensors  $T_1, T_2 \in \Gamma(T^*M \otimes T^*M)$ , their inner product is defined as follows:

$$(24) \quad \langle T_1, T_2 \rangle = \sum_{ij} T_1(e_i, e_j) T_2(e_i, e_j),$$

where  $\{e_i\}$  is an orthonormal basis of  $M$  with respect to  $g$ . For a vector field  $X \in \Gamma(TM)$ , by  $\theta_X$  we denote its dual one form, i.e.,  $\theta_X(Y) = g(X, Y)$ . The covariant derivative of  $\theta_X$  gives a 2-tensor field  $\nabla\theta_X$

$$(25) \quad (\nabla\theta_X)(Y, Z) = (\nabla_Z\theta_X)(Y) = g(\nabla_Z X, Y).$$

If  $X = \nabla\varphi$  is the gradient of some function  $\varphi$  on  $M$ , then  $\theta_X = d\varphi$  and  $\nabla\theta_X = \text{Hess } \varphi$ .

**Lemma 4.9** (cf. [2, 4]). *Let  $T$  be a symmetric  $(0, 2)$ -type tensor field and let  $X$  be a vector field. Then*

$$(26) \quad \text{div}(i_X T) = (\text{div } T)(X) + \langle T, \nabla\theta_X \rangle = (\text{div } T)(X) + \frac{1}{2} \langle T, L_X g \rangle.$$

Let  $D$  be any bounded domain of  $M$  with  $C^1$  boundary. By using the Stokes' theorem, we immediately have the following integral formula

$$(27) \quad \int_{\partial D} T(X, \nu) ds_g = \int_D [\langle T, \frac{1}{2} L_X g \rangle + \text{div}(T)(X)] dv_g$$

where  $\nu$  is the unit outward normal vector field along  $\partial D$ .

By (22) and (3), we have

$$(28) \quad \int_{\partial D} S_{F,2}(X, \nu) ds_g = \int_D \left[ \langle S_{F,2}, \frac{1}{2} L_X g \rangle - F'' \left( \frac{\|\tau(u)\|^2}{2} \right) X \left( \frac{\|\tau(u)\|^4}{4} \right) \right] dv_g.$$




Go back

Full Screen

Close

Quit



When  $F(t) = t$ , the equation (28) turns into the following equation

$$(29) \quad \int_{\partial D} S_2(X, \nu) ds_g = \int_D \langle S_2, \frac{1}{2} L_X g \rangle dv_g.$$

## 5. MONOTONICITY FORMULAS FOR BIHARMONIC MAPS

In this section, we investigate the special case of  $F$ -biharmonic maps, i.e., biharmonic maps.

Let  $(M^m, g)$  be a complete Riemannian manifold with pole  $x_0$ . By  $r(x)$  denote the  $g$ -distance function relative to the pole  $x_0$ , that is,  $r(x) = \text{dist}_g(x, x_0)$ . Set  $B(r) = \{x \in M^m : r(x) \leq r\}$ . By  $\lambda_{\max}$  (resp.  $\lambda_{\min}$ ) denote the maximum (resp. minimal) eigenvalues of  $\text{Hess}(r^2) - dr \otimes dr$  at each point of  $M - \{x_0\}$ .

**Theorem 5.1.** *Let  $u: (M, g) \rightarrow (N, h)$  be an isometric immersion. Assume that there is a constant  $\sigma > 0$  such that*

$$(30) \quad \frac{m-1}{2} \lambda_{\min} + 1 - 2 \max\{2, \lambda_{\max}\} \geq \sigma.$$

*If  $u$  is a biharmonic map and  $h(\tau(u), \tilde{\nabla}_{\frac{\partial}{\partial r}} du(\frac{\partial}{\partial r})) \geq 0$ , then we have*

$$(31) \quad \frac{\int_{B(\rho_1)} \frac{\|\tau(u)\|^2}{2} dv_g}{\rho_1^\sigma} \leq \frac{\int_{B(\rho_2)} \frac{\|\tau(u)\|^2}{2} dv_g}{\rho_2^\sigma}$$

*for any  $0 < \rho_1 \leq \rho_2$ .*

*Proof.* Since  $u: M^m \rightarrow N$  is an isometric immersion, we have  $\tau(u) = mH$ , where  $H$  is the mean curvature vector field of  $M$  in  $N$ , so we know that

$$(32) \quad h(\tau(u), du(X)) = h(mH, du(X)) = 0$$



Go back

Full Screen

Close

Quit

for any tangent vector field  $X$  on  $M$ .

Taking  $D = B(r)$  and  $X = r \frac{\partial}{\partial r}$  in (29), we have

$$(33) \quad \int_{\partial B(r)} S_2 \left( r \frac{\partial}{\partial r}, \frac{\partial}{\partial r} \right) ds_g = \int_{B(r)} \langle S_2, \frac{1}{2} L_{r \frac{\partial}{\partial r}} g \rangle dv_g = \frac{1}{2} \int_{B(r)} \langle S_2, \text{Hess}(r^2) \rangle dv_g.$$

Let  $\{e_i\}_{i=1}^m$  be an orthonormal basis on  $M$  and  $e_m = \frac{\partial}{\partial r}$ . We may assume that  $\text{Hess}(r^2)$  becomes a diagonal matrix with respect to  $\{e_i\}$ .

$$(34) \quad \begin{aligned} -\frac{1}{2} \langle S_2, \text{Hess}(r^2) \rangle &= -\frac{1}{2} \sum_{i,j} S_2(e_i, e_j) \text{Hess}(r^2)(e_i, e_j) \\ &= -\frac{1}{2} \left\{ \sum_i \frac{\|\tau(u)\|^2}{2} \text{Hess}(r^2)(e_i, e_i) \right. \\ &\quad + \sum_k h(\tilde{\nabla}_{e_k} \tau(u), du(e_k)) \sum_i \text{Hess}(r^2)(e_i, e_i) \\ &\quad \left. - 2 \sum_{i,j} h(du(e_i), \tilde{\nabla}_{e_j} \tau(u)) \text{Hess}(r^2)(e_i, e_j) \right\} \\ &= -\frac{1}{2} \left\{ -\frac{\|\tau(u)\|^2}{2} \sum_i \text{Hess}(r^2)(e_i, e_i) \right. \\ &\quad \left. + 2 \sum_i h(\tau(u), \tilde{\nabla}_{e_i} du(e_i)) \text{Hess}(r^2)(e_i, e_i) \right\} \\ &\geq \frac{\|\tau(u)\|^2}{2} \left[ \frac{m-1}{2} \lambda_{\min} + 1 - 2 \max\{2, \lambda_{\max}\} \right] \\ &\geq \sigma \frac{\|\tau(u)\|^2}{2}, \end{aligned}$$

where the equation (32) is used for the third equality and the equation (30) for the last inequality.



Go back

Full Screen

Close

Quit

On the other hand, by the coarea formula, we have

$$\begin{aligned}
 (35) \quad - \int_{\partial B(r)} S_2 \left( r \frac{\partial}{\partial r}, \frac{\partial}{\partial r} \right) ds_g &= - \int_{\partial B(r)} \left\{ \left[ \frac{\|\tau(u)\|^2}{2} + \sum_k h(du(e_k), \tilde{\nabla}_{e_k} \tau(u)) \right] g \left( r \frac{\partial}{\partial r}, \frac{\partial}{\partial r} \right) \right. \\
 &\quad \left. - 2rh \left( du \left( \frac{\partial}{\partial r} \right), \tilde{\nabla}_{\frac{\partial}{\partial r}} \tau(u) \right) \right\} ds_g \\
 &= \int_{\partial B(r)} \left\{ r \frac{\|\tau(u)\|^2}{2} - rh \left( \tau(u), \tilde{\nabla}_{\frac{\partial}{\partial r}} du \left( \frac{\partial}{\partial r} \right) \right) \right\} ds_g \\
 &\leq \int_{\partial B(r)} r \frac{\|\tau(u)\|^2}{2} ds_g = r \frac{d}{dr} \int_{B(r)} \frac{\|\tau(u)\|^2}{2} dv_g,
 \end{aligned}$$

where the condition  $h(\tau(u), \tilde{\nabla}_{\frac{\partial}{\partial r}} du(\frac{\partial}{\partial r})) \geq 0$  is used for the inequality.

From (33), (34) and (35), we have

$$(36) \quad \sigma \int_{B(r)} \frac{\|\tau(u)\|^2}{2} dv_g \leq r \frac{d}{dr} \int_{B(r)} \frac{\|\tau(u)\|^2}{2} dv_g$$

i.e.

$$(37) \quad \frac{d}{dr} \frac{\int_{B(r)} \frac{\|\tau(u)\|^2}{2} dv_g}{r^\sigma} \geq 0.$$

Therefore,

$$\frac{\int_{B(\rho_1)} \frac{\|\tau(u)\|^2}{2} dv_g}{\rho_1^\sigma} \leq \frac{\int_{B(\rho_2)} \frac{\|\tau(u)\|^2}{2} dv_g}{\rho_2^\sigma}$$

for any  $0 < \rho_1 \leq \rho_2$ . □




Go back

Full Screen

Close

Quit



**Lemma 5.2** ([4, 8]). Let  $(M^m, g)$  be a complete Riemannian manifold with a pole  $x_0$ . By  $K_r$  denote the radial curvature of  $M$  as follows

(i) if  $-\alpha^2 \leq K_r \leq -\beta^2$  with  $\alpha \geq \beta > 0$ , then

$$\beta \coth(\beta r)[g - dr \otimes dr] \leq \text{Hess}(r) \leq \alpha \coth(\alpha r)[g - dr \otimes dr],$$

(ii) if  $-\frac{A}{(1+r^2)^{1+\varepsilon}} \leq K_r \leq \frac{B}{(1+r^2)^{1+\varepsilon}}$  with  $\varepsilon > 0$ ,  $A \geq 0$  and  $0 \leq B < 2\varepsilon$ , then

$$\frac{1 - B/2\varepsilon}{r}[g - dr \otimes dr] \leq \text{Hess}(r) \leq \frac{e^{A/2\varepsilon}}{r}[g - dr \otimes dr],$$

(iii) if  $-\frac{a^2}{1+r^2} \leq K_r \leq \frac{b^2}{1+r^2}$  with  $a \geq 0$  and  $b^2 \in [0, \frac{1}{4}]$ , then

$$\frac{1 + \sqrt{1 - 4b^2}}{2r}[g - dr \otimes dr] \leq \text{Hess}(r) \leq \frac{1 + \sqrt{1 + 4a^2}}{2r}[g - dr \otimes dr].$$

**Lemma 5.3.** Let  $(M^m, g)$  be a complete Riemannian manifold with a pole  $x_0$ . By  $K_r$  denote the radial curvature of  $M$  as follows

(i) if  $-\alpha^2 \leq K_r \leq -\beta^2$  with  $\alpha \geq \beta > 0$  and  $(m-1)\beta - 4\alpha \geq 0$ , then

$$\frac{(m-1)}{2}\lambda_{\min} + 1 - 2\max\{2, \lambda_{\max}\} \geq m - \frac{4\alpha}{\beta}.$$

(ii) if  $-\frac{A}{(1+r^2)^{1+\varepsilon}} \leq K_r \leq \frac{B}{(1+r^2)^{1+\varepsilon}}$  with  $\varepsilon > 0$ ,  $A \geq 0$  and  $0 \leq B < 2\varepsilon$ , then

$$\frac{(m-1)}{2}\lambda_{\min} + 1 - 2\max\{2, \lambda_{\max}\} \geq 1 + (m-1)\left(1 - \frac{B}{2\varepsilon}\right) - 4e^{\frac{A}{2\varepsilon}}.$$




Go back

Full Screen

Close

Quit



(iii) if  $-\frac{a^2}{1+r^2} \leq K_r \leq \frac{b^2}{1+r^2}$  with  $a \geq 0$  and  $b^2 \in [0, \frac{1}{4}]$ , then

$$\begin{aligned} & \frac{(m-1)}{2} \lambda_{\min} + 1 - 2 \max\{2, \lambda_{\max}\} \\ & \geq [1 + (m-1) \frac{1 + \sqrt{1-4b^2}}{2} - 4 \frac{1 + \sqrt{1+4a^2}}{2}]. \end{aligned}$$

*Proof.* If  $K_r$  satisfies (i), then by Lemma 5.2, for every  $r > 0$ , we have on  $B(r) - \{x_0\}$ ,

$$\begin{aligned} & \frac{1}{2} [(m-1) \lambda_{\min} + 2 - 4 \max\{2, \lambda_{\max}\}] \\ & \geq \frac{1}{2} [(m-1) 2\beta r \coth(\beta r) + 2 - 4 \times 2\alpha r \coth(\alpha r)] \\ & = 1 + \beta r \coth(\beta r) \left( m - 1 - \frac{4\alpha}{\beta} \frac{\coth(\alpha r)}{\coth(\beta r)} \right) \\ & \geq 1 + 1 \cdot \left( m - 1 - \frac{4\alpha}{\beta} \right) = m - \frac{4\alpha}{\beta}. \end{aligned}$$

where the second inequality is valid the increasing function  $\beta r \coth(\beta r) \rightarrow 1$  as  $r \rightarrow 0$ , and  $\frac{\coth(\alpha r)}{\coth(\beta r)} < 1$  for  $0 < \beta < \alpha$ . Similarly, from Lemma 5.2, the above inequality holds for the cases (ii) and (iii) on  $B(r)$ . □

**Theorem 5.4.** Let  $(M, g)$  be an  $m$ -dimensional complete manifold with a pole  $x_0$ . Assume that the radial curvature  $K_r$  of  $M$  satisfies one of the following three conditions:

- (i) if  $-\alpha^2 \leq K_r \leq -\beta^2$  with  $\alpha \geq \beta > 0$  and  $(m-1)\beta - 4\alpha \geq 0$ ,
- (ii) if  $-\frac{A}{(1+r^2)^{1+\varepsilon}} \leq K_r \leq \frac{B}{(1+r^2)^{1+\varepsilon}}$  with  $\varepsilon > 0$ ,  $A \geq 0$ ,  $0 \geq B < 2\varepsilon$  and  $1 + (m-1)(1 - \frac{B}{2\varepsilon}) - 4e^{\frac{A}{2\varepsilon}} > 0$ ,



Go back

Full Screen

Close

Quit

(iii) if  $-\frac{a^2}{1+r^2} \leq K_r \leq \frac{b^2}{1+r^2}$  with  $a \geq 0$ ,  $b^2 \in [0, \frac{1}{4}]$  and  $1 + (m-1)\frac{1+\sqrt{1-4b^2}}{2} - 4\frac{1+\sqrt{1+4a^2}}{2} > 0$ .

If  $u: (M, g) \rightarrow (N, h)$  is a biharmonic isometric immersion and  $h(\tau(u), \tilde{\nabla}_{\frac{\partial}{\partial r}} du(\frac{\partial}{\partial r})) \geq 0$ , then

$$(38) \quad \frac{\int_{B(\rho_1)} \frac{\|\tau(u)\|^2}{2} dv_g}{\rho_1^\Lambda} \leq \frac{\int_{B(\rho_2)} \frac{\|\tau(u)\|^2}{2} dv_g}{\rho_2^\Lambda}$$

for any  $0 < \rho_1 \leq \rho_2$ , where

$$(39) \quad \Lambda = \begin{cases} m - \frac{4\alpha}{\beta}, & \text{if } K_r \text{ satisfies (i)} \\ 1 + (m-1)\left(1 - \frac{B}{2\varepsilon}\right) - 4e^{\frac{A}{2\varepsilon}}, & \text{if } K_r \text{ satisfies (ii)} \\ 1 + (m-1)\frac{1+\sqrt{1-4b^2}}{2} - 4\frac{1+\sqrt{1+4a^2}}{2}, & \text{if } K_r \text{ satisfies (iii)} \end{cases}$$

*Proof.* From the proof of Theorem 5.1 and Lemma 5.3, we have

$$\frac{d}{dr} \frac{\int_{B(r)} \frac{\|\tau(u)\|^2}{2} dv_g}{r^\Lambda} \geq 0.$$

Therefore, we get the monotonicity formula

$$\frac{\int_{B(\rho_1)} \frac{\|\tau(u)\|^2}{2} dv_g}{\rho_1^\Lambda} \leq \frac{\int_{B(\rho_2)} \frac{\|\tau(u)\|^2}{2} dv_g}{\rho_2^\Lambda}$$

for any  $0 < \rho_1 \leq \rho_2$ . □




Go back

Full Screen

Close

Quit



**Corollary 5.5.** *Let  $M, K_r$  and  $\Lambda$  be as in Theorem 5.4. Assume that  $u: (M, g) \rightarrow (N, h)$  is a biharmonic isometric immersion and  $h(\tau(u), \tilde{\nabla}_{\frac{\partial}{\partial r}} du(\frac{\partial}{\partial r})) \geq 0$ . If*

$$\int_{B(R)} \frac{\|\tau(u)\|^2}{2} dv_g = o(R^\Lambda),$$

*then  $u$  is harmonic.*

We say the bienergy  $E_2(u)$  of  $u$  is slowly divergent if there exists a positive function  $\psi(r)$  with  $\int_{R_0}^\infty \frac{dr}{r\psi(r)} = +\infty$  ( $R_0 > 0$ ) such that

$$(40) \quad \lim_{R \rightarrow \infty} \int_{B(R)} \frac{\|\tau(u)\|^2}{2\psi(r(x))} dv_g < \infty.$$

**Theorem 5.6.** *Let  $u: (M, g) \rightarrow (N, h)$  be a biharmonic isometric immersion. Assume that there is a constant  $\sigma > 0$  such that*

$$\frac{m-1}{2} \lambda_{\min} + 1 - 2 \max\{2, \lambda_{\max}\} \geq \sigma.$$

*If  $E_2(u)$  is slowly divergent and  $h(\tau(u), \tilde{\nabla}_{\frac{\partial}{\partial r}} du(\frac{\partial}{\partial r})) \geq 0$ , then  $u$  is harmonic, i.e.,  $\tau(u) = 0$ .*

*Proof.* From the proof of Theorem 5.1, we have

$$(41) \quad \sigma \int_{B(r)} \frac{\|\tau(u)\|^2}{2} dv_g \leq r \int_{\partial B(r)} \frac{\|\tau(u)\|^2}{2} ds_g.$$

Now suppose that  $u$  is not harmonic, so there exists  $R_0 > 0$  such that for  $R \geq R_0$ ,

$$(42) \quad \sigma \int_{B(R)} \frac{\|\tau(u)\|^2}{2} dv_g \geq c_1,$$




Go back

Full Screen

Close

Quit



where  $c_1$  is a positive constant. From (41) and (42), we have

$$(43) \quad c_1\sigma \leq R \int_{\partial B(R)} \frac{\|\tau(u)\|^2}{2} ds_g.$$

for  $R \geq R_0$  and

$$\begin{aligned} \lim_{R \rightarrow \infty} \int_{B(R)} \frac{\|\tau(u)\|^2}{2\psi(r(x))} dv_g &= \int_0^\infty \frac{dR}{\psi(R)} \int_{\partial B(R)} \frac{\|\tau(u)\|^2}{2} ds_g \\ &\geq \int_{R_0}^\infty \frac{dR}{\psi(R)} \int_{\partial B(R)} \frac{\|\tau(u)\|^2}{2} ds_g \\ &\geq c_1\sigma \int_{R_0}^\infty \frac{dR}{R\psi(R)} = \infty, \end{aligned}$$

which contradicts (40), therefore,  $u$  is harmonic. □

From the proof of Theorem 5.6, we immediately get the following theorem.

**Theorem 5.7.** *Let  $M, K_r$  and  $\Lambda$  be as in Theorem 5.4. If  $u: (M, g) \rightarrow (N, h)$  is a biharmonic isometric immersion, the bienergy  $E_2(u)$  is slowly divergent and  $h(\tau(u), \tilde{\nabla}_{\frac{\partial}{\partial r}} du(\frac{\partial}{\partial r})) \geq 0$ , then  $u$  is harmonic.*

**Acknowledgment.** The authors would like to thank the referee whose valuable suggestions make this paper more perfect. Yingbo Han was supported by NSFC No.11201400, No.10971029, No.11026062, Project of Henan Provincial Department of Education No.2011A110015 and Talent youth teacher fund of Xinyang Normal University.



Go back

Full Screen

Close

Quit



1. Ara M., *Geometry of  $F$ -harmonic maps*, Kodai Math. J. **22**(1999), 243–263.
2. Baird P., *Stress-energy tensors and the Linchernerowicz Laplacian*, J. Geo. and Phys. **58** (2008), 1329–1342.
3. Caddeo R., Montaldo S. and Piu P., *On biharmonic maps*, Contemp. Math. **288** (2001), 286–290.
4. Dong Y. X. and Wei S. S., *On vanishing theorems for vector bundle valued  $p$ -forms and their applications*, Comm. Math. Phys. **304** (2011), 329–368.
5. Dong Y. X., Lin H. Z. and Yang G. L., *Liouville theorems for  $F$ -harmonic maps and their applications*, arXiv:1111.1882v1 [math.DG] 8 Nov 2011.
6. Hornung P. and Moser R., *Intrinsically  $p$ -biharmonic maps*, preprint (Opus: University of Bath online publication store).
7. Eells J., and Lemaire L., *Selected topics in harmonic maps*, CBMS 50, Amer. Math. Soc. 1983.
8. Greene R. E. and Wu H., *Function theory on manifolds which possess pole*, In: Lecture Notes in Mathematics, V. 699, Springer-verlag, Berlin, Heidelberg, New York 1979.
9. Jiang G. Y., *2-harmonic maps and their first and second variational formulas*, Chinese Ann. Math. **7A** (1986), 388–402; the English translation, Note di Matematica, **28** (2009), 209–232.
10. ———, *The conservation law for 2-harmonic maps between Riemannian manifolds*, Acta Math. Sin. **30** (1987), 220–225.
11. Kassi M., *A Liouville theorems for  $F$ -harmonic maps with finite  $F$ -energy*, Electronic Journal Diff. Equa. **15** (2006), 1–9.
12. Liu J. C., *Liouville theorems of stable  $F$ -harmonic maps for compact convex hypersurfaces*, Hiroshima Math. J. **36** (2006), 221–234.
13. Loubeau E. and Oniciuc C., *The index of biharmonic maps in spheres*, Compositio Math. **141** (2005), 729–745.
14. Loubeau E., Montaldo S. and Oniciuc C., *The stress-energy tensor for biharmonic maps*, Math. Z. **259** (2008), 503–524.
15. Nakauchi N., Urakawa H. and Gudmundsson S., *Biharmonic maps into a Riemannian manifold of non-positive curvature*, arXiv:1201.6457v4.
16. Oniciuc C., *Biharmonic maps between Riemannian manifolds*, Analele stiintifice ale Univer. "Al. I. Cusa", din Iasi (2002), 237–248.
17. Ouakkas S., Nasri R., and Djaa M., *On the  $f$ -harmonic and  $f$ -biharmonic maps*, JP J. Geom. Topol. **10**(1) (2010), 11–27.




Go back

Full Screen

Close

Quit



18. Wang C., *Remarks on biharmonic maps into spheres*, Calc. Var. Partial Differential Equations, **21** (2004), 221–242.

19. ———, *Biharmonic maps from  $R^4$  into Riemannian manifold*. Math. Z. **247** (2004), 65–87.

Yingbo Han, College of Mathematics and Information Science, Xinyang Normal University, Xinyang, 464000, Henan, P. R. China, *e-mail*: [hyb@xynu.edu.cn](mailto:hyb@xynu.edu.cn)

Shuxiang Feng, College of Mathematics and Information Science, Xinyang Normal University, Xinyang, 464000, Henan, P. R. China, *e-mail*: [shuxiangfeng78@163.com](mailto:shuxiangfeng78@163.com)



*Go back*

*Full Screen*

*Close*

*Quit*