



NORMAL GENERATION OF UNITARY GROUPS OF CUNTZ ALGEBRAS BY INVOLUTIONS

A. AL-RAWASHDEH

ABSTRACT. In purely infinite factors, P. de la Harpe proved that a normal subgroup of the unitary group which contains a non-trivial self-adjoint unitary contains all self-adjoint unitaries of the factor. Also he proved the same result in finite continuous factors. In a previous work the author proved a similar result in some types of unital AF-algebras. In this paper we extend the result of de la Harpe, concerning the purely infinite factors to a main example of purely infinite C^* -algebras called the Cuntz algebras \mathcal{O}_n ($2 \leq n \leq \infty$) and we prove that $\mathcal{U}(\mathcal{O}_n)$ is normally generated by some non-trivial involution. In particular, in the Cuntz algebra \mathcal{O}_∞ we prove that $\mathcal{U}(\mathcal{O}_\infty)$ is normally generated by self-adjoint unitary of odd type.

1. INTRODUCTION

Let \mathcal{A} be any unital C^* -algebra. The group of unitaries and the set of projections of \mathcal{A} are denoted by $\mathcal{U}(\mathcal{A})$, $\mathcal{P}(\mathcal{A})$ respectively. The involutions of \mathcal{A} are the set of self-adjoint unitaries ($*$ -symmetries). In several types of C^* -algebras, we have that the involutions generate all the unitaries. In the case of von Neumann factors, M. Broise in [3]; proved the following main theorem.

Theorem 1.1. [3, Theorem 1] *If \mathcal{B} is a factor of type II_1 or III , then the set of involutions generates $\mathcal{U}(\mathcal{B})$.*

Received June 11, 2006; revised January 13, 2008.

2000 *Mathematics Subject Classification.* Primary 46L05; 46L80.

Key words and phrases. Cuntz algebras; involutions; K -Theory.

Also, in the case of simple, purely infinite C^* -algebras, M. Leen proved the following result.

Theorem 1.2. [10, Theorem 3.8] *If A is a simple, unital purely infinite C^* -algebra, then the set of $*$ -symmetries of A forms a set of generators for $\mathcal{U}_0(A)$, (where $\mathcal{U}_0(A)$ denotes the identity component of the unitary group of A).*

The Cuntz algebras are interesting examples of simple, unital purely infinite C^* -algebras, which was introduced by Cuntz in [5] this C^* -algebra is generated by isometries that have orthogonal ranges (for more information see [6, V. 4]). As shown in [5] the unitary group of the Cuntz algebras are connected. Now let us recall these definitions.

Definition 1.3. The Cuntz algebra \mathcal{O}_n , where $2 \leq n$, is the universal C^* -algebra which is generated by isometries s_1, s_2, \dots, s_n , such that

$$(1) \quad \sum_{i=1}^n s_i s_i^* = 1$$

with $s_i^* s_j = 0$, when $i \neq j$. The Cuntz algebra \mathcal{O}_∞ is generated by infinite number of such isometries.

Remark 1.4. [6, V. 4] Recall that a universal C^* -algebra \mathcal{O}_n means, whenever t_1, t_2, \dots, t_n form another set of isometries satisfying (1), then there is a unique $*$ -homomorphism ρ of \mathcal{O}_n onto $C^*({t_1, t_2, \dots, t_n})$ such that $\rho(s_i) = t_i$, for all $1 \leq i \leq n$.

In this paper, the projection $s_i s_i^*$ is denoted by p_i , and these projections are called the standard projections of the Cuntz algebras. The corresponding involution $1 - 2p_i$ is denoted by u_i .

Let us recall the following main results concerning the Cuntz algebras.

Theorem 1.5. [5] *The Cuntz algebras \mathcal{O}_n ($2 \leq n \leq \infty$) are simple unital purely infinite C^* -algebras.*

Using the fact that $K_1(\mathcal{O}_n) \cong 0$ (see [4, 3.8]) and $K_1(A) \simeq \mathcal{U}(A)/\mathcal{U}(A)_0$ (see [4, p. 188], M. Leen's result (Theorem 1.2) shows that the set of $*$ -symmetries of \mathcal{O}_n ($2 \leq n \leq \infty$) generates the unitary group $\mathcal{U}(\mathcal{O}_n)$.

Definition 1.6. A group G is normally generated by an element x if the only normal subgroup of G containing x is G itself.

If $u = 1 - 2p$ is an involution in a factor \mathcal{B} , then P. de la Harpe defined the notion of the type of u to be the pair (x, y) , where $x = D(1 - p)$ and $y = D(p)$, as D denotes a normalized dimension function on \mathcal{B} , see [7]. He proved that any normal subgroup \mathcal{N} of $\mathcal{U}(\mathcal{B})$, which is not contained in the circle \mathbb{S}^1 , contains a non-trivial involution, and then contains all the involutions of \mathcal{B} (see [8, Proposition 2]). Afterwards, P. de la Harpe used Broise's result (Theorem 1.1), and he proved the following theorem.

Theorem 1.7. [8] *If \mathcal{B} is a factor of type II_1 or III and \mathcal{N} is any normal subgroup of $\mathcal{U}(\mathcal{B})$, which is not contained in the circle \mathbb{S}^1 , then $\mathcal{N} = \mathcal{U}(\mathcal{B})$.*

If v is an involution of \mathcal{O}_n ($2 \leq n \leq \infty$), then as introduced in [1], we define the type of v to be the element $[p]$ in $K_0(\mathcal{O}_n)$, where $v = 1 - 2p$. Since the $K_0(\mathcal{O}_n)$ is a cyclic group, the type of v is an integer. In Section 2, we show that a normal subgroup \mathcal{N} of $\mathcal{U}(\mathcal{O}_n)$, $n < \infty$ contains all the involutions if

1. \mathcal{N} contains an involution of the type 1 (i.e. [1]), or
2. \mathcal{N} contains a non-trivial involution and $n - 1$ is a prime number, or
3. \mathcal{N} contains a non-trivial involution such that its type and $n - 1$ are relatively prime. Then using M. Leen's result in Theorem 1.2, we prove that $\mathcal{U}(\mathcal{O}_n)$ is normally generated by a non-trivial involution.

In Section 3, we show that if \mathcal{N} contains an involution of odd type, then \mathcal{N} contains all the involutions of \mathcal{O}_∞ . Consequently, we use M. Leen's result in order to prove that $\mathcal{U}(\mathcal{O}_n)$ is normally generated by an involution of odd type.

Now, let us recall main results concerning purely infinite C^* -algebras, that might be used throughout this paper.

Proposition 1.8. [4, 1.5] *In any C^* -algebra A , the following hold:*

- (i) *If p, q are infinite projections and $pq = 0$, then $p + q$ is an infinite projection.*
- (ii) *If p is an infinite projection, and $p' \sim p$, then p' is an infinite projection.*
- (iii) *If p and q are infinite projections, then there exists an infinite projection p' such that $p \sim p'$ and $p' < q$, moreover $q - p'$ is an infinite projection.*

Theorem 1.9. [2, 6.11.9] *Two infinite projections in a simple unital C^* -algebra are equivalent if and only if they have the same K_0 -class. Two non-trivial projections with the same K_0 -class in a purely infinite C^* -algebra are unitarily equivalent.*

2. THE $\mathcal{O}_n(2 \leq n < \infty)$ CASE

We prove the following result which is valid for the Cuntz algebras $\mathcal{O}_n(2 \leq n \leq \infty)$. The proof is similar to [1, Lemma 2.2], in the case of the UHF-algebras. For completeness we have.

Lemma 2.1. *Let u and v be two involutions of $\mathcal{O}_n(2 \leq n \leq \infty)$. Then u is conjugate to v if and only if they have the same type.*

Proof. If u and v are conjugate involutions of $\mathcal{O}_n(2 \leq n \leq \infty)$, then as in [1, Lemma 2.2], there exists a unitary w in $\mathcal{U}(\mathcal{O}_n)$ such that $u = wvw^*$. But $u = 1 - 2e$ and $v = 1 - 2f$ for some projections e, f in A , so $u = w(1 - 2f)w^* = 1 - 2wf w^*$, therefore $e = wf w^*$ and by Theorem 1.9, $[e] = [f]$.

Conversely, assume that the involutions u and v have the same type. Then $u = 1 - 2p$ and $v = 1 - 2q$, for some $p, q \in \mathcal{P}(\mathcal{O}_n)$ with $[p] = [q]$ in $K_0(\mathcal{O}_n)$ group. Then by Theorem 1.9, the projections p and q are unitarily equivalent, and therefore $u = wvw^*$ for some $w \in \mathcal{U}(\mathcal{O}_n)$. \square

Proposition 2.2. *In $\mathcal{O}_n(2 \leq n \leq \infty)$, the involution $u_i(i = 1, \dots, n)$ has type 1.*

Proof. As $u_i = 1 - 2p_i$, the type of u_i is $[p_i]$. By definition $p_i = s_i s_i^*$ and $s_i^* s_i = 1$; therefore by Theorem 1.9, we have $[p_i] = [1]$. \square

The following result is based on [4, 3.7, 3.8]; that is $K_0(\mathcal{O}_n) \simeq \mathbb{Z}_{n-1}$.

Proposition 2.3. *If $0 \leq k \leq n - 2$; $n < \infty$, then there exists an involution in \mathcal{O}_n of type k (in fact, of type $k[1]$).*

Proof. Let p_1, p_2, \dots, p_n be the standard projections of \mathcal{O}_n , and $v_k = 1 - 2(p_1 + p_2 + \dots + p_k)$; for $0 \leq k \leq n - 2$. Then v_k is an involution in \mathcal{O}_n of type equal to k . \square

Lemma 2.4. *If \mathcal{N} is a normal subgroup of $\mathcal{U}(\mathcal{O}_n)(n < \infty)$, which contains an involution of the type 1([1]), then \mathcal{N} contains an involution of any given type.*

Proof. As \mathcal{N} is a normal subgroup of $\mathcal{U}(\mathcal{O}_n)$, and it contains an involution of the type 1, then by Lemma 2.1, \mathcal{N} contains $u_i(i = 1, \dots, n)$. Then $u_1 u_2 = (1 - 2p_1)(1 - 2p_2) = 1 - 2(p_1 + p_2)$, which is an involution of type 2, contained in \mathcal{N} . Also $u_1 u_2 u_3$ is an involution in \mathcal{N} of type 3. Keep going we have $u_1 u_2 \dots u_k = 1 - 2(p_1 + p_2 + \dots + p_k)$ is an involution in \mathcal{N} of type $k(1 \leq k \leq n - 2)$, hence \mathcal{N} contains an involution of any given type, which proves the required. \square

Lemma 2.5. *Let \mathcal{N} be a normal subgroup of $\mathcal{U}(\mathcal{O}_n)$, and suppose that $n - 1$ is a prime number. If \mathcal{N} contains a non-trivial involution of \mathcal{O}_n , then \mathcal{N} contains the involution u_1 .*



Proof. Suppose that $v \in \mathcal{N}$ such that $v = 1 - 2p$ and v is of type k , i.e. $[p] = k$. If $k = n - 1$, then $[p] = 0 \in K_0(\mathcal{O}_n)$, by Proposition 1.8(ii) we must have $p = 0$ and then $v = 1$ which gives a contradiction as v is non-trivial. Therefore we consider $1 \leq k \leq n - 2$. We may assume that $p < 1$, since if $p = 1$, then $v = -1$ which is an involution of type one, and this ends the proof. As $n - 1$ is a prime, there exist integers s and t such that $sk + t(n - 1) = 1$, then $sk = 1$ in \mathbb{Z}_{n-1} . By Proposition 1.8(iii), we can find mutually orthogonal projections q_1, q_2, \dots, q_s , with $[q_i] = [p]$, $i = 1, \dots, s$. Let $v_i = 1 - 2q_i$, $i = 1, \dots, s$. Then for every i , v_i there is an involution of the type k , which belongs to \mathcal{N} as it is conjugate to v . Therefore

$$v_1 v_2 \dots v_s = 1 - 2(q_1 + q_2 + \dots + q_s)$$

is an involution in \mathcal{N} , and the type of $v_1 v_2 \dots v_s$ is $sk = 1 \in \mathbb{Z}_{n-1}$. \square

By imitating the same proof in the previous result, we can rewrite Lemma 2.5 as follows:

Lemma 2.6. *Let \mathcal{N} be a normal subgroup of $\mathcal{U}(\mathcal{O}_n)$. If \mathcal{N} contains an involution of type k such that k and $n - 1$ are relatively primes, then \mathcal{N} contains an involution of type 1.*

Therefore, we have the following theorem.

Theorem 2.1. *A non-trivial involution u normally generates the group $\mathcal{U}(\mathcal{O}_n)$ if either*

- (1) $n - 1$ is a prime number, or
- (2) the type of u is relatively prime to $n - 1$.

Proof. If \mathcal{N} is a normal subgroup of $\mathcal{U}(\mathcal{O}_n)$ that contains a non-trivial involution with hypothesis of either (1) or (2), then by either Lemma 2.5 or Lemma 2.6, \mathcal{N} contains an involution of type 1, therefore by Lemma 2.4, \mathcal{N} contains an involution of any given type, then by Lemma 2.1 it contains all involutions, hence by Leen's result $\mathcal{N} = \mathcal{U}(\mathcal{O}_n)$. \square



3. THE \mathcal{O}_∞ CASE

In this section, we discuss the case of the Cuntz algebra \mathcal{O}_∞ . We may ask, if a normal subgroup of $\mathcal{U}(\mathcal{O}_\infty)$ contains a non-trivial involution u_0 , then does it contain all the involutions of \mathcal{O}_∞ ? Hence by using Leen's result in Theorem 1.2, \mathcal{O}_∞ is normally generated by a non-trivial involution u_0 . We give a positive answer to the question under some conditions on the non-trivial involution u_0 .

Recall that the Cuntz algebra \mathcal{O}_∞ is the universal unital C^* -algebra generated by an infinite sequence of isometries s_1, s_2, s_3, \dots with mutually orthogonal projections $p_j = s_j s_j^*$. The involution $1 - 2p_j$ is denoted by u_j ($1 \leq j \leq \infty$).

Now let us recall the following main results concerning \mathcal{O}_∞ .

Theorem 3.1. [4, 3.11]

- (i) $K_0(\mathcal{O}_\infty) \cong \mathbb{Z}$.
- (ii) $K_1(\mathcal{O}_\infty) \cong 0$.

Theorem 3.2. [4, 3.12] *In \mathcal{O}_∞ , every projection is equivalent to a projection either of the form $\sum_{i=1}^k s_i s_i^*$ ($1 \leq k < \infty$) or $1 - \sum_{i=1}^k s_i s_i^*$ ($1 \leq k < \infty$).*

In \mathcal{O}_∞ , the type of an involution v is $n[1]$, for some $n \in \mathbb{Z}$, and we write that v has the type $n \in \mathbb{Z}$. Recall that Lemma 2.1 is also valid for \mathcal{O}_∞ .

Now we start by proving the following lemma, which is similar to Lemma 2.4 in the case of \mathcal{O}_n , where n is a finite number.

Lemma 3.3. *If \mathcal{N} is a normal subgroup of $\mathcal{U}(\mathcal{O}_\infty)$, which contains an involution of the type 1, then \mathcal{N} contains an involution of any given type.*

Proof. As \mathcal{N} contains an involution of type 1, and \mathcal{N} is a normal subgroup of $\mathcal{U}(\mathcal{O}_\infty)$, we have that \mathcal{N} contains all the involutions u_i $i = 1, 2, \dots$. Then $u_1 u_2$ is an involution in \mathcal{N} of type 2 indeed,

if $k \in \mathbb{Z}^+$, then $u_1 u_2 \dots u_k = 1 - 2(p_1 + p_2 + \dots + p_k)$ is an involution in \mathcal{N} of type k . Also, \mathcal{N} contains an involution of type 0, as $1 \in \mathcal{N}$.

Now it is enough to prove that \mathcal{N} contains an involution of any negative type. Recall that if p is a projection of \mathcal{O}_∞ , then by Theorem 3.2, either p is equivalent to $\sum_{i=1}^k s_i s_i^*$, hence $[p] = k[1]$ or p is equivalent to $1 - \sum_{i=1}^k s_i s_i^*$, hence $[p] = (1 - k)[1]$, for some $k \in \mathbb{Z}^+$. As \mathcal{N} contains involutions of type 1, then the involution -1 belongs to \mathcal{N} . Hence for each $k \in \mathbb{Z}^+$, $-u_1 u_2 \dots u_k \in \mathcal{N}$, and

$$\begin{aligned} -u_1 u_2 \dots u_k &= -(1 - 2(p_1 + p_2 + \dots + p_k)) \\ &= -1 + 2(p_1 + p_2 + \dots + p_k) \\ &= 1 - 2(1 - (p_1 + p_2 + \dots + p_k)), \end{aligned}$$

therefore, $-u_1 u_2 \dots u_k$ is an involution of type $1 - k$ and the lemma has been checked. \square

Therefore, we have the following main result

Theorem 3.4. *Any involution of type 1 normally generates the group $\mathcal{U}(\mathcal{O}_\infty)$.*

Proof. Suppose that \mathcal{N} is a normal subgroup of $\mathcal{U}(\mathcal{O}_\infty)$ that contains an involution of the type 1. By using Lemma 3.3, we have that \mathcal{N} contains an involution of any given type, therefore by Lemma 2.1, \mathcal{N} contains all the involutions, hence by Leen's result in Theorem 1.2, $\mathcal{N} = \mathcal{U}(\mathcal{O}_\infty)$. \square

Let us now prove our main result.

Theorem 3.5. *Any involution of odd type normally generates the group $\mathcal{U}(\mathcal{O}_\infty)$.*

Proof. Case 1: Suppose that \mathcal{N} contains an involution of type $2k + 1$, for some positive integer k . By normality of \mathcal{N} , we may assume that $v = 1 - 2 \sum_{i=1}^{2k+1} p_i \in \mathcal{N}$, also $u = 1 - 2 \sum_{i=2}^{2k+2} p_i \in \mathcal{N}$.

Therefore, we have that

$$vu = (1 - 2 \sum_{i=1}^{2k+1} p_i)(1 - 2 \sum_{i=2}^{2k+2} p_i) = 1 - 2(p_1 + p_{2k+2}),$$

which is an involution in \mathcal{N} of type 2, hence \mathcal{N} contains all involutions of the type 2. Then

$$(1 - 2(p_1 + p_2))(1 - 2(p_3 + p_4)) \dots (1 - 2(p_{2k-1} + p_{2k})) = 1 - 2 \sum_{i=1}^{2k} p_i \in \mathcal{N}.$$

Therefore \mathcal{N} contains the involution

$$(1 - 2 \sum_{i=1}^{2k+1} p_i)(1 - 2 \sum_{i=1}^{2k} p_i) = 1 - 2p_{2k+1},$$

which is of the type 1, hence by Theorem 3.4, we have the desired.

Case 2: Suppose that \mathcal{N} contains an involution v of the type $-k$, where $k \in \mathbb{Z}^+$, which is odd. Then by normality of \mathcal{N} and Lemma 2.1, the involution $w_1 = 1 - 2(1 - (p_1 + p_2 + \dots + p_{k+1}))$ belongs to \mathcal{N} , as its type is $-k$. In fact, $w_1 = -u_1 u_2 \dots u_k u_{k+1}$. Similarly, the involution $w_2 = 1 - 2(1 - (p_2 + p_3 + \dots + p_{k+2}))$ belongs to \mathcal{N} and $w_2 = -u_2 u_3 \dots u_{k+2}$. Therefore, the involution $w_1 w_2 = u_1 u_{k+2} \in \mathcal{N}$, hence \mathcal{N} contains all involutions of type 2, by using Lemma 2.1. As $k+1$ is an even integer, we get $w_3 = (u_1 u_2)(u_3 u_4) \dots (u_k u_{k+1}) \in \mathcal{N}$. Therefore we have that $w_1 w_3 = -1 \in \mathcal{N}$, which is an involution of type 1, hence by Theorem 3.4, the proof is completed. \square

Finally, we conclude by noting that similar arguments show that a normal subgroup of $\mathcal{U}(\mathcal{O}_n)$ which contains a non-trivial involution (of any type) necessarily contains all the involutions of even type.

Home Page

Title Page

Contents



Page 10 of 10

Go Back

Full Screen

Close

Quit

1. Al-Rawashdeh A., *On Normal Subgroups of Unitary Groups of Some Unital AF-Algebras*, Sarajevo Journal of Mathematics, **3**(16), (2007), 233–240.
2. Blackadar B., *K-Theory for Operator Algebras*, Second Edition, MSRI Publications, **5**, Cambridge University Press, Cambridge 1998.
3. Broise M., *Commutateurs Dans le Groupe Unitaire d'un Facteur*, J. Math. Pures et appl., **46** (1967), 299–312.
4. Cuntz J., *K-Theory for Certain C^* -Algebras*, Ann. of Math., **113** (1981), 181–197.
5. ———, *Simple C^* -Algebras Generated by Isometries*, Comm. Math. Phys., **57** (1977), 173–185.
6. Davidson K. R., *C^* -Algebras by Example*, Fields Institute Monographs, **6**, Amer. Math. Soc., Providences, RI 1996.
7. Dixmier J., *Les algèbres d'opérateurs dans l'espace hilbertien (algèbres de von Neumann)*, Sécond Édition, Gauthier-Villars 1969.
8. de la Harpe P., *Simplicity of the Projective Unitary Groups Defined by Simple Factors*, Comment. Math. Helv., **54** (1979), 334–345.
9. de la Harpe P. and Jones V. F. R., *An Introduction to C^* -Algebras*, Université de Genève, 1995.
10. Leen M., *Factorization in the Invertible Group of a C^* -Algebra*, Canad. J. Math., **49**(6) (1997), 1188–1205.

A. Al-Rawashdeh, Department of Mathematics and Statistics, Jordan University of Science and Technology, Irbid 22110, Jordan, e-mail: rahmed72@just.edu.jo