

**SOME PROPERTIES FOR SUBCLASSES OF MEROMORPHIC
MULTIVALENT FUNCTIONS ASSOCIATED WITH NEW
OPERATOR**

M.K. AOUF, A.O. MOSTAFA, H.M. ZAYED

ABSTRACT. The purpose of this paper is to introduce two subclasses of meromorphic multivalent functions by using new operator and investigate various properties for these subclasses.

2010 *Mathematics Subject Classification:* 30C45, 30C50.

Keywords: Meromorphic multivalent functions, Hadamard product (or convolution), linear operator.

1. INTRODUCTION

Let Σ_p be the class of meromorphic p -valent functions of the form:

$$f(z) = \frac{1}{z^p} + \sum_{n=1}^{\infty} a_{n-p} z^{n-p} \quad (p \in \mathbb{N} = \{1, 2, \dots\}), \quad (1)$$

which are analytic in the punctured unit disc $\mathbb{U}^* = \{z : z \in \mathbb{C} \text{ and } 0 < |z| < 1\} = \mathbb{U} \setminus \{0\}$.

Let $\mathcal{P}_k(\rho)$ be the class of functions $p(z)$ analytic in \mathbb{U} satisfying the properties $p(0) = 1$ and

$$\int_0^{2\pi} \left| \frac{\operatorname{Re} p(z) - \rho}{1 - \rho} \right| d\theta \leq k\pi, \quad (2)$$

where $k \geq 2$ and $0 \leq \rho < 1$. This class was introduced by Padmanabhan and Parvatham [10]. For $\rho = 0$, the class $\mathcal{P}_k(0) = \mathcal{P}_k$ introduced by Pinchuk [12]. Also, $\mathcal{P}_2(\rho) = \mathcal{P}(\rho)$, where $\mathcal{P}(\rho)$ is the class of functions with positive real part greater

than ρ and $\mathcal{P}_2(0) = \mathcal{P}$, is the class of functions with positive real part. From (2), we have $p(z) \in \mathcal{P}_k(\rho)$ if and only if there exist $p_1, p_2 \in \mathcal{P}(\rho)$ such that

$$p(z) = \left(\frac{k}{4} + \frac{1}{2}\right) p_1(z) - \left(\frac{k}{4} - \frac{1}{2}\right) p_2(z) \quad (z \in \mathbb{U}). \quad (3)$$

It is known that the class $\mathcal{P}_k(\rho)$ is a convex set (see [7]).

The Hadamard product (or convolution) $f * g$ of $f(z)$ given by (1) and $g(z)$ given by

$$g(z) = \frac{1}{z^p} + \sum_{n=1}^{\infty} b_{n-p} z^{n-p}, \quad (4)$$

is defined by

$$(f * g)(z) = \frac{1}{z^p} + \sum_{n=1}^{\infty} a_{n-p} b_{n-p} z^{n-p} = (g * f)(z). \quad (5)$$

Define the functions

$$f_{\alpha}(z) = \frac{1}{z^p} + \sum_{n=1}^{\infty} \left(\frac{n+\lambda}{\lambda}\right)^{\alpha} z^{n-p} \quad (\alpha \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}; \lambda > 0),$$

and

$$\tilde{\varphi}(a, b; z) = \frac{1}{z^p} F(1, a; b; z) = \frac{1}{z^p} + \sum_{n=1}^{\infty} \left| \frac{(a)_n}{(b)_n} \right| z^{n-p},$$

where $a \in \mathbb{C}^* = \mathbb{C} \setminus \{0\}$, $b \in \mathbb{R} \setminus \mathbb{Z}_0^-$, $\mathbb{Z}_0^- = \{0, -1, -2, \dots\}$, $F(a, b; c; z)$ is the (Gaussian) hypergeometric function defined by

$$F(a, b; c; z) = \sum_{n=0}^{\infty} \left| \frac{(a)_n (b)_n}{(c)_n (1)_n} \right| z^n \quad (c \in \mathbb{R} \setminus \mathbb{Z}_0^-),$$

and

$$(\eta)_n = \begin{cases} 1 & \text{if } n = 0, \\ \eta(\eta+1)(\eta+2)\dots(\eta+n-1) & \text{if } n \in \mathbb{N}. \end{cases}$$

Also, let the associated function $f_{\alpha, \mu}^*(z)$ be defined by the Hadamard product (or convolution):

$$f_{\alpha}(z) * f_{p, \alpha, \mu}^*(z) = \frac{1}{z^p (1-z)^{\mu}} \quad (\mu > 0; z \in \mathbb{U}^*).$$

Then, we have

$$\begin{aligned} \mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z) &= f_{p,\alpha,\mu}^*(z) * \tilde{\varphi}(a, b; z) * f(z) \\ &= \frac{1}{z^p} + \sum_{n=1}^{\infty} \left| \frac{(a)_n}{(b)_n} \right| \left(\frac{\lambda}{n+\lambda} \right)^\alpha \frac{(\mu)_n}{(1)_n} a_{n-p} z^{n-p}. \end{aligned} \quad (6)$$

It is easily verified from (6) that

$$z \left(\mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z) \right)' = \mu \mathcal{I}_{a,b,\mu+1}^{p,\alpha,\lambda} f(z) - (\mu + p) \mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z), \quad (7)$$

and

$$z \left(\mathcal{I}_{a,b,\mu}^{p,\alpha+1,\lambda} f(z) \right)' = \lambda \mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z) - (\lambda + p) \mathcal{I}_{a,b,\mu}^{p,\alpha+1,\lambda} f(z). \quad (8)$$

We note that:

- (i) $\mathcal{I}_{1,1,\mu}^{p,\alpha,\ell} f(z) = I_{p,\mu}^\alpha(\ell) f(z)$ (see El-Ashwah and Aouf [4, with $\lambda = 1$]);
- (ii) $\mathcal{I}_{1,1,\mu}^{1,\alpha,\lambda} f(z) = \mathcal{I}_{\lambda,\mu}^\alpha f(z)$ (see Cho et al. [3] and Piejko and Sokol [11]) (see also Aouf et al. [2]).

Next, by using the operator $\mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda}$, we introduce two subclasses of meromorphic multivalent functions of Σ_p as follows:

Definition 1. A function $f(z) \in \Sigma_p$ is said to be in the class $\mathcal{M}_{a,b,\mu}^{p,\alpha,\lambda}(\beta, \gamma, \rho, k)$ if it satisfies the condition:

$$\begin{aligned} \left[(1 - \gamma) \left(z^p \mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z) \right)^\beta + \gamma \left(\frac{\mathcal{I}_{a,b,\mu+1}^{p,\alpha,\lambda} f(z)}{\mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z)} \right) \left(z^p \mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z) \right)^\beta \right] \in \mathcal{P}_k(\rho), \\ (k \geq 2; a \in \mathbb{C}^*; b \in \mathbb{R} \setminus \mathbb{Z}_0^-; \alpha \in \mathbb{N}_0; \beta, \gamma, \mu, \lambda > 0; 0 \leq \rho < 1). \end{aligned} \quad (9)$$

We note that:

$$\mathcal{M}_{1,1,\mu}^{1,\alpha,\lambda}(\beta, \gamma, \rho, k) = \mathcal{M}_{\lambda,\mu}^\alpha(\beta, \gamma, \rho, k) \text{ (see Aouf et al. [2]).}$$

Also, we note that:

$$\begin{aligned} \mathcal{M}_{1,1,\mu}^{p,\alpha,\ell}(\beta, \gamma, \rho, k) = \mathcal{M}_\mu^{p,\alpha,\ell}(\beta, \gamma, \rho, k) = \{ f(z) \in \Sigma_p : \\ \left[(1 - \gamma) \left(z^p I_{p,\mu}^\alpha(\ell) f(z) \right)^\beta + \gamma \left(\frac{I_{p,\mu+1}^\alpha(\ell) f(z)}{I_{p,\mu}^\alpha(\ell) f(z)} \right) \left(z^p I_{p,\mu}^\alpha(\ell) f(z) \right)^\beta \right] \in \mathcal{P}_k(\rho) \}. \end{aligned}$$

Definition 2. A function $f(z) \in \Sigma_p$ is said to be in the class $\mathcal{N}_{a,b,\mu}^{p,\alpha,\lambda}(\beta, \gamma, \rho, k)$ if it satisfies the condition:

$$\left[(1 - \gamma) \left(z^p \mathcal{I}_{a,b,\mu}^{p,\alpha+1,\lambda} f(z) \right)^\beta + \gamma \left(\frac{\mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z)}{\mathcal{I}_{a,b,\mu}^{p,\alpha+1,\lambda} f(z)} \right) \left(z^p \mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z) \right)^\beta \right] \in \mathcal{P}_k(\rho),$$

$$(k \geq 2; a \in \mathbb{C}^*; b \in \mathbb{R} \setminus \mathbb{Z}_0^-; \alpha \in \mathbb{N}_0; \beta, \gamma, \mu, \lambda > 0; 0 \leq \rho < 1). \quad (10)$$

We note that:

$$\mathcal{N}_{1,1,\mu}^{1,\alpha,\lambda}(\beta, \gamma, \rho, k) = \mathcal{N}_{\lambda,\mu}^\alpha(\beta, \gamma, \rho, k) \text{ (see Aouf et al. [2]).}$$

Also, we note that:

$$\mathcal{N}_{1,1,\mu}^{p,\alpha,\ell}(\beta, \gamma, \rho, k) = \mathcal{N}_\mu^{p,\alpha,\ell}(\beta, \gamma, \rho, k) = \left\{ f(z) \in \Sigma_p : \left[(1 - \gamma) \left(z^p I_{p,\mu}^{\alpha+1}(\ell) f(z) \right)^\beta + \gamma \left(\frac{I_{p,\mu}^{\alpha,\ell} f(z)}{I_{p,\mu}^{\alpha+1}(\ell) f(z)} \right) \left(z^p I_{p,\mu}^{\alpha+1}(\ell) f(z) \right)^\beta \right] \in \mathcal{P}_k(\rho) \right\}.$$

2. MAIN RESULTS

Unless otherwise mentioned, we assume throughout this paper that $k \geq 2$, $a \in \mathbb{C}^*$, $b \in \mathbb{R} \setminus \mathbb{Z}_0^-$, $\alpha \in \mathbb{N}_0$, $\beta, \gamma, \mu, \lambda > 0$ and $0 \leq \rho < 1$.

To establish our results, we need the following lemma due to Miller and Mocanu [5].

Lemma 2.1 [5]. Let $\phi(u, v)$ be a complex valued function $\phi : D \rightarrow \mathbb{C}$, $D \subset \mathbb{C}^2$ and let $u = u_1 + iu_2$, $v = v_1 + iv_2$. Suppose that the function $\phi(u, v)$ satisfies

- (i) $\phi(u, v)$ is continuous in D ;
- (ii) $(1, 0) \in D$ and $Re \{ \phi(1, 0) \} > 0$;
- (iii) for all $(iu_2, v_1) \in D$ such that $v_1 \leq -\frac{n}{2}(1 + u_2^2)$, $Re \{ \phi(iu_2, v_1) \} \leq 0$.

Let $p(z) = 1 + p_n z^n + p_{n+1} z^{n+1} + \dots$ be regular in \mathbb{U} such that $(p(z), zp'(z)) \in D$ for all $z \in \mathbb{U}$. If $Re \{ \phi(p(z), zp'(z)) \} > 0$ for all $z \in \mathbb{U}$, then $Rep(z) > 0$.

Employing the techniques used by Owa [9] for univalent functions, Noor and Muhammad [8] and Aouf and Seoudy [1] for multivalent functions and Mostafa et al. [6] for meromorphic multivalent functions, we prove the following theorems.

Theorem 2.1. If $f(z) \in \mathcal{M}_{a,b,\mu}^{p,\alpha,\lambda}(\beta, \gamma, \rho, k)$, then

$$\left(z^p \mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z)\right)^\beta \in \mathcal{P}_k(\rho_1), \quad (11)$$

where ρ_1 is given by

$$\rho_1 = \frac{2\mu\beta\rho + n\gamma}{2\mu\beta + n\gamma} \quad (0 \leq \rho_1 < 1). \quad (12)$$

Proof. Let

$$\begin{aligned} \left(z^p \mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z)\right)^\beta &= (1 - \rho_1)p(z) + \rho_1 \\ &= \left(\frac{k}{4} + \frac{1}{2}\right) [(1 - \rho_1)p_1(z) + \rho_1] - \left(\frac{k}{4} - \frac{1}{2}\right) [(1 - \rho_1)p_2(z) + \rho_1], \end{aligned} \quad (13)$$

where $p_i(z)$ is analytic in \mathbb{U} with $p_i(0) = 1$ for $i = 1, 2$. Differentiating (13) with respect to z , and using identity (7) in the resulting equation, we get

$$\begin{aligned} &\left[(1 - \gamma) \left(z^p \mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z)\right)^\beta + \gamma \left(\frac{\mathcal{I}_{a,b,\mu+1}^{p,\alpha,\lambda} f(z)}{\mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z)}\right) \left(z^p \mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z)\right)^\beta \right] \\ &= [(1 - \rho_1)p(z) + \rho_1] + \frac{\gamma(1 - \rho_1)zp'(z)}{\mu\beta} \in \mathcal{P}_k(\rho). \end{aligned}$$

This implies that

$$\frac{1}{1 - \rho} \left\{ [(1 - \rho_1)p_i(z) + \rho_1] - \rho + \frac{\gamma(1 - \rho_1)zp'_i(z)}{\mu\beta} \right\} \in \mathcal{P} \quad (z \in \mathbb{U}; i = 1, 2).$$

Defining the function

$$\phi(u, v) = [(1 - \rho_1)u + \rho_1] - \rho + \frac{\gamma(1 - \rho_1)v}{\mu\beta}$$

where $u = p_i(z) = u_1 + iu_2$, $v = zp'_i(z) = v_1 + iv_2$, we have

- (i) $\phi(u, v)$ is continuous in $D = \mathbb{C}^2$;
- (ii) $(1, 0) \in D$ and $Re \{\phi(1, 0)\} = 1 - \rho > 0$;
- (iii) for all $(iu_2, v_1) \in D$ such that $v_1 \leq -\frac{n}{2}(1 + u_2^2)$,

$$\begin{aligned} Re \{\phi(iu_2, v_1)\} &= \rho_1 - \rho + \frac{\gamma(1 - \rho_1)v_1}{\mu\beta} \\ &\leq \rho_1 - \rho - \frac{n\gamma(1 - \rho_1)(1 + u_2^2)}{2\mu\beta} \\ &= \frac{A + Bu_2^2}{2C}, \end{aligned}$$

where $A = 2(\rho_1 - \rho)\mu\beta - n\gamma(1 - \rho_1)$, $B = -n\gamma(1 - \rho_1)$, $C = \mu\beta > 0$. We note that $Re\{\phi(iu_2, v_1)\} < 0$ if and only if $A = 0$, $B < 0$, this is true from (12). Therefore, by applying Lemma 2.1, $p_i(z) \in \mathcal{P}$ ($i = 1, 2$) and consequently $p(z) \in \mathcal{P}_k(\rho_1)$ for $z \in \mathbb{U}$. This completes the proof of Theorem 2.1.

Using similar arguments to those in the proof of Theorem 2.1 and the identity (8) instead of (7), we obtain the following theorem for the class $\mathcal{N}_{a,b,\mu}^{p,\alpha,\lambda}(\beta, \gamma, \rho, k)$.

Theorem 2.2. If $f(z) \in \mathcal{N}_{a,b,\mu}^{p,\alpha,\lambda}(\beta, \gamma, \rho, k)$, then

$$\left(z^p \mathcal{I}_{a,b,\mu}^{p,\alpha+1,\lambda} f(z)\right)^\beta \in \mathcal{P}_k(\rho_2), \tag{14}$$

where ρ_2 is given by

$$\rho_2 = \frac{2\lambda\rho\beta + n\gamma}{2\lambda\beta + n\gamma} \quad (0 \leq \rho_2 < 1). \tag{15}$$

Theorem 2.3. If $f(z) \in \mathcal{M}_{a,b,\mu}^{p,\alpha,\lambda}(\beta, \gamma, \rho, k)$, then

$$\left(z^p \mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z)\right)^{\beta/2} \in \mathcal{P}_k(\rho_3), \tag{16}$$

where ρ_3 is given by

$$\rho_3 = \frac{n\gamma + \sqrt{(n\gamma)^2 + 4(\mu\beta + n\gamma)\mu\beta\rho}}{2(\mu\beta + n\gamma)} \quad (0 \leq \rho_3 < 1). \tag{17}$$

Proof. Let

$$\begin{aligned} \left(z^p \mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z)\right)^{\beta/2} &= (1 - \rho_3)p(z) + \rho_3 \\ &= \left(\frac{k}{4} + \frac{1}{2}\right) [(1 - \rho_3)p_1(z) + \rho_3] - \left(\frac{k}{4} - \frac{1}{2}\right) [(1 - \rho_3)p_2(z) + \rho_3], \end{aligned} \tag{18}$$

where $p_i(z)$ is analytic in \mathbb{U} with $p_i(0) = 1$ for $i = 1, 2$. Differentiating (18) with respect to z , and using identity (7) in the resulting equation, we get

$$\begin{aligned} &\left[(1 - \gamma) \left(z^p \mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z)\right)^\beta + \gamma \left(\frac{\mathcal{I}_{a,b,\mu+1}^{p,\alpha,\lambda} f(z)}{\mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z)}\right) \left(z^p \mathcal{I}_{a,b,\mu}^{p,\alpha,\lambda} f(z)\right)^\beta \right] \\ &= [(1 - \rho_3)p(z) + \rho_3]^2 + \frac{2\gamma(1 - \rho_3) [(1 - \rho_3)p(z) + \rho_3] zp'(z)}{\mu\beta} \in \mathcal{P}_k(\rho). \end{aligned}$$

This implies that

$$\frac{1}{1-\rho} \left\{ [(1-\rho_3)p_i(z) + \rho_3]^2 - \rho + \frac{2\gamma(1-\rho_3)[(1-\rho_3)p_i(z) + \rho_3]zp'_i(z)}{\mu\beta} \right\} \in \mathcal{P} \quad (z \in \mathbb{U}; i = 1, 2).$$

Defining the function

$$\phi(u, v) = [(1-\rho_3)u + \rho_3]^2 - \rho + \frac{2\gamma(1-\rho_3)[(1-\rho_3)u + \rho_3]v}{\mu\beta}$$

where $u = p_i(z) = u_1 + iu_2$, $v = zp'_i(z) = v_1 + iv_2$, we have

- (i) $\phi(u, v)$ is continuous in $D = \mathbb{C}^2$;
- (ii) $(1, 0) \in D$ and $Re \{ \phi(1, 0) \} = 1 - \rho > 0$;
- (iii) for all $(iu_2, v_1) \in D$ such that $v_1 \leq -\frac{n}{2}(1 + u_2^2)$,

$$\begin{aligned} Re \{ \phi(iu_2, v_1) \} &= -(1-\rho_3)^2u_2^2 + \rho_3^2 - \rho + \frac{2\gamma\rho_3(1-\rho_3)v_1}{\mu\beta} \\ &\leq -(1-\rho_3)^2u_2^2 + \rho_3^2 - \rho - \frac{n\gamma\rho_3(1-\rho_3)(1+u_2^2)}{\mu\beta} \\ &= \frac{A + Bu_2^2}{C}, \end{aligned}$$

where $A = \mu\beta\rho_3^2 - \mu\beta\rho - n\gamma\rho_3(1-\rho_3)$, $B = -(1-\rho_3)[\mu\beta(1-\rho_3) + \gamma n\rho_3]$, $C = \mu\beta > 0$. We note that $Re \{ \phi(iu_2, v_1) \} < 0$ if and only if $A = 0$, $B < 0$, this is true from (17) and $0 \leq \rho_3 < 1$. Therefore, by applying Lemma 2.1, $p_i(z) \in \mathcal{P}$ ($i = 1, 2$) and consequently $p(z) \in \mathcal{P}_k(\rho_3)$ for $z \in \mathbb{U}$. This completes the proof of Theorem 2.3.

Using similar arguments to those in the proof of Theorem 2.3 and the identity (8) instead of (7), we obtain the following theorem for the class $\mathcal{N}_{a,b,\mu}^{p,\alpha,\lambda}(\beta, \gamma, \rho, k)$.

Theorem 2.4. If $f(z) \in \mathcal{N}_{a,b,\mu}^{p,\alpha,\lambda}(\beta, \gamma, \rho, k)$, then

$$\left(z^p \mathcal{I}_{a,b,\mu}^{p,\alpha+1,\lambda} f(z) \right)^\beta \in \mathcal{P}_k(\rho_4), \tag{19}$$

where ρ_4 is given by

$$\rho_4 = \frac{n\gamma + \sqrt{(n\gamma)^2 + 4(\lambda\beta + n\gamma)\lambda\beta\rho}}{2(\lambda\beta + n\gamma)} \quad (0 \leq \rho_4 < 1). \tag{20}$$

Remark. (i) Putting $p = a = b = 1$ in our results, we will obtain the result obtained by Aouf et al [2];

(ii) Putting $a = b = 1$ in our results, we will obtain new results for the classes $\mathcal{M}_\mu^{p,\alpha,\ell}(\beta, \gamma, \rho, k)$ and $\mathcal{N}_\mu^{p,\alpha,\ell}(\beta, \gamma, \rho, k)$ mentioned in the introduction.

REFERENCES

- [1] M. K. Aouf and T. M. Seoudy, *Some properties of certain subclasses of p -valent Bazilevič functions associated with the generalized operator*, Appl. Math. Letters, 24 (2011), no. 11, 1953–1958.
- [2] M. K. Aouf, A. O. Mostafa and H. M. Zayed, *Some characterizations for subclasses of meromorphic Bazilevič functions associated with Cho-Kwon-Srivastava operator*, Int. J. Open Problems Complex Analysis, (To appear).
- [3] N. E. Cho, O. S. Kwon and H. M. Srivastava, *Inclusion and argument properties for certain subclasses of meromorphic functions associated with a family of multiplier transformations*, J. Math. Anal. Appl., 300 (2004), 505–520.
- [4] R. M. El-Ashwah and M. K. Aouf, *Inclusion relationships of certain classes of meromorphic p -valent functions*, Southeast Asian Bull. Math., 36 (2012), 801–810
- [5] S. S. Miller and P. T. Mocanu, *Second order differential inequalities in the complex plane*, J. Math. Anal. Appl., 65 (1978), no. 2, 289–305.
- [6] A. O. Mostafa, M. K. Aouf, A. Shamandy and E. A. Adwan, *Certain subclasses of p -valent meromorphic functions associated with a new operator*, J. Complex Analysis, (2013), 1-4.
- [7] K. I. Noor, *On subclasses of close-to-convex functions of higher order*, Internat. J. Math. Math. Sci., 15 (1992), 279–290.
- [8] K. I. Noor and A. Muhammad, *Some properties of the subclass of p -valent Bazilevič functions*, Acta Univ. Apulensis, no. 17 (2009), 189–197.
- [9] S. Owa, *On certain Bazilevič functions of order β* , Internat. J. Math. Math. Sci., 15 (1992), no. 3, 613-616.
- [10] K. S. Padmanabhan and R. Parvatham, *Properties of a class of functions with bounded boundary rotation*, Ann. Polon. Math., 31 (1975), 311–323.
- [11] K. Piejko and J. Sokol, *Subclasses of meromorphic functions associated with the Cho-Kwon-Srivastava operator*, J. Math. Anal. Appl., 337 (2008), 1261–1266.
- [12] B. Pinchuk, *Functions with bounded boundary rotation*, Israel J. Math., 10 (1971), 7–16.

M. K. Aouf

Department of Mathematics, Faculty of Science, Mansoura University

Mansoura 35516, Egypt

Email: mkaouf127@yahoo.com

A. O. Mostafa

Department of Mathematics, Faculty of Science, Mansoura University

Mansoura 35516, Egypt
Email: *adelaeg254@yahoo.com*

H. M. Zayed
Department of Mathematics, Faculty of Science, Menofia University
Shebin Elkom 32511, Egypt
Email: *hanaa_zayed42@yahoo.com*