

QR-Hypersurfaces of Quaternionic Kähler Manifolds

Vittorio Mangione

**Dedicated to the Memory of Grigorios TSAGAS (1935-2003),
President of Balkan Society of Geometers (1997-2003)**

Abstract

We prove that the basic manifold of a submersion from a QR-hypersurface of a quaternionic Kähler manifold to an almost quaternionic Hermitian manifold is quaternionic Kähler. Then we prove some results involving the sectional curvatures.

Mathematics Subject Classification: 53B35, 53C25, 53C26.

Key words: Kähler manifolds, Hermitian manifolds, quaternionic manifolds, CR-manifolds.

Introduction

Real hypersurfaces of quaternionic space forms have been studied by many authors ([1], [2], [3], [4], [5], [11], [12]) under conditions concerning their shape operator. It is known that real hypersurface of quaternionic Kähler manifolds are not CR-hypersurface in general ([2]).

The study of CR-submanifolds of a quaternionic Kähler manifolds has been carried out in the paper [1]. S. Kobayashi considered the similarity between the total space of a Riemannian submersion and a CR-submanifold of a Kähler manifold in terms of distributions ([9]). In this paper we study Riemannian submersions from QR-hypersurface of a quaternionic Kähler manifold over an almost quaternionic Hermitian manifold (second section). In the last section we study some curvature properties induced on the basic manifold by the submersion.

1 Hypersurfaces of quaternionic Kähler manifolds

We say that a $4(m+1)$ -dimensional manifold \tilde{M} with a metric \tilde{g} is a *quaternionic Kähler manifold* ($m \geq 1$) if there exists a 3-dimensional vector bundle V of tensors of type $(1,1)$ on \tilde{M} satisfying the following conditions:

- (a) In any coordinate neighborhood \tilde{U} on \tilde{M} there is a local basis of almost Hermitian structures $\{\mathcal{J}_a, \tilde{g}\}$, such that $\mathcal{J}_a^2 = -Id$, $a \in \{1, 2, 3\}$ and $\mathcal{J}_a \circ \mathcal{J}_b = -\mathcal{J}_b \circ \mathcal{J}_a = \mathcal{J}_c$ for any cyclic permutation (a, b, c) of $(1, 2, 3)$.
- (b) For any local section φ of V and any tangent vector X to \tilde{M} , $\tilde{\nabla}_X \varphi$ is also a local section in V , where $\tilde{\nabla}$ denotes the Levi-Civita connection of \tilde{g} .

Condition (b) is equivalent to the following:

- (b') There exist local 1-forms ω_{ab} , $a, b \in \{1, 2, 3\}$ on \tilde{U} such that $\omega_{ab} + \omega_{ba} = 0$, and

$$(1) \quad \tilde{\nabla}_x \mathcal{J}_a = \omega_{ab}(x) \mathcal{J}_b + \omega_{ac}(x) \mathcal{J}_c$$

for any cyclic permutation (a, b, c) of $(1, 2, 3)$.

Given two local bases $\{\mathcal{J}_a\}$ and $\{\mathcal{J}'_a\}$ of V defined on coordinate neighborhoods \tilde{U} and \tilde{U}' such that $\tilde{U} \cap \tilde{U}' \neq \emptyset$, we have on $\tilde{U} \cap \tilde{U}'$:

$$(2) \quad \mathcal{J}'_a = \sum_{b=1}^3 C_{ab} \mathcal{J}_b$$

where $[C_{ab}]$ is an element of the special orthogonal group $SO(3)$ (see [8]).

Let \tilde{M} be an orientable hypersurface of \tilde{M} and ξ a unit normal field defined on M . On \tilde{U} $\xi_a = -\mathcal{J}_a(\xi)$, $a \in \{1, 2, 3\}$ defines a tangent vector field to M . Similarly, we define ξ'_a on \tilde{U}' and on $\tilde{U} \cap \tilde{U}' \neq \emptyset$ we have:

$$(3) \quad \xi'_a = \sum_{b=1}^3 C_{ab} \xi_b, \quad b \in \{1, 2, 3\}$$

so that one obtains a distribution \mathcal{V} on M which is locally represented by $\{\xi_a\}$, $1 \leq a \leq 3$, on \tilde{U} . Let \mathcal{H} be the orthogonal complementary distribution to \mathcal{V} with respect to the Riemannian metric g induced by \tilde{g} on M .

We see that for each $x \in M$, \mathcal{H}_x is \mathcal{J}_a -invariant, but \mathcal{V}_x is not an anti-invariant subspace of $T_x \tilde{M}$ with respect \mathcal{J}_a , $a = \{1, 2, 3\}$. It is easy to see that $\mathcal{J}_a(\mathcal{V}_x) = T_x M^\perp$, $x \in M$, where $T_x M^\perp$ is the normal space at x to the hypersurface M in \tilde{M} . In general, when the previous conditions are satisfied, we say that M is a *QR-hypersurface* of \tilde{M} (see [3]). Now, let B be the second fundamental form of M in \tilde{M} . Then, for any $E, F \in \Gamma(TM)$ we have the Gauss formula

$$(4) \quad \tilde{\nabla}_E F = \nabla_E F + B(E, F),$$

where $\tilde{\nabla}$ and ∇ are the Levi-Civita connections on \tilde{M} and M , respectively.

If L denotes the fundamental tensor of Weingarten with respect to ξ , we have the Weingarten formula

$$(5) \quad \tilde{\nabla}_E \xi = -L(E),$$

and for any $E, F \in \Gamma(TM)$ the following formula

$$(6) \quad g(L(E), F) = g(B(E, F), \xi)$$

holds.

The integrability of the distributions \mathcal{V} and \mathcal{H} on M has been studied by A. Bejancu ([2]).

We recall that the vertical distribution \mathcal{V} is integrable if and only if

$$(7) \quad B(U, X) = 0$$

for any $U \in \Gamma(\mathcal{V})$ and $X \in \Gamma(\mathcal{H})$.

If (7) is satisfied, we say that M is a *mixed geodesic QR-hypersurface* of \tilde{M} .

2 Riemannian submersions of QR-hypersurfaces

Let M be a mixed geodesic QR-hypersurface of a quaternionic Kähler manifold \tilde{M} . We denote by (M', g', \mathcal{J}'_a) , $a \in \{1, 2, 3\}$, an almost quaternionic Hermitian manifold (i.e. satisfying the condition (a)). We say that a Riemannian submersion $\pi: M \rightarrow M'$ is a *QR-submersion* if the following conditions are satisfied:

- i) \mathcal{V} is the kernel of π_* ;
- ii) for each $x \in M$, $\pi_* : \mathcal{H}_x \rightarrow T_{\pi(x)}M'$ is an isometry with respect to each complex structure of H_x and $T_{\pi(x)}M'$, where $T_{\pi(x)}M'$ denotes the tangent space to M' at $\pi(x)$.

As in the paper [10], the letters U, V, W, W' will always denote vertical vector fields and X, Y, Z, Z' horizontal vector fields. A horizontal vector field X on M is said to be *basic* if it is π -related to a vector field X' on M' .

We denote by T and A O'Neill's fundamental tensors (see [13], [11]).

Lemma 2.1 *Let X and Y be basic vector fields on M . Then the following conditions hold:*

- a) *The horizontal component $h[X, Y]$ of $[X, Y]$ is a basic vector field and $\pi_* h[X, Y] = [X', Y'] \circ \pi$;*
- b) *$h(\nabla_X Y)$ is basic vector field corresponding to $\nabla'_{X'} Y'$ where ∇ and ∇' are the Levi-Civita connections on M and M' , respectively;*
- c) *$[X, U] \in \Gamma(\mathcal{V})$, for any vertical field $U \in \Gamma(\mathcal{V})$;*

where h denotes the horizontal component of a vector E on M .

We define a skew-symmetric tensor field C by

$$(1) \quad \tilde{\nabla}_x Y = h\tilde{\nabla}_x Y + C(X, Y)$$

for all $X, Y \in \Gamma(\mathcal{H})$.

The second fundamental form B of M in \tilde{M} is:

$$(2) \quad B(E, F) = \tilde{\nabla}_E F - \nabla_E F$$

for all $E, F \in \Gamma(TM)$.

Theorem 2.2 *Let M be a mixed geodesic QR-hypersurface of a quaternionic Kähler manifold \tilde{M} . If $\pi : M \rightarrow M'$ is a QR-submersion of M on an almost quaternionic Hermitian manifold, then M' is a quaternionic Kähler manifold.*

Proof. By using Gauss formula and (1) we obtain

$$(3) \quad h\nabla_X \mathcal{J}_a Y - \mathcal{J}_a(h\nabla_X Y) = \omega_{ab}(X) \mathcal{J}_b Y + \omega_{ac}(X) \mathcal{J}_c Y,$$

for any local basic vector fields X, Y on M and for any cyclic permutation (a, b, c) of $(1, 2, 3)$. Then we can define 1-forms ω'_{ab} on M' by

$$(4) \quad \omega'_{ab}(X') \circ \pi = \omega_{ab}(X), \quad a, b, c \in \{1, 2, 3\},$$

for any local vector field X' on M' and X a real basic vector field on M such that $\pi_* X = X'$.

On the other hand, by the definition of a QR-submersion we have

$$(5) \quad \pi_* \circ \mathcal{J}_a = \mathcal{J}'_a \circ \pi_*.$$

Using Lemma 2.1, from (3)-(5) we obtain

$$h(\nabla'_X \mathcal{J}'_a) Y' = \omega'_{ab}(X') \mathcal{J}'_b Y' + \omega'_{ac}(X') \mathcal{J}'_c Y',$$

where ∇' is the Levi-Civita connection on M' and X', Y' any local vector fields on M' . We conclude that (M', \mathcal{J}'_a, g') is a quaternionic Kähler manifold. \square

3 Totally umbilical QR-hypersurfaces

In the sequel we shall denote by $\langle \cdot, \cdot \rangle$ the scalar product induced on the tangent spaces of M and \tilde{M} by the Riemannian metric g . We recall that a hypersurface M of \tilde{M} is totally umbilical if the first and the second fundamental forms are proportional, that is

$$(1) \quad B(E, F) = \langle E, F \rangle H$$

for any $E, F \in \Gamma(TM)$, where H is the mean curvature vector of M , defined by the formula,

$$(2) \quad H = \frac{1}{4m+3} \text{Trace} B.$$

We have the Gauss equation:

$$(3) \quad \tilde{R}(E, E', F, F') = R(E, E', F, F') - \langle B(E, F), B(E', F') \rangle + \langle B(E, F'), B(F, E') \rangle.$$

Taking account of the formula (1), the Gauss equation for a totally umbilical hypersurface M in \tilde{M} becomes:

$$(4) \quad \begin{aligned} \tilde{R}(E, E', F, F') &= R(E, E', F, F') - (\langle E, F \rangle \langle E', F' \rangle + \\ &\quad - \langle E, F' \rangle \langle F, F' \rangle) \|H\|^2, \end{aligned}$$

where $\|H\|^2 = \langle H, H \rangle$.

We see that, if M is a totally umbilical QR-hypersurface of \tilde{M} , then it is a mixed geodesic QR-hypersurface, i.e. $B(V, X) = 0$ for any $V \in \Gamma(\mathcal{V})$ and $X \in \Gamma(\mathcal{H})$. Consequently, the vertical distribution \mathcal{V} is integrable.

Moreover, it is easy to check that each leaf of \mathcal{V} is totally geodesic in M (see, for example [3], p. 121). Then we conclude that the first fundamental tensor T of the Riemannian submersion $\pi : M \rightarrow M'$ vanishes, because $T_U V$ is the second fundamental form of each fibre for any $U, V \in \Gamma(\mathcal{V})$ (see [7], [13]).

Let us now recall the following two Gray-O'Neill curvature equations for a Riemannian submersion:

$$(5) \quad \begin{aligned} R(U, V, U', V') &= \hat{R}(U, V, U', V') + \langle T_U V', T_V U' \rangle + \\ &\quad - \langle T_V V', T_U U' \rangle, \end{aligned}$$

$$(6) \quad \begin{aligned} R(X, Y, X', Y') &= R^*(X, Y, X', Y') + 2\langle C(X, Y), C(X', Y') \rangle + \\ &\quad + \langle C(Y, X'), C(X, Y') \rangle - \langle C(X, X'), C(Y, Y') \rangle, \end{aligned}$$

for all $U, V, U', V' \in \Gamma(\mathcal{V})$ and $X, Y, X', Y' \in \Gamma(\mathcal{H})$, where, for any quadruplet of horizontal vector fields (X, Y, X', Y') , $R^*(X, Y, X', Y') = R'(\pi_* X, \pi_* Y, \pi_* X', \pi_* Y') \circ \pi$, with R^* Riemannian curvature on the fibres of \mathcal{H} . Here R' is the Riemannian curvature of the metric g' on M' .

Lemma 3.1 *Let M be a totally umbilical, not totally geodesic, QR-hypersurface of a quaternionic Kähler manifold. Then the tensor field C which measures the integrability of the horizontal distribution \mathcal{H} , is given by the formula*

$$(7) \quad C(X, Y) = \|H\| \sum_{a=1}^3 \langle X, \mathcal{J}_a Y \rangle \xi_a.$$

Proof. Using (1), (4), (5) and (7), we obtain

$$(8) \quad \mathcal{J}_a(LX) - \nabla_X \xi_a = \omega_{ac}(X) \xi_b - \omega_{ab}(X) \xi_c,$$

for any $X \in \Gamma(\mathcal{H})$. Now, by (6) in (8), we have

$$(9) \quad \langle \nabla_X Y, \xi_a \rangle = \langle B(X, \mathcal{J}_a Y), \xi \rangle,$$

for any $X, Y \in \Gamma(\mathcal{H})$, and $a \in \{1, 2, 3\}$. Taking into account that the mean curvature vector H of M is a global vector field and it is non vanishing on M (see [3]), we take

$\xi = \frac{H}{\|H\|}$. Then we have

$$(10) \quad C(X, Y) = V \nabla_X Y = \|H\| \sum_{a=1}^3 \langle X, \mathcal{J}_a Y \rangle \xi_a,$$

from which formula (7) follows. \square

Theorem 3.1 *Let M be a totally umbilical, not totally geodesic, QR-hypersurface of a quaternionic Kähler manifold. Then,*

- a) $\tilde{K}(U, V) = K(U, V) - \|H\|^2$, where $\{U, V\}$ is an orthonormal basis of the vertical 2-plane α , $\alpha \subset \mathcal{V}_x$, $x \in M$, and \tilde{K}, K denote the sectional curvatures of α on \tilde{M}, M , respectively.
- b) $K(X, Y) = K'(X', Y') - 3\|H\|^2 \sum_{a=1}^3 \langle X, \mathcal{J}_a Y \rangle^2$, where X, Y is an orthonormal basis of a horizontal 2-plane $\alpha \subset \mathcal{H}_x$, $K(X, Y)$ denoting the sectional curvature of α , and $K'(X', Y')$ denotes the sectional curvature in M' of the 2-plane spanned by $X' = \pi_* X$ and $Y' = \pi_* Y$.

Proof. Property a) is easily obtained from (4) and (5). From (6), as an immediate consequence of the skew-symmetry of C , we have

$$(11) \quad R(X, Y, X, Y) = R'(X', Y', X', Y') - 3\|C(X, Y)\|^2.$$

Lemma 3.1 and (11) directly give b). \square

We recall that a totally umbilical, not totally geodesic, hypersurface M of a Riemannian manifold \tilde{M} is an extrinsic hypersphere if the mean curvature vector field H is parallel with respect to the linear normal connection ∇^\perp or, equivalently, $\|H\| = c$ is a constant $c \neq 0$ on M .

Then we have the following

Theorem 3.2 *Let M be an extrinsic hypersurface of a flat quaternionic Kähler manifold \tilde{M} and $\pi : M \rightarrow M'$ a QR-submersion of M on a quaternionic Kähler manifold M' . Then M' is a quaternionic Kähler manifold with constant quaternionic sectional curvature $c > 0$.*

Proof. By (4), (6) and Lemma 3.1 we have

$$\begin{aligned} R'(X', Y')Z' &= \|H\|^2 \{g'(Y', Z')X' - g'(X', Z')Y' + \\ &+ \sum_{a=1}^3 (g'(\mathcal{J}'_a Y', Z')\mathcal{J}'_a X' - g'(\mathcal{J}'_a X', Z')\mathcal{J}'_a Y') + \\ &+ 2g'(X', \mathcal{J}'_a Y')\mathcal{J}'_a Z'\}. \end{aligned}$$

where $\|H\|$ is a constant on M' and $X', Y', Z' \in \Gamma(TM')$. \square

Remark There exist no proper totally umbilical QR-submanifolds in positively or negatively curved quaternionic Kähler manifolds (see [3]).

Acknowledgment: The author thanks to Prof. V. Balan for his suggestions on improving the exposition and for pointing out some errors.

References

- [1] M. Barros; B.Y. Chen; F. Urbano, *Quaternion CR-submanifolds of quaternion manifolds*, Kodai Math.J.t. 4 (1981), 399-417.
- [2] A. Bejancu, *Geometry of CR-submanifolds*, Kluwer, 1986.
- [3] A. Bejancu, *On totally umbilical QR-Submanifold of quaternionic Kählerian manifolds*, Bull.Austral.Math.Soc. 62 (2000), 95-103.
- [4] J. Berndt, *Real hypersurfaces in quaternionic spaces forms*, J. reine angew. Math. 419 (1991), 9-26.
- [5] B.Y. Chen, *Geometry of submanifolds and its applications*, Sci. University, Tokyo, 1981.
- [6] A. Gray, *Minimal varieties and almost Hermitian manifolds*, Michigan Math. J. 12 (1955), 273-287.
- [7] A. Gray, *Pseudo-Riemannian almost product manifold and submersions*, J. Math. Mech. 16 (1967), 715-737.
- [8] S. Ishihara, *Quaternion Kählerian manifolds*, J. Diff. Geometry 9 (1974), 483-500.
- [9] S. Kobayashi, *Submersions of CR-submanifolds*, Tohoku Math. J. 39 (1987), 95-100.
- [10] V. Mangione, *Some submersion of CR-hypersurfaces of Kähler-Einstein manifolds*, International J. Math. Math. Sci.; to appear.
- [11] S. Marchiafava, *Sulla geometria locale delle varietà Kähleriane quaternionali*, Boll. UMI 7 (1991), 417-447.
- [12] A. Martinez; J. Perez, *Real hypersurfaces in quaternionic projective space*, Annali Mat. Pura Appl. 145 (1986), 355-384.
- [13] B. O'Neill, *The Fundamental equations of a submersion*, Michigan Math.J. 13 (1966), 459-469.
- [14] C. Udriste, *Structures presque coquaternionnes*, Bull. Soc. Sc. Math. Roumanie 12 (1969), 487-507.

Dipartimento di Matematica, Università
degli Studi di Parma
Via M. D'Azeglio, 85A, 43100 Parma, Italia.
email: vittorio.mangione@unipr.it