# An Almost Paracontact Structure on the Indicatrix Bundle of a Finsler Space

Manuela Gîrţu

#### Abstract

In a paper by I. Hasegawa, K. Yamaguchi and H. Shimada, [2], it was proved that the indicatrix bundle of a Finsler space  $F^n = (M, L)$  has a natural almost contact structure. On a different way, the same structure was found by M. Anastasiei in [1]. Adopting the approach from [1] we prove that the indicatrix bundle of  $F^n$  carries also an almost paracontact structure.

#### Mathematics Subject Classifications 2000: 53C60

Key words and phrases: tangent manifold, framed f-structure, indicatrix bundle, almost paracontact structure.

### 1 Introduction

Let  $F^n = (M, L)$  be a Finsler space. Here M is a real  $C^{\infty}$  manifold of dimension n with local coordinates  $(x^i), i, j, k... = 1, ..., n$ . For the tangent manifold TM with the projection  $\tau$  over M we take the local coordinates  $(x^i \circ \tau, y^i)$ , where  $y^i$  are the components of a vector from  $T_pM$ , in the natural basis  $\partial_i = \frac{\partial}{\partial x^i}$ .

The function  $L: T_0M: TM \setminus \{0\} \to \mathbf{R}_+$  is smooth, positively homogeneous of degree 1 with respect to  $y^i$  and the matrix  $\left(g_{ij}(x,y) = \frac{1}{2}\frac{\partial^2 L^2}{\partial y^i \partial y^j}\right)$  is of rank n. We

set  $\dot{\partial}_i = \frac{\partial}{\partial u^i}$ .

The homogeneity of L implies

$$L^{2}(x,y) = g_{ij}(x,y)y^{i}y^{j} = y^{i}y_{i}$$
 for  $y_{i} = g_{ij}y^{j}$ .

The functions  $N_j^i(x,y) = \frac{1}{2}\dot{\partial}_j(\gamma_{00}^i)$ , for  $\gamma_{00}^i = \gamma_{jk}^i(x,y)y^jy^k$  and  $\gamma_{jk}^i(x,y)$  the "generalized" Christoffel symbols, are the local coefficients of the nonlinear Cartan connection. See [Ch. VIII, 4] for details. One considers a new local basis  $\{\delta_i, \dot{\partial}_i\}$ , with  $\delta_i = \partial_i - N_i^k(x,y)\dot{\partial}_k$ , on  $T_0M$ . Its dual basis is  $(dx^i, \delta y^i)$  with  $\delta y^i = dy^i + N_k^i(x,y)dx^k$ . If we assume that the quadratic form  $g_{ij}(x,y)\xi^i\xi^j$ ,  $\xi \in \mathbf{R}^n$  is positive definite, then

Bałkan Journal of Geometry and Its Applications, Vol.7, No.2, 2002, pp. 43-48.

<sup>©</sup> Balkan Society of Geometers, Geometry Balkan Press 2002.

$$G_S = g_{ij}(x, y) dx^i \otimes dx^j + g_{ij}(x, y) \delta y^i \otimes \delta y^j$$

is a Riemannian metric on  $T_0M$ .

The linear operator P given in the local basis by

(1.1) 
$$P(\delta_i) = \delta_i, \quad P(\dot{\partial}_i) = -\dot{\partial}_i,$$

defines an almost product structure on  $T_0M$  and we have

(1.2) 
$$G_S(PX, PY) = G_S(X, Y), \quad X, Y \in \chi(T_0M).$$

Here  $\chi(T_0M)$  is the module of vector fields on  $T_0M$ . The vector field  $C = y^i \dot{\partial}_i$  is called the Liouville vector field on  $T_0M$  and  $S = y^i \delta_i$  is the geodesic spray of  $F^n$ .

An almost paracontact structure on a manifold N is a set  $(\varphi, \xi, \eta)$ , where  $\varphi$  is a tensor field of type (1, 1),  $\xi$  a vector field and  $\eta$  an 1-form such that

(1.3) 
$$\eta(\xi) = 1, \ \varphi(\xi) = 0, \ \eta \circ \varphi = 0, \ \varphi^2 = +I - \eta \otimes \xi,$$

where I denotes the Kronecker tensor field.

This structure generalizes as follows. One considers on a manifold N of dimension (2n + s) a tensor field f of type (1, 1). If there exists on N the vector fields  $(\xi_{\alpha})$  and the 1- forms  $(\eta^{\alpha})$   $(\alpha = 1, 2, ...s)$  such that

(1.4) 
$$\eta^{\alpha}(\xi_{\beta}) = \delta^{\alpha}_{\beta}, f(\xi_{\alpha}) = 0, \eta^{\alpha} \circ f = 0, f^{2} = I - \sum_{\alpha} \eta^{\alpha} \otimes \xi_{\alpha},$$

then  $(f, (\xi_{\alpha}), (\eta^{\alpha}))$  is called a framed f(3, -1)- structure. The term was suggested by the equation  $f^3 - I = 0$ . This is in some sense dual to the framed f-structure which generalizes the almost contact structure and which may be called a framed f(3, +1)structure. For an account of such kind of structures we refer to the book [3].

In the following (Section 2) we show that the slit tangent bundle  $T_0M$  of a Finsler space carries a natural framed f(3, -1)- structure. The set  $I(M) = \{(x, y) \mid L(x, y) = 1\}$  is a (2n - 1)- dimensional submanifold of  $T_0M$ . In Section 3 we prove that the framed f(3, -1)- structure on  $T_0M$  induces on I(M) an almost paracontact structure. We note that it was known that I(M) carries an almost contact structure [2], [1] but only the approach from [1] allowed us to construct this almost paracontact structure.

# **2** A framed f(3, -1)- structure on $T_0M$

Let us put  $\xi_1 := S = y^i \delta_i$  and  $\xi_2 := C = y^i \dot{\partial}_i$ . By a direct calculation one finds (*P* is the almost product structure (1.1)).

**Lemma 2.1.**  $P(\xi_1) = \xi_1$ ,  $P(\xi_2) = -\xi_2$ . We consider the 1- forms

$$\eta^1 = \frac{y_i}{L^2} dx^i, \eta^2 = \frac{y_i}{L^2} \delta y^i$$

and we prove

**Lemma 2.2.**  $\eta^1 \circ P = \eta^1, \eta^2 \circ P = -\eta^2$ . **Proof.** It is sufficient to check these equalities on the adapted basis  $(\delta_i, \dot{\partial}_i)$ . We have

$$(\eta^1 \circ P)(\delta_i) = \eta^1(P(\delta_i)) = \eta^1(\delta_i) \text{ and } (\eta^1 \circ P)(\dot{\partial}_i) = -\eta^1(\dot{\partial}_i) = 0$$

An Almost Paracontact Structure

Then

$$(\eta^2 \circ P)(\delta_i) = \eta^2(\delta_i) = 0$$
 and  $(\eta^2 \circ P)(\dot{\partial}_i) = -\eta^2(\dot{\partial}_i).$ 

Let be  $G = \frac{1}{L^2} G_S$  a Riemannian metric which is conformal with  $G_S$ . Lemma 2.3.  $\eta^1(X) = G(X, \xi_1), \ \eta^2(X) = G(X, \xi_2), \forall X \in \chi(T_0M)$ . Proof. It is sufficient to check these equalities on the basis  $(\delta_i, \dot{\partial}_i)$ . We have:  $\eta^1(\delta_j) = \frac{y_j}{L^2} = di \frac{1}{L^2} g_{jk} y^k$  and  $G(\delta_j, \xi_1) = \frac{1}{L^2} G_S(\delta_j, y^k \delta_k) = \frac{1}{L^2} y^k G_S(\delta_j, \delta_k) = \frac{1}{L^2} y^k g_{jk}$ . Further,  $\eta^1(\dot{\partial}_i) = 0$  and  $G(\dot{\partial}_j, \xi_1) = \frac{1}{L^2} G_S(\dot{\partial}_i, y^k \delta_k) = 0$ . Similarly, one checks the equation  $\eta^2(X) = G(X, \xi_2)$ .

Now we define a tensor field p of type (1,1) on  $T_0M$  by

(2.1) 
$$p(X) = P(X) - \eta^{1}(X)\xi_{1} + \eta^{2}(X)\xi_{2}, X \in \chi(T_{0}M).$$

This can be written in a more compact form as  $p = P - \eta^1 \otimes \xi_1 + \eta^2 \otimes \xi_2$ . **Theorem 2.1.** For the data  $(p, (\xi_a), (\eta^a))$ , a = 1, 2 the following hold

(i)  $\eta^{a}(\xi_{b}) = \delta^{a}_{b}, \ p(\xi_{a}) = 0, \ \eta^{a} \circ p = 0,$ 

(ii) 
$$p^2 = I - \eta^1 \otimes \xi_1 - \eta^2 \otimes \xi_2, X \in \chi(T_0 M),$$

(iii) *p* is of rank 2n - 2 and  $p^3 - p = 0$ .

**Proof.** (i) follows easily from Lemmas 2.1, 2.2 and the formula (2.1). For (ii) we have

$$p^{2}(X) = p(p(X)) = P(P(X) - \eta^{1}(X)\xi_{1} + \eta^{2}(X)\xi_{2}) - \eta^{1}(P(X) - \eta^{1}(X)\xi_{1} + \eta^{2}(X)\xi_{2}) + \eta^{2}(P(X) - \eta^{1}(X)\xi_{1} + \eta^{2}(X)\xi_{2}) = +X - \eta^{1}(X)\xi_{1} - \eta^{2}(X)\xi_{2},$$

the other terms vanish or cancel because of Lemmas 2.1, 2.2 and (i). Applying p to the equality (ii) and using again the Lemmas 2.1, 2.2 and (i) one gets  $p^3 - p = 0$ . From the second equation in (i) we see that the subspace  $span(\xi_1, \xi_2)$  is contained in Ker p. Let now  $X = X^i \delta_i + Y^i \dot{\partial}_i \in Kerp$ . On using (2.1),

$$p(X) = X^i \delta_i - Y^i \dot{\partial}_i - (X^i \frac{y_i}{L^2})\xi_1 + Y^i \frac{y_i}{L^2}\xi_2 = (X^i - \frac{(X^k y_k)}{L^2}y^i)\delta_i - (Y^i - (Y^k \frac{y_k}{L^2})y^i)\dot{\partial}_i = 0$$

equivalent to

$$X^i = \frac{X^k y_k}{L^2} y^i, \quad Y^i = \frac{(Y^k y_k)}{L^2} y^i.$$

Hence  $X = \frac{X^k y_k}{L^2} \xi_1 + \frac{Y^k y_k}{L^2} \xi_2$  that is X belongs to  $span(\xi_1, \xi_2)$ . In other words, Ker  $p = span(\xi_1, \xi_2)$ . Thus rank p = 2n - 2.

**Theorem 2.2.** The Riemannian metric  $G = \frac{1}{L^2}G_S$  satisfies

(2.2) 
$$G(pX, pY) = G(X, Y) - \eta^{1}(X)\eta^{1}(Y) - \eta^{2}(X)\eta^{2}(Y), X, Y \in \chi(T_{0}M).$$

**Proof.** Use (2.1) and Lemma 2.3 and Lemma 2.1 as well as  $G(\xi_1, \xi_1) = 1$ ,  $G(\xi_2, \xi_2) = 1$ ,  $G(\xi_1, \xi_2) = 0$  to obtain

$$\begin{split} G(pX,pY) &= G(PX,PY) - \eta^1(Y)G(PX,\xi_1) + \eta^2(Y)G(PX,\xi_2) - \\ &-\eta^1(X)G(\xi_1,PY) + \eta^1(X)\eta^1(Y) + \eta^2(X)G(\xi_2,PY) + \eta^2(X)\eta^2(Y) = \\ &= G(X,Y) - \eta^1(Y)\eta^1(P(X)) + \eta^2(Y)\eta^2(PX) - \eta^1(X)\eta^1(PY) + \eta^2(X)\eta^2(PY) + \\ &+ \eta^1(X)\eta^1(Y) + \eta^2(X)\eta^2(Y) = G(X,Y) - \eta^1(X)\eta^1(Y) - \eta^2(X)\eta^2(Y) \end{split}$$

**Remark.** In the local basis  $(\delta_i, \dot{\partial}_i)$ , we get

(2.3)  

$$G(p(\delta_i), p(\delta_j)) = \frac{1}{L^2} (g_{ij} - \frac{y_i y_j}{L^2}), \ G(p(\delta_i), p(\dot{\partial}_i)) = 0,$$

$$G(p(\dot{\partial}_i), \ p(\dot{\partial}_j)) = \frac{1}{L^2} (g_{ij} - \frac{y_i y_j}{L^2}).$$

Let us put

(2.4) 
$$h(X,Y) = G(pX,Y), X, Y \in \chi(T_0M).$$

We have

**Theorem 2.3.** The map h is a symmetric bilinear form on  $T_0M$  of rank 2n-2, with the null space span $(\xi_1, \xi_2)$ .

**Proof.** h is bilinear since G is so. As for the symmetry we have

$$\begin{split} h(Y,X) &= G(pY,X) = G(pY,p^2X + \eta^1(X)\xi_1 + \eta^2(X)\xi_2) = \\ &= G(pY,p(pX)) + \eta^1(X)G(pY,\xi_1) + \eta^2(X)G(pY,\xi_2) = \\ &= G(pY,p(pX)) + \eta^1(X)\eta^1(PY) + \eta^2(X)\eta^2(PY) = \\ &= G(Y,pX) - \eta^1(Y)\eta^1(pX) - \eta^2(Y)\eta^2(pX) = G(Y,pX) = h(X,Y). \end{split}$$

Then we have  $h(\xi_1, \xi_1) = h(\xi_2, \xi_2) = 0$ . Thus  $span(\xi_1, \xi_2)$  is contained in the null space of h. Conversely, if  $X = X^i \delta_i$  is such that  $h(X, X) = 0 \iff G(pX, X) = 0$  it results  $X = \frac{X^k y_k}{L^2} \xi_1$  and similarly, if  $X = Y^i \dot{\partial}_i$  is such that h(X, X) = 0, it results  $X = \frac{Y^k y_k}{L^2} \xi_2$ . Thus the null space of h is just  $span(\xi_1, \xi_2)$  and the proof is finished.

**Remark.** The map h is a singular pseudo-Riemannian metric on  $T_0M$ . Locally it looks as follows

$$h = \frac{1}{L^2} (g_{ij} - \frac{y_i y_j}{L^2}) dx^i \otimes dx^j - \frac{1}{L^2} (g_{ij} - \frac{y_i y_j}{L^2}) \delta y^i \otimes \delta y^j,$$

with

$$\operatorname{rank}\left(g_{ij} - \frac{y_i y_j}{L^2}\right) = n - 1$$

since

$$(g_{ij} - \frac{y_i y_j}{L^2})y^j = y_i - y_i = 0$$
  $(y_j y^j = L^2).$ 

46

## 3 An almost paracontact structure on the indicatrix bundle of the Finsler space $F^n = (M, L)$

The indicatrix bundle of  $F^n$  is the submanifold

$$I(M) = \{(x, y) \in T_0M \mid L(x, y) = 1\}$$

of  $T_0M$  projected over M. It is well-known that  $\xi_2 = y^i \dot{\partial}_i$  is normal to I(M) and this is unitary with respect to G since

$$G(\xi_1, \xi_2) = \frac{1}{L^2} y^i y^j g_{ij} = 1.$$

We consider  $T_0M$  with the Riemannian metric G and then I(M) appears as a hypersurface of  $T_0M$  with normal vector field  $\xi_2$ . We restrict to I(M) all the objects introduced above and indicate this fact by putting a bar over the letters denoting those objects. We have:

- $\overline{\xi}_1 = \xi_1$  since  $\xi_1$  is tangent to I(M),
- $\overline{\eta}^2 = 0$  on I(M) since  $\eta^2(X) = G(X, \xi_2) = 0$  for  $X \in \chi(I(M))$ ,
- $\overline{G} = G_S \mid_{I(M)}$  because  $L^2 = 1$  on I(M),
- $\overline{p}(X) = P(X) \overline{\eta}^1(X)\xi_1$  for  $X \in \chi(I(M))$ .
- The map  $\overline{p}$  is an endomorphism of the tangent bundle to I(M) since  $G(\overline{p}X,\xi_2) = 0$ .

We put  $\overline{\xi}_1 = \overline{\xi}, \overline{\eta}^1 = \overline{\eta}$  and as a consequence of the Theorem 2.1 we get **Theorem 3.1.** The triple  $(\overline{p}, \overline{\xi}, \overline{\eta})$  defines an almost paracontact structure on I(M), that is,

(i)  $\overline{\eta}(\overline{\xi}) = 1, \overline{p}(\overline{\xi}) = 0, \overline{\eta} \circ \overline{p} = 0,$ 

(ii) 
$$\overline{p}^2(X) = X - \overline{\eta}(X)\xi, X \in \chi(I(M))),$$

(iii)  $\overline{p}^3 - p = 0$ , rank  $\overline{p} = 2n - 2 = (2n - 1) - 1$ .

Using the restriction to I(M) and the Theorem 2.2 one infers **Theorem 3.2.** The Riemannian metric  $\overline{G}$  satisfies

(3.1) 
$$\overline{G}(\overline{p}X,\overline{p}Y) = \overline{G}(X,Y) - \overline{\eta}(X)\overline{\eta}(Y), X, Y \in \chi(I(M)).$$

From the last two theorems we see that the ensemble  $(\overline{p}, \overline{\xi}, \overline{\eta}, \overline{G})$  defines an almost metrical paracontact structure on I(M).

### References

 Anastasiei, M., A framed f-structure on tangent manifold of a Finsler space, Analele Univ. Bucureşti, Mat.- Inf., XLIX, 2000, 3-9.

- [2] Hasegawa, I., Yamaguchi, K., Shimada, H., Sasakian structures on Finsler manifolds in P.L. Antonelli and R. Miron (eds.), Lagrange and Finsler Geometry, Kluwer Academic Publishers, 1996, p. 75–80.
- [3] Mihai, I., Roşca, R., Verstraelen, L., Some aspects of the differential geometry of vector fields, PADGE, Katholieke Universiteit Leuven, vol.2, 1996.
- [4] Miron, R., Anastasiei, M., The geometry of Lagrange spaces: theory and applications, Kluwer Academic Publ., FTPH 59, 1994.

Faculty of Sciences University of Bacău 5500, Bacău, Romania