

REMARKS ON FILTRATIONS
OF THE HOMOLOGY OF REAL VARIETIES

JEREMIAH HELLER AND MIRCEA VOINEAGU¹

Received: October 16, 2011

Revised: March 30, 2012

Communicated by Alexander Merkurjev

ABSTRACT. We demonstrate that a conjecture of Teh which relates the niveau filtration on Borel-Moore homology of real varieties and the images of generalized cycle maps from reduced Lawson homology is false. We show that the niveau filtration on reduced Lawson homology is trivial and construct an explicit class of examples for which Teh's conjecture fails by generalizing a result of Schülting. We compare different cycle maps and in particular we show that the Borel-Haefliger cycle map naturally factors through the reduced Lawson homology cycle map.

2010 Mathematics Subject Classification: 14F43, 14C25

CONTENTS

1. Introduction	642
2. Preliminaries	644
3. Birational invariants and examples	646
4. Coniveau spectral sequences	649
5. Filtrations in homology	652
6. Cycle Maps	653
References	659

¹The second author was partially supported by JSPS Grant in Aid (B), No. 23740006

1. INTRODUCTION

Let X be a quasi-projective real variety. In [Teh10] the reduced Lawson homology groups $RL_q H_n(X)$ are introduced as homotopy groups of certain spaces of “reduced” algebraic cycles. For $q = 0$ we have $RL_0 H_n(X) = H_n(X(\mathbb{R}); \mathbb{Z}/2)$, where $H_n(X(\mathbb{R}); \mathbb{Z}/2)$ is the Borel-Moore homology. At the other extreme we have that $RL_n H_n(X)$ is a quotient of the Chow group $CH_n(X)$. There are generalized cycle maps

$$cyc_{q,n} : RL_q H_n(X) \rightarrow H_n(X(\mathbb{R}); \mathbb{Z}/2)$$

and the images of these cycle maps form a filtration of the homology

$$Im(cyc_{n,n}) \subseteq Im(cyc_{n-1,n}) \subseteq \cdots \subseteq Im(cyc_{0,n}) = H_n(X(\mathbb{R}); \mathbb{Z}/2).$$

The first step of this filtration is the image of the Borel-Haeflinger cycle map $Im(cyc_{n,n}) = H_n(X(\mathbb{R}); \mathbb{Z}/2)_{alg}$ (see Theorem 6.3).

The construction of the reduced Lawson homology is based on Friedlander’s construction of Lawson homology groups for complex varieties. Friedlander-Mazur [FM94] have conjectured a relationship between the filtration on singular homology of the space of complex points given by images of the generalized cycle map and the niveau filtration. Teh makes an analogous conjecture for the reduced Lawson groups.

CONJECTURE 1.1 ([Teh10, Conjecture 7.9]). *Let X be a smooth projective real variety. Then $Im(cyc_{q,n}) = N_{2n-q} H_n(X(\mathbb{R}); \mathbb{Z}/2)$ for any $0 \leq q \leq n$.*

Here $N_p H_n(X(\mathbb{R}); \mathbb{Z}/2)$ is the niveau filtration which is the sum over all images

$$Im(H_n(V(\mathbb{R}); \mathbb{Z}/2) \rightarrow H_n(X(\mathbb{R}); \mathbb{Z}/2))$$

such that $\dim V \leq p$.

In the complex case, Friedlander-Mazur’s conjecture is a very difficult and interesting question. It is known to be true with arbitrary finite coefficients as a corollary of the Beilinson-Lichtenbaum conjecture. With integer coefficients, it is known that Suslin’s conjecture for a smooth complex quasi-projective variety X implies Friedlander-Mazur’s conjecture for X . Moreover, Beilinson showed that Grothendieck’s conjecture B for a smooth complex quasi-projective variety X is equivalent to the rational Friedlander-Mazur conjecture for X [Bei10] (see also [Fri95]).

Surprisingly, the real case is totally different.

THEOREM 1.2. *Conjecture 1.1 is false.*

To see the failure of this conjecture we first observe that the niveau filtration on reduced Lawson homology is uninteresting (the case of Borel-Moore homology is $q = 0$). Specifically we have that

$$N_j RL_q H_n(X) = \begin{cases} RL_q H_n(X) & j \geq n \\ 0 & \text{else.} \end{cases}$$

This is a consequence of Corollary 4.4 which asserts that the coniveau spectral sequence for the reduced morphic cohomology collapses. Conjecture 1.1 is therefore equivalent to the surjectivity of the generalized cycle maps. It is known that in general $H_n(X(\mathbb{R}); \mathbb{Z}/2) \neq H_n(X(\mathbb{R}); \mathbb{Z}/2)_{alg}$ although it is difficult to find explicit examples. In Example 5.2 and Proposition 5.3 we give an explicit class of such examples. Our examples are based on a decomposition given in Theorem 3.1 of the reduced cycle spaces of a blow-up with smooth center. This decomposition is a generalization to reduced cycle spaces of the main result of [Sch85].

As another application of this decomposition we give in Corollary 5.5 examples of smooth rational varieties X such that

$$cyc_{1,1} : RL^1H^1(X) \rightarrow H^1(X(\mathbb{R}), \mathbb{Z}/2)$$

is not injective. This is in contrast to the complex case where the group of divisors modulo algebraic equivalence of an irreducible complex variety always injects into the corresponding homology group. This also gives examples of thin divisors which represent non-trivial classes in the reduced Lawson homology group (see Remark 5.6).

The collapse of the coniveau spectral sequence for reduced morphic cohomology is a consequence of local vanishing of motivic cohomology in degrees larger than the weight together with the vanishing theorem proved in [HV12b]. For the purposes of seeing that Conjecture 1.1 is false one does not need the full strength of the collapsing, it suffices to use only the local vanishing of motivic cohomology. However, an interesting consequence of the collapse of this spectral sequence is that we can compute reduced morphic cohomology as the sheaf cohomology

$$H_{Zar}^n(X; \mathcal{R}\mathcal{L}^q\mathcal{H}^0) = RL^qH^n(X).$$

As a consequence, we identify a family of birational invariants given by

$$RL^qH^0(X) = H_{Zar}^0(X; \mathcal{R}\mathcal{L}^q\mathcal{H}^0),$$

for any $q \geq 0$. In case $q = \dim(X)$ we obtain that the number s of connected components of $X(\mathbb{R})$ is a birational invariant (i.e. $RL^{\dim(X)}H^0(X) = (\mathbb{Z}/2)^s$). The purely algebraic nature of s forms part of the main result of [CTP90], where they use étale cohomology. The relation between reduced morphic cohomology and étale cohomology is discussed in the final section. As an application of these birational invariants we compute reduced Lawson homology of a real rational surface in Corollary 3.10.

In Section 6 we discuss cycle maps. We show that there is basically one cycle map from the mod-2 motivic cohomology to mod-2 singular cohomology of the space of real points. As a consequence we see in Theorem 6.9 that the Borel-Haeflinger cycle map factors through the cycle map from reduced morphic cohomology to singular cohomology.

We thank the anonymous referee for some helpful comments.

2. PRELIMINARIES

Let Y be a projective complex variety and $\mathcal{C}_q(Y)$ the Chow variety of effective q -cycles on Y . Write $\mathcal{Z}_q(Y) = (\mathcal{C}_q(Y)(\mathbb{C}))^+$ for the group completion of this monoid. The group completion is done algebraically and $\mathcal{Z}_q(Y)$ is given the quotient topology. It turns out that this naive group completion is actually a homotopy group completion [FG93], [LF92]. When U is quasi-projective with projectivization $U \subseteq \overline{U}$ then define $\mathcal{Z}_q(U) = \mathcal{Z}_q(\overline{U})/\mathcal{Z}_q(U_\infty)$ where $U_\infty = \overline{U} \setminus U$ (and the quotient is a group quotient). This definition is independent of choice of projectivization [LF92], [FG93].

If X is a real variety then $G = \mathbb{Z}/2$ acts on $\mathcal{Z}_q(X_{\mathbb{C}})$ via complex conjugation. The space of real cycles on X is defined to be the subgroup $\mathcal{Z}_q(X_{\mathbb{C}})^G$ of cycles invariant under conjugation. Write $\mathcal{Z}_q(X_{\mathbb{C}})^{av}$ for the subgroup generated by cycles of the form $\alpha + \overline{\alpha}$. The space of reduced cycles on X is defined to be the quotient group

$$\mathcal{R}_q(X) = \frac{\mathcal{Z}_q(X_{\mathbb{C}})^G}{\mathcal{Z}_q(X_{\mathbb{C}})^{av}}.$$

DEFINITION 2.1 ([Teh10]). Let X be a quasi-projective real variety. The *reduced Lawson homology* of X is defined by

$$RL_q H_{q+i}(X) = \pi_i \mathcal{R}_q(X).$$

When $q = 0$ we have that $\mathcal{R}_0(X) = \mathcal{Z}_0(X(\mathbb{R}))/2$ so by the Dold-Thom theorem $RL_0 H_i(X) = H_i(X(\mathbb{R}); \mathbb{Z}/2)$ is the Borel-Moore homology of $X(\mathbb{R})$. In general $RL_q H_{q+i}(X)$ are all $\mathbb{Z}/2$ -vector spaces. It is not known whether these are finitely-generated vector spaces or not however we do have the following vanishing theorem.

THEOREM 2.2 ([HV12b]). *Let X be a quasi-projective real variety. Then*

$$RL_q H_n(X) = 0$$

if $n > \dim(X)$.

There is also a space $\mathcal{R}^q(X) = \mathcal{Z}^q(X_{\mathbb{C}})^G/\mathcal{Z}^q(X_{\mathbb{C}})^{av}$ of reduced algebraic cocycles on X when X is normal and projective. We refer to [Teh10] for the details of its construction. It is convenient to extend this definition to quasi-projective normal varieties, which is done in [HV12b] although not introduced formally as such an extension. We avoid difficulties with point-set topology by giving the extension as a simplicial abelian group. Define the simplicial abelian group of reduced cocycles on a quasi-projective normal real variety

$$\widetilde{\mathcal{R}}^q(X) = \frac{\text{Sing}_\bullet \mathcal{Z}^q/2(X_{\mathbb{C}})^G}{\text{Sing}_\bullet \mathcal{Z}^q/2(X_{\mathbb{C}})^{av}}.$$

If X is a projective, normal real variety then $\widetilde{\mathcal{R}}^q(X) \xrightarrow{\simeq} \text{Sing}_\bullet \mathcal{R}^q(X)$ is a homotopy equivalence [HV12b, Lemma 6.7].

DEFINITION 2.3 ([Teh10]). Let X be a normal quasi-projective real variety. Define the reduced morphic cohomology of X by

$$RL^q H^{q-i}(X) = \pi_i \tilde{\mathcal{R}}^q(X).$$

The reduced Lawson homology and reduced morphic cohomology are related by a Poincaré duality. This is proved in [Teh10, Theorem 6.2] for smooth projective varieties. We give a quick proof below which applies to both the projective and quasi-projective case. If T is a topological abelian group we write $\tilde{T} = \text{Sing}_\bullet T$ for the associated simplicial abelian group. If M is a G -module and σ is the nontrivial element of G we write $N = 1 + \sigma$ and define $M^{av} = \text{Im}(N)$. If in addition M is 2-torsion then we have the two fundamental short exact sequences of abelian groups $0 \rightarrow M^G \rightarrow M \xrightarrow{N} M^{av} \rightarrow 0$ and $0 \rightarrow M^{av} \rightarrow M^G \rightarrow M^G/M^{av} \rightarrow 0$.

THEOREM 2.4. *Let X be a smooth quasi-projective real variety of dimension d . The inclusion*

$$\tilde{\mathcal{R}}^q(X) \rightarrow \tilde{\mathcal{R}}_d(X \times \mathbb{A}^q)$$

is a homotopy equivalence of simplicial abelian groups. Consequently there is a natural isomorphism $RL^q H^n(X) \cong RL_{d-q} H_{d-n}(X)$.

Proof. This follows from consideration of the following comparison diagrams of homotopy fiber sequences of simplicial abelian groups where the right-hand horizontal maps are all surjective,

$$\begin{array}{ccccc} \tilde{\mathcal{Z}}^q/2(X_{\mathbb{C}})^G & \longrightarrow & \tilde{\mathcal{Z}}^q/2(X_{\mathbb{C}}) & \xrightarrow{N} & \tilde{\mathcal{Z}}^q/2(X_{\mathbb{C}})^{av} \\ \downarrow \simeq & & \downarrow \simeq & & \downarrow \\ \tilde{\mathcal{Z}}_d/2(X_{\mathbb{C}} \times_{\mathbb{C}} \mathbb{A}_{\mathbb{C}}^q)^G & \longrightarrow & \tilde{\mathcal{Z}}_d/2(X_{\mathbb{C}} \times_{\mathbb{C}} \mathbb{A}_{\mathbb{C}}^q) & \xrightarrow{N} & \tilde{\mathcal{Z}}_d/2((X_{\mathbb{C}} \times_{\mathbb{C}} \mathbb{A}_{\mathbb{C}}^q)^{av}) \end{array}$$

and

$$\begin{array}{ccccc} \tilde{\mathcal{Z}}^q/2(X_{\mathbb{C}})^{av} & \longrightarrow & \tilde{\mathcal{Z}}^q/2(X_{\mathbb{C}})^G & \longrightarrow & \tilde{\mathcal{R}}^q(X) \\ \downarrow & & \downarrow \simeq & & \downarrow \\ \tilde{\mathcal{Z}}_d/2(X_{\mathbb{C}} \times_{\mathbb{C}} \mathbb{A}_{\mathbb{C}}^q)^{av} & \longrightarrow & \tilde{\mathcal{Z}}_d/2(X_{\mathbb{C}} \times_{\mathbb{C}} \mathbb{A}_{\mathbb{C}}^q)^G & \longrightarrow & \tilde{\mathcal{R}}_d(X \times \mathbb{A}_{\mathbb{R}}^q). \end{array}$$

The displayed homotopy equivalences follow from [Fri98, Theorem 5.2] and [HV12b, Corollary 4.20]

□

The inclusion of algebraic cocycles into topological cocycles defines a generalized cycle map

$$cyc_{q,n} : RL^q H^n(X) \rightarrow H^n(X(\mathbb{R}); \mathbb{Z}/2).$$

If X is smooth and $q \geq \dim X$ then the cycle map $cyc_{q,n}$ is an isomorphism. For X projective this follows from [Teh10, Corollary 6.5, Theorem 8.1] (the results there are stated under the assumption that $X(\mathbb{R})$ is nonempty and connected

but this is unnecessary). The isomorphism for projective varieties implies the isomorphism for quasi-projective varieties (for example by using cohomology with supports and an argument as in [HV12a, Corollary 4.2]).

There are operations called the s -map in both reduced Lawson homology and reduced morphic cohomology $s : RL_q H_n(X) \rightarrow RL_{q-1} H_n(X)$ and $s : RL^q H^n(X) \rightarrow RL^{q+1} H^n(X)$. Iterated compositions of s -maps give rise to generalized cycle maps in reduced Lawson homology

$$cyc_{q,n} : RL_q H_n(X) \xrightarrow{s} RL_{q-1} H_n(X) \xrightarrow{s} \cdots \xrightarrow{s} RL_0 H_n(X) = H_n(X(\mathbb{R}); \mathbb{Z}/2).$$

The s -map in morphic cohomology is compatible with the cycle map in the sense that $cyc_{q,n}$ agrees with the composition

$$cyc_{q,n} : RL^q H^n(X) \xrightarrow{s} RL^{q+1} H^n(X) \xrightarrow{cyc_{q+1,n}} H^n(X(\mathbb{R}); \mathbb{Z}/2).$$

Let X be a normal quasi-projective real variety and $Z \subseteq X$ a closed subvariety. The *reduced morphic cohomology with supports* is defined in the usual way with $RL^q H^{q-i}(X)_Z = \pi_i \tilde{\mathcal{R}}^q(X)_Z$ where $\tilde{\mathcal{R}}^q(X)_Z = \text{hofib}(\tilde{\mathcal{R}}^q(X) \rightarrow \tilde{\mathcal{R}}^q(X - Z))$.

THEOREM 2.5 (Cohomological purity for reduced morphic cohomology). *Let X be a smooth, quasi-projective real variety of dimension d and $Z \subset X$ a closed smooth subvariety of codimension p . There are homotopy equivalences $\tilde{\mathcal{R}}^q(X)_Z \simeq \tilde{\mathcal{R}}^{q-p}(Z)$, which induce natural isomorphisms*

$$RL^q H^n(X)_Z = RL^{q-p} H^{n-p}(Z).$$

Proof. This follows from the localization sequence for reduced Lawson homology [Teh10, Corollary 3.14] together with Poincaré duality, Theorem 2.4, between reduced Lawson homology and reduced morphic cohomology. \square

Recall that a presheaf $F(-)$ of cochain complexes satisfies *Nisnevich descent* provided that for any smooth X , any étale map $f : Y \rightarrow X$, and open embedding $i : U \subseteq X$ such that $f : Y - f^{-1}(U) \rightarrow X - U$ is an isomorphism, we have a Mayer-Vietoris exact triangle (in the derived category of abelian groups):

$$F(X) \rightarrow F(Y) \oplus F(U) \rightarrow F(f^{-1}(V)) \rightarrow F(X)[1].$$

COROLLARY 2.6. *The presheaf $\tilde{\mathcal{R}}^q(-)$ is homotopy invariant theory and satisfies Nisnevich descent.*

Proof. It is homotopy invariant by [Teh10, Theorem 5.13]. Nisnevich descent follows immediately from Theorem 2.5. \square

3. BIRATIONAL INVARIANTS AND EXAMPLES

We use the following decomposition theorem for spaces of reduced cycles of blow-ups with smooth center in order to obtain a decomposition of the cokernels of the cycle map from reduced Lawson homology. Later we use this decomposition to exhibit spaces whose cycle map has nontrivial cokernel. Recall that $\mathcal{R}_{-q}(X) = \mathcal{R}_0(X \times \mathbb{A}^q)$ for $q \geq 0$.

THEOREM 3.1. *Let X be a smooth projective real variety and $Z \subset X$ a smooth closed irreducible subvariety of codimension $d > 1$. Let $\pi : \text{Bl}_Z(X) \rightarrow X$ be the blow up of X with smooth center Z . Then, for any $0 \leq q \leq \dim(X)$ we have a homotopy equivalence of topological abelian groups*

$$(3.2) \quad \mathcal{R}_q(\text{Bl}_Z(X)) \stackrel{h.e.}{\simeq} \mathcal{R}_q(X) \oplus \mathcal{R}_{q-d+1}(Z) \oplus \mathcal{R}_{q-d+2}(Z) \oplus \cdots \oplus \mathcal{R}_{q-1}(Z).$$

Moreover this decomposition is compatible with s -maps.

Proof. We follow the ideas used to prove [Voi, Theorem 2.5] and the main theorem of [Sch85]. We work in $\mathcal{H}^{-1}AbTop$, the category of topological abelian groups with a CW-structure with inverted homotopy equivalences. Recall that $\pi^{-1}(Z) \rightarrow Z$ is the projective bundle $p : \mathbb{P}(N_Z X) \rightarrow Z$ of dimension $d - 1$. The decomposition in the statement of the theorem is given as follows. The first component of the map is π_* (notice that $\pi_* \circ \pi^* = id$). The other maps are given via compositions

$$\phi_l : \mathcal{R}_{q-d+1+l}(Z) \xrightarrow{p^*} \mathcal{R}_{q+l}(\text{Bl}_Z(X)) \xrightarrow{-\cap c_1(O(1))^l} \mathcal{R}_q(\text{Bl}_Z(X)),$$

where $p^* : \mathcal{R}_q(Z) \rightarrow \mathcal{R}_{q+d-1}(\mathbb{P}(N_Z X)) \xrightarrow{i_*} \mathcal{R}_{q+d-1}(\text{Bl}_Z(X))$.

Using the Mayer-Vietoris sequence (see Corollary 2.6), we have the homotopy equivalence

$$\mathcal{R}_k(\text{Bl}_Z(X)) \stackrel{h.e.}{\simeq} \mathcal{R}_k(X) \oplus \text{Ker}(p_*),$$

where $p_* : \mathcal{R}_k(N_Z(X)) \rightarrow \mathcal{R}_k(Z)$. In $\mathcal{H}^{-1}AbTop$

$$(3.3) \quad \text{Ker}(p_*) \stackrel{h.e.}{\simeq} \mathcal{R}_{q-d+1}(Z) \oplus \mathcal{R}_{q-d+2}(Z) \oplus \cdots \oplus \mathcal{R}_{q-1}(Z).$$

The Segre classes $s_l(N_Z X) \cap - : \mathcal{R}_k(Z) \rightarrow \mathcal{R}_{k+d-1-l}(Z)$ satisfy $s_l(N_Z(X)) \cap - = 0$ if $l < 0$ and $s_0(N_Z(X)) \cap - = id$. Using this one can prove the projective bundle formula for the reduced cycle groups in the usual way, see [Teh10]. The projective bundle formula is valid as well for negative indexes. We thus obtain

$$\mathcal{R}_k(N_Z(X)) = \bigoplus_{0 \leq l \leq d-1} \mathcal{R}_{k-d+1+l}(Z).$$

Now one can conclude the homotopy equivalence (3.3).

By [Teh10] the s -maps are compatible with all of the maps involved in the decomposition (3.2) therefore this decomposition is preserved by the s -maps. \square

The generalized cycle maps $cyc_{q,n} : RL_q H_n(X) \rightarrow H_n(X(\mathbb{R}), \mathbb{Z}/2)$, are defined as a composite of s -maps together with the Dold-Thom isomorphism. Write

$$T_{q,n}(X) = \text{coker}(cyc_{q,n} : RL_q H_n(X) \rightarrow H_n(X(\mathbb{R}), \mathbb{Z}/2))$$

and

$$K_{q,n}(X) = \text{ker}(cyc_{q,n} : RL_q H_n(X) \rightarrow H_n(X(\mathbb{R}), \mathbb{Z}/2)).$$

Notice that $T_{q,n}(X) = 0 = K_{q,n}(X)$, for $q \leq 0$ and any X .

COROLLARY 3.4. *Let $\pi : \text{Bl}_Z(X) \rightarrow X$ and Z be as in the above theorem. Then*

$$T_{q,n}(\text{Bl}_Z(X)) = T_{q,n}(X) \oplus T_{q-1,n-1}(Z) \oplus \dots \oplus T_{q-d+1,n-d+1}(Z),$$

and

$$(3.5) \quad K_{q,n}(\text{Bl}_Z(X)) = K_{q,n}(X) \oplus K_{q-1,n-1}(Z) \oplus \dots \oplus K_{q-d+1,n-d+1}(Z).$$

Proof. In the case $k = 0$ the decomposition (3.2) gives

$$\mathcal{R}_0(\text{Bl}_Z(X)(\mathbb{R})) \stackrel{h.e.}{\simeq} \mathcal{R}_0(X(\mathbb{R})) \oplus \mathcal{R}_{-d+1}(Z(\mathbb{R})) \oplus \mathcal{R}_{-d+2}(Z(\mathbb{R})) \oplus \dots \oplus \mathcal{R}_{-1}(Z(\mathbb{R})),$$

therefore producing the decomposition of Borel-Moore homology

$$\begin{aligned} H_k(\text{Bl}_Z(X)(\mathbb{R}), \mathbb{Z}/2) = \\ H_k(X(\mathbb{R}), \mathbb{Z}/2) \oplus H_{k-d-1}(Z(\mathbb{R}), \mathbb{Z}/2) \oplus \dots \oplus H_{k-1}(Z(\mathbb{R}), \mathbb{Z}/2). \end{aligned}$$

The s -maps respect the decomposition (3.2) and comparing the decomposition for $q = 0$ and for $q > 0$ yields the result. \square

COROLLARY 3.6. *If $Z(\mathbb{R}) = \emptyset$, then $T_{q,n}(\text{Bl}_Z(X)) = T_{q,n}(X)$ for any $0 \leq q \leq n \leq \dim(X)$.*

REMARK 3.7. An analog of Corollary 3.4 for the cokernel of the Borel-Haefliger cycle map was originally proven by Schülting in [Sch85]. There separate arguments are needed to give a decomposition algebraically and a decomposition topologically. An advantage that our uniform proof has is that is entirely algebraic, the homology of real points being expressed in terms of homotopy of the group of algebraic cycles $\mathcal{R}_0(X)$.

REMARK 3.8. Using similar techniques one can prove a decomposition analogous to (3.2) for the spaces of real cycles defining dos Santos equivariant Lawson homology groups.

COROLLARY 3.9. *The groups $T_{1,n}(X)$ and $K_{1,n}(X)$ are birational invariants for smooth projective real varieties.*

Proof. By [AKMW02, Theorem 0.3.1] every birational map between smooth projective varieties factors as a composition of blow-ups and blow-downs with smooth centers. The result then follows from Corollary 3.4. \square

We close the section with the following computation.

COROLLARY 3.10. *Let X be a rational smooth projective surface i.e. $X \stackrel{\text{birational}}{\simeq} \mathbb{P}_{\mathbb{R}}^2$. Then the cycle map*

$$\text{cyc}_{q,n} : RL_q H_n(X) \rightarrow H_n(X(\mathbb{R}), \mathbb{Z}/2)$$

is an isomorphism for $q \leq n$.

Proof. We have $\mathcal{R}_0(X) = \mathcal{R}_0(X(\mathbb{R}))$. By Corollary 3.9 we have that $T_{1,n}(X)$ and $K_{1,n}(X)$ are birational invariants and we know that $T_{1,n}(\mathbb{P}_{\mathbb{R}}^2) = 0 = K_{1,n}(\mathbb{P}_{\mathbb{R}}^2)$. By Theorem 4.4 the group $\pi_0 \mathcal{R}_2(X)$ is a birational invariant. \square

REMARK 3.11. For X as in the previous corollary we have that $RL_q H_n(X) = 0$ for $n < q$ and $n > 2$ and $RL_0 H_0(X) = RL_0 H_2(X) = RL_1 H_2(X) = RL_2 H_2(X) = \mathbb{Z}/2$.

4. CONIVEAU SPECTRAL SEQUENCES

In this section we show that the coniveau spectral sequence for reduced morphic cohomology collapses. We make use of [CTHK97] for the Bloch-Ogus theorem identifying the E_2 -term of this spectral sequence. Let X be a smooth, quasi-projective real variety and write $X^{(p)}$ for the set of points $x \in X$ whose closure has codimension p . Let h^* be a cohomology theory with supports. Define $h_x^i(X) = \text{colim}_{U \subseteq X} h_{\bar{x} \cap U}^i(U)$ and $h^i(k(x)) = \text{colim}_{U \subseteq \bar{x}} h^i(U)$ (where in both colimits, U ranges over nonempty opens). One may form the Gersten complex

$$0 \rightarrow \bigoplus_{x \in X^{(0)}} h_x^n(X) \rightarrow \bigoplus_{x \in X^{(1)}} h_x^{n+1}(X) \rightarrow \dots \rightarrow \bigoplus_{x \in X^{(p)}} h_x^{n+p}(X) \rightarrow \dots$$

This complex gives rise to the coniveau spectral sequence

$$E_1^{p,q} = \bigoplus_{x \in X^{(p)}} h_x^{p+q}(X) \implies h^{p+q}(X)$$

The associated filtration is $N^p h^n(X) = \cup_Z \text{Im}(h_Z^n(X) \rightarrow h(X))$, where the union is over closed subvarieties $Z \subseteq X$ of codimension p .

PROPOSITION 4.1. [CTHK97, Corollary 5.1.11, Theorem 8.5.1] *Let h^* be a cohomology theory with supports on Sm/\mathbb{R} which satisfies Nisnevich excision and is homotopy invariant. Let \mathcal{H}^q be the Zariski sheafification of the presheaf $U \mapsto h^q(U)$. Then the Gersten complex $E_1^{\bullet,q}$ is a flasque resolution of \mathcal{H}^q and the coniveau spectral sequence has the form*

$$E_2^{p,q} = H_{Zar}^p(X; \mathcal{H}^q) \implies H^{p+q}(X).$$

For every q , the group $H^0(X, \mathcal{H}^q)$ is a birational invariant for smooth proper varieties.

COROLLARY 4.2. *Let X be a smooth quasi-projective real variety. For each k we have spectral sequences*

$$E_1^{p,q}(k) = \bigoplus_{x \in X^{(p)}} RL^{k-p} H^q(k(x)) \implies RL^k H^{p+q}(X).$$

The E_2 -terms are $E_2^{p,q}(k) = H_{Zar}^p(X; \mathcal{R}\mathcal{L}^k \mathcal{H}^q)$ and each $H_{Zar}^0(X; \mathcal{R}\mathcal{L}^k \mathcal{H}^q)$ is a birational invariant for smooth projective real varieties. Moreover, the s -maps induce maps of spectral sequences $\{E_r^{p,q}(k)\} \rightarrow \{E_r^{p,q}(k+1)\}$.

Proof. By Corollary 2.6 reduced morphic cohomology is homotopy invariant and satisfies Nisnevich excision. For $x \in X^{(p)}$ we have that $\bar{x} \cap U$ is smooth for small enough open $U \subseteq X$. Therefore, for small enough open $U \subseteq X$ we can apply Theorem 2.5 to $\bar{x} \cap U \subseteq U$ and we conclude that we have an isomorphism $RL^k H^{p+q}(X)_x = RL^{k-p} H^q(k(x))$. Thus the E_1 -page of the coniveau spectral sequence can be rewritten in the displayed form. The s -maps are natural

transformations and so induce maps of the exact couples defining the coniveau spectral sequence. \square

The following result gives us the collapsing of the coniveau spectral sequence. It is a consequence of the main vanishing theorem of [HV12b] and the local vanishing of equivariant morphic cohomology and morphic cohomology.

THEOREM 4.3. *Let $R = \mathcal{O}_{T, t_1, \dots, t_n}$ be the semi-local ring of a smooth, real variety T at a finite set of points $t_1, \dots, t_n \in T$. Then*

$$RL^k H^q(\mathrm{Spec} R) = 0$$

if $q \neq 0$ and any $k \geq 0$.

Proof. For convenience write $Y = \mathrm{Spec} R$. By definition

$$RL^k H^q(Y) = \mathrm{colim} RL^k H^q(U),$$

where the colimit is over all open $U \subseteq T$ such that all $t_i \in U$. Recall also that filtered colimits commute with homotopy groups and preserve exact sequences. We need to see that $RL^k H^{k-s}(Y) = \pi_s \mathcal{R}^k(Y) = 0$ for $s \neq k$. The main vanishing result in [HV12b, Theorem 6.10] implies that $\pi_s \mathcal{R}^k(Y) = 0$ for $s > k$. Consider the homotopy fiber sequences of simplicial abelian groups

$$\tilde{\mathcal{Z}}^k/2(Y_{\mathbb{C}})^G \longrightarrow \tilde{\mathcal{Z}}^k/2(Y_{\mathbb{C}}) \xrightarrow{N} \tilde{\mathcal{Z}}^k/2(Y_{\mathbb{C}})^{av}$$

and

$$\tilde{\mathcal{Z}}^k/2(Y_{\mathbb{C}})^{av} \longrightarrow \tilde{\mathcal{Z}}^k/2(Y_{\mathbb{C}})^G \longrightarrow \tilde{\mathcal{R}}^k(Y).$$

Because $\pi_s \tilde{\mathcal{Z}}^k/2(Y_{\mathbb{C}})^G = 0 = \pi_s \tilde{\mathcal{Z}}^k/2(Y_{\mathbb{C}})$ if $s \leq k-1$ ([FHW04, Theorem 7.3] and [HV12a, Lemma 3.22]) we see that $\pi_s \tilde{\mathcal{Z}}^k/2(Y_{\mathbb{C}})^{av} = 0$ if $s \leq k-1$. Using the second homotopy fiber sequence we conclude that $\pi_s \tilde{\mathcal{R}}^k(Y) = 0$ if $s \leq k-1$. \square

COROLLARY 4.4. *Let X be a smooth quasi-projective real variety. For any k , the coniveau spectral sequence for reduced morphic cohomology satisfies*

$$E_1^{p,q}(k) = 0$$

for $q \neq 0$. Consequently $E_2^{p,0}(k) = E_{\infty}^{p,0}(k)$ and so we have natural isomorphisms

$$H_{Zar}^p(X; \mathcal{R}\mathcal{L}^k H^0) = RL^k H^p(X).$$

In particular $H^0(X, \mathcal{R}\mathcal{L}^q H^0) = RL^q H^0(X) = \pi_q(\mathcal{R}^q(X))$ is a birational invariant for smooth projective real varieties.

Proof. We have $E_1^{p,q}(k) = \bigoplus_{x \in X^{(p)}} RL^{k-p} H^q(k(x))$. For $x \in X^{(p)}$ we have that $k(x) = \mathcal{O}_{U, \eta}$, where U is the open set of nonsingular points of \bar{x} and η is the generic point. In particular the previous result implies that $RL^{k-p} H^q(k(x))$ for $q \neq 0$. The other statements follow from Proposition 4.1 \square

REMARK 4.5. The case $k = \dim X$ tells us that $\mathcal{H}_{\mathbb{R}}^i = 0$ for any $i > 0$ where $\mathcal{H}_{\mathbb{R}}^i$ is the Zariski sheafification of the presheaf $U \mapsto H^i(U(\mathbb{R}); \mathbb{Z}/2)$. This also follows from [Sch94, Theorem 19.2].

REMARK 4.6. Corollary 4.4 gives us birational invariants $RL^q H^0(X)$ for $0 \leq q \leq d = \dim(X)$. If $q = d$ then we have

$$RL^d H^0(X) = H^0(X(\mathbb{R}); \mathbb{Z}/2) = (\mathbb{Z}/2)^{\oplus s}$$

and therefore $s = s(X) = \#(\text{connected components of } X(\mathbb{R}))$ is a birational invariant of an algebraic nature. This also follows from the main result of [CTP90] where they show that $H^0(X, \mathcal{H}_{et}^n) = H^0(X(\mathbb{R}), \mathbb{Z}/2)$ for any $n \geq \dim(X) + 1$. Here \mathcal{H}_{et} is the sheaf associated to the presheaf $U \rightarrow H_{et}^n(U, \mu_2^{\otimes n})$. At the other extreme if one takes $q = 0$,

$$RL^0 H^0(X) = (\mathbb{Z}/2)^{\oplus r}$$

where $r = r(X) = \#(\text{geometrically irreducible components of } X)$ and so r is also a birational invariant.

REMARK 4.7. By Corollary 4.4 and Corollary 4.2 the s -maps

$$RL^q H^n(X) \xrightarrow{s} RL^{q+1} H^n(X)$$

are obtained as the map induced on Zariski sheaf cohomology by the sheafified s -maps $s : \mathcal{R}\mathcal{L}^q \mathcal{H}^0 \rightarrow \mathcal{R}\mathcal{L}^{q+1} \mathcal{H}^0$. In particular we see that the generalized cycle map

$$cyc_{q,n} : RL^q H^n(X) \rightarrow H^n(X(\mathbb{R}); \mathbb{Z}/2)$$

is obtained from the sheafified cycle map $\mathcal{R}\mathcal{L}^q \mathcal{H}^0 \rightarrow \mathcal{H}_{\mathbb{R}}^0$. In the last section we show that this cycle map is naturally related to the Borel-Haeflinger cycle map [BH61].

We finish by observing that Poincare duality gives the collapsing of the niveau spectral sequence for reduced Lawson homology (of possibly singular varieties).

PROPOSITION 4.8. *Let X be a quasi-projective real variety. Write $X_{(p)}$ for the set of points $x \in X$ whose closure has dimension p . The niveau spectral sequence*

$$E_{p,q}^1(k) = \bigoplus_{x \in X_{(p)}} RL_k H_{p+q}(k(x)) \implies RL_k H_{p+q}(X)$$

satisfies $E_{p,q}^1(k) = 0$ for any $q \neq 0$ and therefore $E_{p,q}^2(k) = E_{p,q}^\infty(k)$.

Proof. The niveau spectral sequence is constructed as in [BO74]. Consider an $x \in X_{(p)}$. For any open $U \subseteq \bar{x}$ which is smooth, by Theorem 2.4 we have $RL_k H_n(U) = RL^{p-k} H^{p-n}(U)$. In particular, we see that $RL_k H_n(k(x)) = RL^{p-k} H^{p-n}(k(x))$. Therefore by Corollary 4.4 we have $RL_k H_{p+q}(k(x)) = RL^{p-k} H^{-q}(k(x)) = 0$ for any $q \neq 0$. \square

5. FILTRATIONS IN HOMOLOGY

Let X be a quasi-projective real variety of dimension d . The generalized cycle map $\phi_{q,n} : RL_q H_n(X) \rightarrow H_n(X(\mathbb{R}), \mathbb{Z}/2)$ is the composition of q iterations of the s -map together with the Dold-Thom isomorphism. Write

$$RT_q H_n(X) = \text{Im}(\phi_{q,n} : RL_q H_n(X) \rightarrow H_n(X(\mathbb{R}); \mathbb{Z}/2).$$

This gives us a decreasing filtration of the homology of the space of real points and is called the topological filtration.

Associated to the niveau spectral sequence is the niveau filtration

$$N_p RL_k H_n(X) = \sum_{\dim V \leq p} \text{Im}(RL_k H_n(V) \rightarrow RL_k H_n(X)).$$

Notice that in the complex case, if Y is a complex variety of dimension d then Weak Lefschetz theorem says that

$$N_n H_n(Y(\mathbb{C}); \mathbb{Z}) = N_{n+1} H_n(Y(\mathbb{C}); \mathbb{Z}) = \cdots = N_d H_n(Y(\mathbb{C}); \mathbb{Z}).$$

In particular, in the complex case there are only $n + 1$ steps in the filtration, and another $d - n$ are equal to the homology. In the real case, we don't have this theorem and so a priori all one has is a filtration.

$$N_0 H_n(X(\mathbb{R}); \mathbb{Z}/2) \subseteq \cdots \subseteq N_d H_n(X(\mathbb{R}); \mathbb{Z}/2) = H_n(X(\mathbb{R}); \mathbb{Z}/2).$$

Teh has formulated the following conjecture which is made in analogy with a conjecture of Friedlander-Mazur [FM94, Conjecture p.71] for complex varieties.

CONJECTURE 5.1 ([Teh10, Conjecture 7.9]). *Let X be a smooth projective real variety. Then $RT_q H_n(X) \subseteq N_{2n-q} H_n(X(\mathbb{R}); \mathbb{Z}/2)$ and moreover this containment is an equality $RT_q H_n(X) = N_{2n-q} H_n(X(\mathbb{R}); \mathbb{Z}/2)$ for any $0 \leq q \leq n$.*

From Proposition 4.8 we have the following equality:

$$E_{n-q,q}^\infty(k) = N_{n-q} RL_k H_n(X) / N_{n-q-1} RL_k H_n(X) = 0$$

for any $q \neq 0$. This means that for any k we have

$$RL_k H_n(X) = N_d RL_k H_n(X) = \cdots = N_{n+1} RL_k H_n(X) = N_n RL_k H_n(X),$$

and

$$0 = N_{-1} RL_k H_n(X) = N_0 RL_k H_n(X) = \cdots = N_{n-1} RL_k H_n(X).$$

The first row of equalities contains the groups that appear in Conjecture 5.1. Consequently the first part of the conjecture is obviously true because by the above we have that $N_j H_n(X(\mathbb{R}); \mathbb{Z}/2) = H_n(X(\mathbb{R}); \mathbb{Z}/2)$ for all $j \geq n$.

The second part of the conjecture is false because the s -maps are not always surjective. Using the material from Section 3 we give an explicit example of this failure. Recall that we write $T_{q,n}(X) = \text{coker}(cyc_{q,n} : RL_q H_n(X) \rightarrow H_n(X(\mathbb{R}); \mathbb{Z}/2))$.

EXAMPLE 5.2. Let $Z \subseteq \mathbb{P}_{\mathbb{R}}^3$ be the smooth irreducible elliptic curve given by the equation $t^2x + ty^2 - x^3 = 0, z = 0$. Then $Z(\mathbb{R})$ is well known to consist of 2 connected components (see for example [BCR98, Example 3.1.2]). Let $X = \text{Bl}_Z \mathbb{P}^3$ be the blow-up of \mathbb{P}^3 along Z . Then $T_{2,2}(X) = T_{2,2}(\mathbb{P}_{\mathbb{R}}^3) \oplus T_{1,1}(Z)$ by Corollary 3.4. Because $RL_1H_1(Z) = \pi_0\mathcal{R}_1(Z) = \mathbb{Z}/2$ and $Z(\mathbb{R})$ has two components we conclude that $T_{2,2}(X) = \mathbb{Z}/2$.

More generally we have the following.

PROPOSITION 5.3. *For each $N \geq 3$, there is a smooth projective real variety X of dimension N (which is topologically connected) such that $T_{q,q}(X) \neq 0$ for all $2 \leq q \leq N - 1$.*

Proof. Let $Z \subseteq \mathbb{P}^N$ be a smooth irreducible real curve such that $Z(\mathbb{R})$ has at least 2 connected components. Let s denote the number of connected components of $Z(\mathbb{R})$. Since Z is irreducible we have that $\mathcal{R}_1(Z) = \mathbb{Z}/2$. Therefore $T_{1,1}(Z) = (\mathbb{Z}/2)^{s-1}$. Since Z is a curve $T_{i,n}(Z) = 0$ for all other values of i and n . We take $X \rightarrow \mathbb{P}^N$ to be the blow up of \mathbb{P}^N along Z . Then $T_{q,q}(X) = T_{1,1}(Z) = \mathbb{Z}/2^{s-1}$ by Corollary 3.4 because $2 \leq q \leq N - 1 = \text{codim}(Z)$. \square

We also have a similar result for the kernel.

PROPOSITION 5.4. *For each $N \geq 3$, there is a smooth projective real variety X , birational to \mathbb{P}^N , such that $K_{q,q}(X) \neq 0$ for all $2 \leq q \leq N - 1$.*

Proof. Let $Z \subseteq \mathbb{P}^N$ be a smooth irreducible real curve such that $Z(\mathbb{R}) = \emptyset$. Then $K_{1,1}(Z) = \mathbb{Z}/2$ and $K_{i,n}(Z) = 0$ for all other values of i and n . Take $X \rightarrow \mathbb{P}^N$ to be the blow up of \mathbb{P}^N along Z . We have $K_{q,q}(X) = K_{1,1}(Z) = \mathbb{Z}/2$ by Corollary 3.4. \square

As an interesting particular case we have the following which is different than the complex analog.

COROLLARY 5.5. *There exists a smooth real variety $X \stackrel{\text{birational}}{\sim} \mathbb{P}_{\mathbb{R}}^N$ such that the cycle map on divisors $RL^1H^1(X) \rightarrow H^1(X(\mathbb{R}), \mathbb{Z}/2)$ is not injective.*

REMARK 5.6. A k -cycle is said to be thin if it is a sum of closed subvarieties $Z \subseteq X$ with $\dim Z(\mathbb{R}) < k$. The kernel of the Borel-Haeflinger cycle map consists entirely of thin cycles by [IS88] and the composite $CH_q(X) \rightarrow RL_qH_q(X) \rightarrow H_q(X(\mathbb{R}); \mathbb{Z}/2)$ agrees with the Borel-Haeflinger cycle map by Theorem 6.9. This means that the proposition above gives examples of nonzero classes which are represented by thin cycles in $RL_qH_q(X)$.

6. CYCLE MAPS

Let X be a smooth quasi-projective real variety. We discuss two natural cycle maps from motivic cohomology to the singular cohomology of $X(\mathbb{R})$. Based on this, we show that Borel-Haeflinger map [BH61] factors through the reduced Lawson homology cycle map. We end the section with a discussion of the maps involved in the Suslin conjecture from the view of the methods in this section.

Recall that $G = \mathbb{Z}/2$. If M is a G -space we write $H_G^i(M; \mathbb{Z}/2)$ for the Borel cohomology with $\mathbb{Z}/2$ -coefficients. The reduced morphic cohomology of X comes equipped with a natural generalized cycle map to the singular cohomology of real points. Composing this with the canonical map from real morphic cohomology and its isomorphism with motivic cohomology (with $\mathbb{Z}/2$ -coefficients) gives us the cycle map

$$(6.1) \quad H_{\mathcal{M}}^p(X; \mathbb{Z}/2(q)) \rightarrow RL^q H^{p-q}(X) \xrightarrow{cyc} H^{p-q}(X(\mathbb{R}); \mathbb{Z}/2).$$

On the other hand the real morphic cohomology maps naturally to the Borel cohomology of the space of complex points. In turn there is a map $H_G^n(X(\mathbb{C}); \mathbb{Z}/2) \rightarrow \oplus H^{n-i}(X(\mathbb{R}); \mathbb{Z}/2)$ obtained by restricting to real points together with the decomposition

$$H_G^n(X(\mathbb{R}), \mathbb{Z}/2) = H^n(X(\mathbb{R}) \times \mathbb{R}P^\infty; \mathbb{Z}/2) = \bigoplus_{0 \leq i \leq n} H^i(X(\mathbb{R}), \mathbb{Z}/2).$$

Composing with the appropriate projection gives us

$$(6.2) \quad H_{\mathcal{M}}^p(X; \mathbb{Z}/2(q)) \rightarrow H_G^p(X(\mathbb{C}); \mathbb{Z}/2) \rightarrow H^{p-q}(X(\mathbb{R}); \mathbb{Z}/2).$$

We show that the cycle maps (6.1) and (6.2) agree with each other. Basically these agree because they can be seen as induced by maps of presheaves of cochain complexes and so Theorem 6.6 applies.

Write $(Top)_{an}$ for the category of topological spaces homeomorphic to a finite dimensional CW-complex given the usual topology and $\phi : (Top)_{an} \rightarrow (Sm/\mathbb{R})_{zar}$ for the map of sites induced by $X \mapsto X(\mathbb{R})$.

THEOREM 6.3. *Let X be a smooth quasi-projective real variety. The cycle maps given by (6.1) and (6.2) agree. Moreover the intermediate maps in (6.2) can be chosen so that the following diagram commutes*

$$\begin{array}{ccccc} H_{\mathcal{M}}^p(X, \mathbb{Z}/2(q)) & \longrightarrow & H_{et}^p(X, \mu_2^{\otimes q}) & \xrightarrow{\cong} & H_G^p(X(\mathbb{C}), \mathbb{Z}/2) \\ \downarrow & & \downarrow & & \downarrow \\ RL^p H^{p-q}(X) & \xrightarrow{cyc} & H^{p-q}(X(\mathbb{R}), \mathbb{Z}/2) & \longleftarrow & H_G^p(X(\mathbb{R}), \mathbb{Z}/2). \end{array}$$

for any $p, q \geq 0$.

Proof. Consider the following complexes of Zariski sheaves on Sm/\mathbb{R} :

$$\begin{aligned} \mathbb{Z}/2(q)(X) &= (z_{equi}(\mathbb{P}_{\mathbb{R}}^{q/q-1}, 0)(X \times_{\mathbb{R}} \Delta_{\mathbb{R}}^{\bullet}) \otimes \mathbb{Z}/2)[-2q] \\ \mathbb{Z}/2(q)^{sst}(X) &= \text{Sing}_{\bullet}(\mathbb{Z}^q/2(X_{\mathbb{C}})^G)[-2q] \\ \mathbb{Z}/2(q)^{top}(X) &= \text{Hom}_{cts}(X(\mathbb{C}) \times \Delta_{top}^{\bullet}, \mathbb{Z}/2_0(\mathbb{A}_{\mathbb{C}}^q))^G[-2q] \\ \mathbb{Z}/2(q)^{Bor}(X) &= \text{Hom}_{cts}(X(\mathbb{C}) \times EG \times \Delta_{top}^{\bullet}, \mathbb{Z}/2_0(\mathbb{A}_{\mathbb{C}}^q))^G[-2q] \\ \mathbb{Z}/2(q)_{\mathbb{R}}^{Bor}(X) &= \text{Hom}_{cts}(X(\mathbb{R}) \times EG \times \Delta_{top}^{\bullet}, \mathbb{Z}/2_0(\mathbb{A}_{\mathbb{C}}^q))^G[-2q], \\ \mathcal{R}(q)(X) &= (\text{Sing}_{\bullet} \mathbb{Z}^q/2(X_{\mathbb{C}})^G / \text{Sing}_{\bullet} \mathbb{Z}^q/2(X_{\mathbb{C}})^{av})[-2q] \end{aligned}$$

These complexes all satisfy Nisnevich descent for standard reasons. See e.g. [HV12b, Section 5] for the first three. Similarly for the complex $\mathbb{Z}/2(q)_{\mathbb{R}}^{Bor}(X)$ because taking real points of a distinguished Nisnevich square of real varieties gives a homotopy pushout square of spaces. The last complex satisfies Nisnevich descent by Proposition 2.6. As a consequence we have

$$\begin{aligned} \mathbb{H}_{Zar}^i(Z; \mathbb{Z}/2(q)) &= H_{\mathcal{M}}^i(X; \mathbb{Z}/2(q)) \\ \mathbb{H}_{Zar}^i(X; \mathbb{Z}/2(q)^{sst}) &= L^q H\mathbb{R}^{i-q,q}(X; \mathbb{Z}/2) \\ \mathbb{H}_{Zar}^i(X; \mathbb{Z}/2(q)^{top}) &= H^{i-q,q}(X(\mathbb{C}); \underline{\mathbb{Z}/2}) \\ \mathbb{H}_{Zar}^i(X; \mathbb{Z}/2(q)^{Bor}) &= H_G^i(X(\mathbb{C}); \mathbb{Z}/2) \\ \mathbb{H}_{Zar}^i(X; \mathbb{Z}/2(q)_{\mathbb{R}}^{Bor}) &= H_G^i(X(\mathbb{R}); \mathbb{Z}/2) \\ \mathbb{H}_{Zar}^i(X; \mathcal{R}(q)) &= RL^q H^{i-q}(X), \end{aligned}$$

where $L^q H\mathbb{R}^{i-q,q}(X; \mathbb{Z}/2)$ denotes Friedlander-Walker’s real morphic cohomology [FW02] and $H^{i-q,q}(X(\mathbb{C}); \underline{\mathbb{Z}/2})$ is Bredon cohomology.

The map (6.1) is induced by the map of complexes

$$(6.4) \quad \mathbb{Z}/2(q) \xrightarrow{1)} \mathcal{R}(q) \xrightarrow{2)} \phi_* \mathbb{Z}/2[-q].$$

The map 1) is given by the “usual” cycle map from motivic cohomology to reduced morphic cohomology. It is defined as the composite $\mathbb{Z}/2(q) \rightarrow \mathbb{Z}/2(q)^{sst} \rightarrow \mathcal{R}(q)$. The map 2) is obtained by adjunction from the composite

$$\phi^*(\mathcal{R}(q)) \xrightarrow{\simeq} Map_{cts}((-)(\mathbb{R}) \times \Delta_{top}^*, \mathcal{R}_0(\mathbb{A}_{\mathbb{R}}^q))[-2q] \xrightarrow{\simeq} \mathbb{Z}/2[-q]$$

which arises because $\mathcal{R}_0(\mathbb{A}_{\mathbb{R}}^q) \simeq K(\mathbb{Z}/2, q)$ and any CW complex has an open cover given by contractibles.

We show that the map (6.2) is induced by a composite of maps:

$$(6.5) \quad \mathbb{Z}/2(q) \rightarrow tr_{\leq 2q} \mathbb{R}\epsilon_* \mathbb{Z}/2 \xrightarrow{3)} \mathbb{Z}/2(q)^{Bor} \xrightarrow{4)} \mathbb{Z}/2(q)_{\mathbb{R}}^{Bor} \xrightarrow{5)} \phi_* \mathbb{Z}/2[-q].$$

of Zariski complexes of sheaves in $D^-(Shv_{Zar}(Sm/\mathbb{R}))$ which we now explain. Write $\epsilon : X_{et} \rightarrow X_{Zar}$ for the usual map of sites. The first unlabeled map is the cycle map from motivic cohomology to etale cohomology. The map 3) is obtained in Proposition 6.7 using Cox’s theorem [Cox79]. The map 4) is obtained by restriction to real points. The map 5) will be obtained from the adjoint of a map $\phi^*(\mathbb{Z}/2(q)_{\mathbb{R}}^{Bor}) \rightarrow \mathbb{Z}/2[-q]$ as follows. Every CW-complex is locally contractible and so

$$\phi^*(\mathbb{Z}/2(q)_{\mathbb{R}}^{Bor}) \simeq Hom_{cts}(EG \times \Delta_{top}^\bullet, \mathcal{Z}/2_0(\mathbb{A}_{\mathbb{C}}^q))^G[-2q]$$

is a quasi-isomorphism of complexes of Zariski sheaves where the right-hand side is the constant sheaf. In $D_{\mathbb{Z}/2}^-(Ab)$, any complex is quasi-isomorphic with the complex given by its cohomology. We have that $H^{p,q}(EG; \underline{\mathbb{Z}/2}) = H^{p+q}(BG; \mathbb{Z}/2)$ and therefore

$$H^k Hom_{cts}(EG \times \Delta_{top}^\bullet, \mathcal{Z}/2_0(\mathbb{A}_{\mathbb{C}}^q))^G[-2q] = H^{q-k,q}(EG; \underline{\mathbb{Z}/2}) = \mathbb{Z}/2$$

for $0 \leq k \leq 2q$ and is 0 otherwise. This gives us the map

$$\phi^*(\mathbb{Z}/2(q)_{\mathbb{R}}^{Bor}) \simeq \bigoplus_{0 \leq i \leq 2q} \mathbb{Z}/2[-i] \rightarrow \mathbb{Z}/2[-i],$$

which induces the i^{th} projection on cohomology

$$H_G^n(X(\mathbb{R}), \mathbb{Z}/2) = \bigoplus_{0 \leq i \leq n} H^{n-i}(X(\mathbb{R}), \mathbb{Z}/2) \rightarrow H^{n-i}(X(\mathbb{R}), \mathbb{Z}/2).$$

In particular the adjoint of this map for $i = q$ gives us the map 5)

$$\mathbb{Z}/2(q)_{\mathbb{R}}^{Bor} \rightarrow \phi_* \mathbb{Z}/2[-q].$$

Because both map 6.5 and map 6.4 induce non-trivial maps in cohomology they have to coincide by Theorem 6.6. \square

In the proof of the previous theorem we made use of the following result.

THEOREM 6.6. *Let $k = \mathbb{R}$ or \mathbb{C} and write $\phi : (Top)_{an} \rightarrow (Sm/k)_{zar}$ the map of sites that send $X \mapsto X(k)$ where (Top) is the category of spaces homeomorphic to finite dimensional CW complexes equipped with the usual topology. Then we have*

- (1) $\text{Hom}_{D^-((Sm/\mathbb{R})_{zar})}(\mathbb{Z}/2(q), \phi_* \mathbb{Z}/2[-q]) = \mathbb{Z}/2,$
- (2) $\text{Hom}_{D^-((Sm/\mathbb{R})_{zar})}(\mathcal{R}(q), \phi_* \mathbb{Z}/2[-q]) = \mathbb{Z}/2,$
- (3) $\text{Hom}_{D^-((Sm/\mathbb{C})_{zar})}(\mathbb{Z}(q)^{sst}, \mathbb{R}\phi_* \mathbb{Z}/n) = \mathbb{Z}/n,$ for any $n \geq 1.$

Proof. We have a quasi-isomorphism $\mathbb{Z}/2(q) \simeq \mathbb{Z}/2(q)^{sst}$ and that

$$\text{Hom}(\mathbb{Z}/2(q)^{sst}, \phi_* \mathbb{Z}/2[-q]) = \text{Hom}(\phi^*(\mathbb{Z}/2(q)^{sst}), \mathbb{Z}/2[-q]).$$

Because every CW complex is locally contractible in $D^-(Top)$ we have

$$\phi^*(\mathbb{Z}/2(q)^{sst})[2q] \simeq \text{Hom}_{cts}(- \times \Delta_{top}^\bullet, \mathcal{Z}/2_0(\mathbb{A}_{\mathbb{C}}^q)^G) \simeq \text{Sing}_\bullet \mathcal{Z}/2_0(\mathbb{A}_{\mathbb{C}}^q)^G.$$

From [dS03, (3.6)] it follows that $\mathcal{Z}/2_0(\mathbb{A}_{\mathbb{C}}^q)^G \simeq \prod_{i=q}^{2q} K(\mathbb{Z}/2, i)$. This yields the result because we then have

$$\text{Hom}_{D^-(Top)}(\phi^*(\mathbb{Z}/2(q)), \mathbb{Z}/2[-q]) = \bigoplus_{i=q}^{2q} H^q(K(\mathbb{Z}/2, i); \mathbb{Z}/2) = \mathbb{Z}/2.$$

In the proof of Theorem 6.3 we observed that $\phi^* \mathcal{R}(q) \simeq \mathbb{Z}/2[-q]$ and so

$$\begin{aligned} & \text{Hom}_{D^-((Sm/\mathbb{R})_{zar})}(\mathcal{R}(q), \phi_* \mathbb{Z}/2[-q]) \\ &= \text{Hom}_{D^-((Sm/\mathbb{R})_{zar})}(\phi^* \mathcal{R}(q), \mathbb{Z}/2[-q]) \\ &= \text{Hom}_{D^-((Sm/\mathbb{R})_{zar})}(\mathbb{Z}/2[-q], \mathbb{Z}/2[-q]) = \mathbb{Z}/2. \end{aligned}$$

The last item follows from the equivalence $\phi^* \mathbb{Z}(q)^{sst} \simeq \mathbb{Z}$. We have

$$\text{Hom}_{D^-((Sm/\mathbb{C})_{zar})}(\mathbb{Z}^{sst}(q), \mathbb{R}\phi_* \mathbb{Z}/n) = \text{Hom}_{D^-((Sm/\mathbb{R})_{zar})}(\mathbb{Z}, \mathbb{Z}/n) = \mathbb{Z}/n,$$

for any $n \geq 1$. \square

In the proof of Theorem 6 we also used the following proposition which relies on Cox’s theorem identifying the étale cohomology of a real variety with Borel cohomology.

PROPOSITION 6.7. *There is a quasi-isomorphism $\rho : tr_{\leq 2q} \mathbb{R}\epsilon_* \mathbb{Z}/2 \rightarrow \mathbb{Z}/2(q)^{Bor}$ of complexes of Zariski sheaves.*

Proof. We show that the canonical map $\mathbb{Z}/2(q)^{Bor} \rightarrow \mathbb{R}\epsilon_* \mathbb{Z}/2$, constructed in [HV12b, Proposition 5.5] induces a quasi-isomorphism $\mathbb{Z}/2(q)^{Bor} \rightarrow tr_{\leq 2q} \mathbb{R}\epsilon_* \mathbb{Z}/2$. The map ρ is its inverse in the derived category. Its hypercohomology gives the cycle map $H_G^n(X, \mathbb{Z}/2) \rightarrow H_{et}^n(X, \mathbb{Z}/2)$, for every $n \geq 0$. There is a quasi-isomorphism $\mathbb{R}\epsilon_* \mu_2^{\otimes q} \xrightarrow{\simeq} \mathbb{R}\epsilon_* \mu_2^{\otimes q+i}$ and a commutative diagram

$$(6.8) \quad \begin{array}{ccc} \mathbb{Z}/2(q)^{Bor} & \longrightarrow & tr_{\leq 2q} \mathbb{R}\epsilon_* \mu_2^{\otimes q} \\ \downarrow \simeq & & \downarrow \simeq \\ tr_{\leq 2q} \mathbb{Z}/2(q+i)^{Bor} & \longrightarrow & tr_{\leq 2q} \mathbb{R}\epsilon_* \mu_2^{\otimes q+i}. \end{array}$$

Take $i = q$. The result follows by showing the bottom map is a quasi-isomorphism. In [HV12b, Section 5] it is shown that the composite

$$\mathbb{Z}/2(2q) \rightarrow tr_{\leq 2q} \mathbb{Z}/2(2q)^{Bor} \rightarrow tr_{\leq 2q} \mathbb{R}\epsilon_* \mu_2^{\otimes 2q}$$

is the usual cycle map $\mathbb{Z}/2(2q) \rightarrow tr_{\leq 2q} \mathbb{R}\epsilon_* \mu_2^{\otimes 2q}$. By Voevodsky’s resolution of the Milnor conjecture [Voe03] this cycle map is a quasi-isomorphism. This implies that $H_G^n(X, \mathbb{Z}/2) \rightarrow H_{et}^n(X, \mathbb{Z}/2)$ is a surjective map between finite-dimensional spaces for $n \leq 2q$. By [Cox79] both vector spaces have the same dimension and so the map is an isomorphism. Therefore $\mathbb{Z}/2(q)^{Bor} \simeq tr_{\leq 2q} \mathbb{R}\epsilon_* \mathbb{Z}/2$. □

Let a_{Zar} denote Zariski sheafification and define the following sheaves

$$\begin{aligned} \mathcal{H}_{\mathcal{M}}^n(q) &= a_{Zar}(U \mapsto H_{\mathcal{M}}^n(U(\mathbb{C}), \mathbb{Z}/2(q))) \\ \mathcal{H}_{\mathbb{C}}^n(G) &= a_{Zar}(U \mapsto H_G^n(U(\mathbb{C}), \mathbb{Z}/2)) \\ \mathcal{H}_{\mathbb{R}}^n(G) &= a_{Zar}(U \mapsto H_G^n(U(\mathbb{R}), \mathbb{Z}/2)), \\ \mathcal{H}_{et}^n(q) &= a_{Zar}(U \mapsto H_{et}^n(U, \mu_2^{\otimes q})), \\ \mathcal{H}_{\mathbb{R}}^n &= a_{Zar}(U \mapsto H^n(U(\mathbb{R}), \mathbb{Z}/2)). \end{aligned}$$

Sheafifying the diagram in Theorem 6.3 for $p = q$ gives the commutative diagram

$$\begin{array}{ccccc} \mathcal{H}_{\mathcal{M}}^q(q) & \xrightarrow{\cong} & \mathcal{H}_{et}^q(q) & \xrightarrow{\cong} & \mathcal{H}_{\mathbb{C}}^q(G) \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{R}\mathcal{L}^q \mathcal{H}^0 & \xrightarrow{cyc_{q,0}} & \mathcal{H}_{\mathbb{R}}^0 & \longleftarrow & \mathcal{H}_{\mathbb{R}}^q(G). \end{array}$$

By Corollary 4.4 and Corollary 4.2 the sheafified cycle map $cyc_{q,0}$ induces the usual cycle map between reduced morphic cohomology and singular cohomology

$$RL^q H^n(X) = H^n_{Zar}(X; \mathcal{R}\mathcal{L}^q \mathcal{H}^0) \rightarrow H^n_{Zar}(X; \mathcal{H}^0_{\mathbb{R}}) = H^n(X(\mathbb{R}); \mathbb{Z}/2).$$

The map $\mathcal{H}^q_{\mathcal{M}}(q) \rightarrow \mathcal{H}^q_{et}(q)$ induces the Bloch-Ogus isomorphism

$$CH^q(X) = H^q_{Zar}(X; \mathcal{H}^q_{\mathcal{M}}(q)) \cong H^q_{Zar}(X; \mathcal{H}^q_{et}(q)).$$

The composite $\mathcal{H}^q_{et}(q) \rightarrow \mathcal{H}^q_{\mathbb{C}}(G) \rightarrow \mathcal{H}^q_{\mathbb{R}}(G) \rightarrow \mathcal{H}^0_{\mathbb{R}}$ induces a map

$$CH^q(X) = \mathcal{H}^q_{Zar}(X; \mathcal{H}^q_{et}(q)) \rightarrow \mathcal{H}^q_{Zar}(X; \mathcal{H}^0_{\mathbb{R}}) = H^q(X(\mathbb{R}); \mathbb{Z}/2)$$

which by [CTS96, Remark 2.3.5] is just the Borel-Haeflinger cycle map sending a closed irreducible $Z \subseteq X$ to the Poincare dual of the fundamental class of Z if $\dim Z(\mathbb{R}) = \dim Z$ and zero otherwise. The above commutative diagram tells us that this agrees with the composite $\mathcal{H}^q(q)_{et} \cong \mathcal{H}^q_{\mathcal{M}}(q) \rightarrow \mathcal{L}\mathcal{R}^q \mathcal{H}^0 \rightarrow \mathcal{H}^0_{\mathbb{R}}$ and so we immediately obtain the following.

THEOREM 6.9. *Let X be a smooth quasi-projective real variety. For any $q \geq 0$, the Borel-Haeflinger cycle map factors as the composite*

$$CH^q(X)/2 \rightarrow RL^q H^q(X) \xrightarrow{cyc_{q,n}} H^q(X(\mathbb{R}), \mathbb{Z}/2),$$

where the first map is the natural quotient.

Next we compare the s -map in reduced morphic cohomology with the operation (-1) in étale cohomology. Recall that the operation $(-1) : H^i_{et}(X; \mu_2^{\otimes q}) \rightarrow H^{i+1}_{et}(X; \mu_2^{\otimes q+1})$ is defined to be multiplication with the class (-1) which is the image of -1 under the boundary map $H^0_{et}(X; \mathbb{G}_m) \rightarrow H^1_{et}(X; \mu_2)$ in the Kummer sequence. By naturality this is equal to the pullback of $(-1) \in H^1_{et}(\text{Spec}(\mathbb{R}); \mu_2)$ under the structure map $X \rightarrow \text{Spec}(\mathbb{R})$. Sheafifying gives the operation on Zariski sheaves $(-1) : \mathcal{H}^q_{et}(q) \rightarrow \mathcal{H}^{q+1}_{et}(q+1)$.

PROPOSITION 6.10. *Let X be a smooth quasi-projective real variety of dimension d . The following square commutes for any $q \geq 0$*

$$\begin{array}{ccc} H^i_{Zar}(X; \mathcal{H}^q_{et}(q)) & \longrightarrow & RL^q H^i(X) \\ \downarrow \cup(-1) & & \downarrow \cup s \\ H^i_{Zar}(X; \mathcal{H}^{q+1}_{et}(q+1)) & \longrightarrow & RL^{q+1} H^i(X). \end{array}$$

For any $q \geq d + 1$ all maps are isomorphisms.

Proof. The s -operation is induced by multiplication with $s \in RL^1 H^0(\text{Spec}(\mathbb{R}))$, where s is the generator. Sheafifying the s -map gives a map of Zariski sheaves and the composite $\mathcal{R}\mathcal{L}^q \mathcal{H}^0 \rightarrow \mathcal{R}\mathcal{L}^{q+1} \mathcal{H}^0 \rightarrow \mathcal{H}^0_{\mathbb{R}}$ induces the usual s -map on sheaf cohomology.

The class $(-1) \in H^1_{et}(\text{Spec}(\mathbb{R}); \mu_2)$ and (-1) maps to s under the isomorphism $H^1_{et}(\text{Spec}(\mathbb{R}); \mu_2) \cong RL^1 H^0(\text{Spec}(\mathbb{R}))$ because they both map to the generator of $H^0(pt; \mathbb{Z}/2)$. Therefore the above square commutes. When $q \geq d$ then the vertical maps are isomorphisms, both $H^i_{Zar}(X; \mathcal{H}^q_{et}(q)) \rightarrow H^i(X(\mathbb{R}); \mathbb{Z}/2)$ and $RL^q H^i(X) \rightarrow H^i(X(\mathbb{R}); \mathbb{Z}/2)$ are isomorphisms. \square

COROLLARY 6.11. *Let X be a smooth quasi-projective real variety. We have the isomorphism of rings*

$$H_{\text{ét}}^*(X, \mu_2^{\otimes *})[s^{-1}] \cong RL^*H^*(X)[s^{-1}] \cong H^*(X(\mathbb{R}), \mathbb{Z}/2),$$

where $s = (-1)$ under the left-hand isomorphism.

REMARK 6.12. The map $\mathcal{H}_{\text{ét}}^q(q) \rightarrow \mathcal{R}\mathcal{L}^q H^0$ is not in general an isomorphism of sheaves as we can see for the case of a smooth projective variety of dimension $\dim(X) = q$. In this case the first group surjects with non-trivial kernel in codimension 0, 1, 2 (under some mild conditions on X) onto the cohomology of real points by [CTS96]. On the other hand the latter group is the cohomology of real points by [Teh10].

REFERENCES

- [AKMW02] Dan Abramovich, Kalle Karu, Kenji Matsuki, and Jarosław Włodarczyk, *Torification and factorization of birational maps*, J. Amer. Math. Soc. 15 (2002), no. 3, 531–572 (electronic). MR 1896232 (2003c:14016)
- [BCR98] Jacek Bochnak, Michel Coste, and Marie-Françoise Roy, *Real algebraic geometry*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 36, Springer-Verlag, Berlin, 1998, Translated from the 1987 French original, Revised by the authors. MR 1659509 (2000a:14067)
- [Bei10] A. Beilinson, *Remarks on Grothendieck’s standard conjectures*, ArXiv e-prints (2010).
- [BH61] Armand Borel and André Haefliger, *La classe d’homologie fondamentale d’un espace analytique*, Bull. Soc. Math. France 89 (1961), 461–513. MR 0149503 (26 #6990)
- [BO74] Spencer Bloch and Arthur Ogus, *Gersten’s conjecture and the homology of schemes*, Ann. Sci. École Norm. Sup. (4) 7 (1974), 181–201 (1975). MR MR0412191 (54 #318)
- [Cox79] David A. Cox, *The étale homotopy type of varieties over \mathbf{R}* , Proc. Amer. Math. Soc. 76 (1979), no. 1, 17–22. MR MR534381 (80f:14009)
- [CTHK97] Jean-Louis Colliot-Thélène, Raymond T. Hoobler, and Bruno Kahn, *The Bloch-Ogus-Gabber theorem*, Algebraic K-theory (Toronto, ON, 1996), Fields Inst. Commun., vol. 16, Amer. Math. Soc., Providence, RI, 1997, pp. 31–94. MR MR1466971 (98j:14021)
- [CTP90] J.-L. Colliot-Thélène and R. Parimala, *Real components of algebraic varieties and étale cohomology*, Invent. Math. 101 (1990), no. 1, 81–99. MR 1055712 (91j:14015)
- [CTS96] J.-L. Colliot-Thélène and C. Scheiderer, *Zero-cycles and cohomology on real algebraic varieties*, Topology 35 (1996), no. 2, 533–559. MR 1380515 (97a:14009)

- [dS03] Pedro F. dos Santos, *Algebraic cycles on real varieties and $\mathbb{Z}/2$ -equivariant homotopy theory*, Proc. London Math. Soc. (3) 86 (2003), no. 2, 513–544. MR MR1971161 (2004c:55026)
- [FG93] Eric M. Friedlander and Ofer Gabber, *Cycle spaces and intersection theory*, Topological methods in modern mathematics (Stony Brook, NY, 1991), Publish or Perish, Houston, TX, 1993, pp. 325–370. MR MR1215970 (94j:14010)
- [FHW04] Eric M. Friedlander, Christian Haesemeyer, and Mark E. Walker, *Techniques, computations, and conjectures for semi-topological K -theory*, Math. Ann. 330 (2004), no. 4, 759–807. MR MR2102312
- [FM94] Eric M. Friedlander and Barry Mazur, *Filtrations on the homology of algebraic varieties*, Mem. Amer. Math. Soc. 110 (1994), no. 529, x+110, With an appendix by Daniel Quillen. MR MR1211371 (95a:14023)
- [Fri95] Eric M. Friedlander, *Filtrations on algebraic cycles and homology*, Ann. Sci. École Norm. Sup. (4) 28 (1995), no. 3, 317–343. MR 1326671 (96i:14004)
- [Fri98] ———, *Algebraic cocycles on normal, quasi-projective varieties*, Compositio Math. 110 (1998), no. 2, 127–162. MR MR1602068 (2000a:14024)
- [FW02] Eric M. Friedlander and Mark E. Walker, *Semi-topological K -theory of real varieties*, Algebra, arithmetic and geometry, Part I, II (Mumbai, 2000), Tata Inst. Fund. Res. Stud. Math., vol. 16, Tata Inst. Fund. Res., Bombay, 2002, pp. 219–326. MR MR1940670 (2003h:19005)
- [HV12a] Jeremiah Heller and Mircea Voineagu, *Equivariant semi-topological invariants, Atiyah’s KR -theory, and real algebraic cycles*, Trans. Amer. Math. Soc. 364 (2012), no. 12, 6565–6603. MR 2958948
- [HV12b] ———, *Vanishing theorems for real algebraic cycles*, Amer. J. Math. 134 (2012), no. 3, 649–709.
- [IS88] Friedrich Ischebeck and Heinz-Werner Schülting, *Rational and homological equivalence for real cycles*, Invent. Math. 94 (1988), no. 2, 307–316. MR 958834 (89k:14004)
- [LF92] Paulo Lima-Filho, *Lawson homology for quasiprojective varieties*, Compositio Math. 84 (1992), no. 1, 1–23. MR MR1183559 (93j:14007)
- [Sch85] Heinz-Werner Schülting, *Algebraische und topologische reelle Zykeln unter birationalen Transformationen*, Math. Ann. 272 (1985), no. 3, 441–448. MR 799672 (87g:14018)
- [Sch94] Claus Scheiderer, *Real and étale cohomology*, Lecture Notes in Mathematics, vol. 1588, Springer-Verlag, Berlin, 1994. MR 1321819 (96c:14018)
- [Teh10] Jyh-Haur Teh, *A homology and cohomology theory for real projective varieties*, Indiana Univ. Math. J. 59 (2010), no. 1, 327–384. MR 2666482 (2011g:14018)

- [Voe03] Vladimir Voevodsky, *Motivic cohomology with $\mathbf{Z}/2$ -coefficients*, Publ. Math. Inst. Hautes Études Sci. (2003), no. 98, 59–104. MR MR2031199 (2005b:14038b)
- [Voi] M. Voineagu, *Cylindrical Homomorphisms and Lawson Homology*, Journal of K-theory. To appear. Preprint available at <http://arxiv.org/abs/0904.3374>.

Jeremiah Heller
Bergische Universität Wuppertal
Gaußstr. 20
D-42119 Wuppertal
Germany
heller@math.uni-wuppertal.de

Mircea Voineagu
IPMU
The University of Tokyo
5-1-5 Kashiwanoha
Kashiwa 277-8583
Japan
mircea.voineagu@ipmu.jp

